

**MULTI – CRITERIA GROUP DECISION MAKING MODEL IN NEUTROSOPHIC REFINED SET AND ITS APPLICATION****Surapati Pramanik\*, Durga Banerjee, B. C. Giri**

\* Department of Mathematics, Nandalal Ghosh B.T. College, Panpur, P.O.-Narayanpur, District –North 24 Parganas, West Bengal, India-743126

Ranaghat Yusuf Institution, P. O. Ranaghat, Dist. Nadia, India-741201

Department of Mathematics, Jadavpur University, Jadavpur, West Bengal, India-700032

DOI: 10.5281/zenodo.55307

**KEYWORDS:** Neutrosophic set, Single valued neutrosophic set, Neutrosophic refined sets, Tangent similarity measure, Multi criteria decision making.**ABSTRACT**

In the paper, multi criteria group decision making model has been presented based on tangent similarity measure of neutrosophic refined set. Simplified form of tangent similarity measure in neutrosophic refined set has been presented. New ranking method has been proposed based on refined tangent similarity measure. The proposed approach has been illustrated by solving a teacher selection problem in neutrosophic refined set environment.

**INTRODUCTION**

The term “neutrosophy” has been coined by Smarandache [1] in his study of new branch of philosophy. According to Smarandache neutrosophy means knowledge of neutral thoughts. From neutrosophy, neutrosophic set has been derived [1] which is the generalization of fuzzy set [2] and intuitionistic fuzzy set [3]. Wang et al. [4] studied single valued neutrosophic sets to deal realistic problems. Neutrosophic set and single valued neutrosophic sets have been applied in different areas of research such as medical diagnosis [5], decision making [6, 7], social problems [8, 9], conflict resolution [10], etc.

Similarity measure is a very important tool to determine the degree of similarity between two objects. Many researchers have studied different similarity measures and their properties. Broumi and Smarandache [11] studied Hausdroff distance between neutrosophic sets. Broumi and Smarandache [12] also studied correlation coefficient between interval neutrosophic sets. Broumi and Smarandache [13] also proposed cosine similarity measure and its properties based on Bhattacharjee’s distance [14]. Using distances, a matching function, membership grades Majumder and Samanta [15] established different similarity measures of single valued neutrosophic sets (SVNS) and proposed an entropy measure for SVNS. Ye and Zhang [16] described similarity measures using minimum and maximum operators and proposed a multi criteria decision making method based on the weighted similarity measure for SVNS. Ye [17] presented clustering method for single valued neutrosophic sets and a clustering algorithm for single valued neutrosophic data. Multi criteria group decision making approach under simplified neutrosophic environment has been proposed by Mondal and Pramanik [18]. They have used hybrid score accuracy function to make decision and apply their approach in teacher selection in higher education.

Biswas et al. [19] studied cosine similarity measure based on multi attribute decision making with trapezoidal fuzzy neutrosophic numbers. Pramanik and Mondal [20] applied cosine similarity measure of rough neutrosophic set in medical diagnosis. Decision making based on some similarity measures under interval rough neutrosophic environment has been studied by Mondal and Pramanik [21]. Mondal and Pramanik [22] further developed some rough neutrosophic similarity measures and their application to multi attribute decision making.

Yager [23] introduced the concept of multiset. Sebastian and Ramakrishnan [24] developed the concept and properties of multi fuzzy set. This concept has been extended to intuitionistic fuzzy multiset by Shinoj and John [25]. Smarandache [26] proposed n- valued refined neutrosophic logic and its application. Broumi and Smarandache [27] defined neutrosophic refined similarity measure based on cosine function. Dice similarity



measure between single valued neutrosophic multi sets and its application have been established by Ye and Ye [28].

Pramanik and Mondal [29] proposed weighted fuzzy similarity measure based on tangent function and applied it to medical diagnosis. Pramanik and Mondal [30] further studied tangent similarity measure in neutrosophic environments. Mondal and Pramanik [31] discussed neutrosophic refined similarity measure based on tangent function. Mondal and Pramanik [32] also proposed refined cotangent similarity measure and its application in multi attribute decision making.

We have presented simplified form of tangent similarity measure. We have developed multi criteria group decision making model based on tangent similarity measure of neutrosophic refined set.

Rest of the paper has been designed as follows: Some relevant definitions are presented in the section 2. Tangent similarity measure for neutrosophic refined sets and some of its properties have been stated in the next section. Section 4 is devoted to present decision making model based on refined tangent similarity measure. The application of refined tangent similarity measure to the problem on teacher section is shown in the next section. Finally, section 6 presents the conclusion and future works of the proposed approach.

## SOME RELEVANT DEFINITIONS

### Neutrosophic Set [1]

Let  $X$  be an universe of discourse. Then the neutrosophic set  $P$  is of the form  $P = \{ \langle x: T_P(x), I_P(x), F_P(x) \rangle | x \in X \}$ , where the functions  $T, I, F: X \rightarrow ]-0, 1+[$  are defined respectively the degree of membership, the degree of indeterminacy, and the degree of non-membership of the element  $x \in X$  to the set  $N$  satisfying the following condition.

$$-0 \leq \sup T_P(x) + \sup I_P(x) + \sup F_P(x) \leq 3^+$$

For two neutrosophic sets,  $P = \{ \langle x: T_P(x), I_P(x), F_P(x) \rangle | x \in X \}$  and  $Q = \{ \langle x, T_Q(x), I_Q(x), F_Q(x) \rangle | x \in X \}$  the two relations are defined as follows:

- (1)  $P \subseteq Q$  if and only if  $T_P(x) \leq T_Q(x), I_P(x) \geq I_Q(x), F_P(x) \geq F_Q(x)$
- (2)  $P = Q$  if and only if  $T_P(x) = T_Q(x), I_P(x) = I_Q(x), F_P(x) = F_Q(x)$

### Single valued neutrosophic set [4]

Let  $X$  be a space of points with generic elements in  $X$  denoted by  $x$ . A SVNS  $P$  in  $X$  is characterized by a truth-membership function  $T_P(x)$ , an indeterminacy-membership function  $I_P(x)$ , and a falsity membership function  $F_P(x)$ , for each point  $x$  in  $X$ ,  $T_P(x), I_P(x), F_P(x) \in [0, 1]$ . When  $X$  is continuous, a SVNS  $P$  can be written as:

$$P = \int_x \frac{\langle T_P(x), I_P(x), F_P(x) \rangle}{x} : x \in X$$

When  $X$  is discrete, a SVNS  $P$  can be written as:

$$P = \sum_{i=1}^n \frac{\langle T_P(x_i), I_P(x_i), F_P(x_i) \rangle}{x_i} : x_i \in X$$

For two SVNSs,  $P_{SVNS} = \{ \langle x: T_P(x), I_P(x), F_P(x) \rangle | x \in X \}$  and  $Q_{SVNS} = \{ \langle x, T_Q(x), I_Q(x), F_Q(x) \rangle | x \in X \}$  the two relations are defined as follows:

- (1)  $P_{SVNS} \subseteq Q_{SVNS}$  if and only if  $T_P(x) \leq T_Q(x), I_P(x) \geq I_Q(x), F_P(x) \geq F_Q(x)$
- (2)  $P_{SVNS} = Q_{SVNS}$  if and only if  $T_P(x) = T_Q(x), I_P(x) = I_Q(x), F_P(x) = F_Q(x)$  for any  $x \in X$

### Neutrosophic refined set [29]

Let  $P$  be a neutrosophic refined set (NRS)



## Global Journal of Engineering Science and Research Management

$P = \{ \langle x, (T_P^1(x_i), T_P^2(x_i), \dots, T_P^r(x_i)), (I_P^1(x_i), I_P^2(x_i), \dots, I_P^r(x_i)), (F_P^1(x_i), F_P^2(x_i), \dots, F_P^r(x_i))) \rangle : x \in X \}$  where,  $T_P^1(x_i), T_P^2(x_i), \dots, T_P^r(x_i) \in [0, 1]$ ,  $I_P^1(x_i), I_P^2(x_i), \dots, I_P^r(x_i) \in [0, 1]$  and  $F_P^1(x_i), F_P^2(x_i), \dots, F_P^r(x_i) \in [0, 1]$  such that  $0 \leq \sup T_P^j(x_i) + \sup I_P^j(x_i) + \sup F_P^j(x_i) \leq 3$ , for  $j = 1, 2, \dots, r$  for any  $x \in X$ . Now,  $(T_P^1(x_i), T_P^2(x_i), \dots, T_P^r(x_i)), (I_P^1(x_i), I_P^2(x_i), \dots, I_P^r(x_i)), (F_P^1(x_i), F_P^2(x_i), \dots, F_P^r(x_i)))$  are respectively called the truth-membership sequence, indeterminacy-membership sequence and falsity-membership sequence of the element  $x$ . Also,  $r$  is called the dimension of neutrosophic refined sets  $P$ .

### TANGENT SIMILARITY MEASURE FOR SINGLE VALUED REFINED NEUTROSOPHIC SETS [31]

Let  $P$  and  $Q$  be two neutrosophic refined sets where

$P = \{ \langle x, (T_P^1(x_i), T_P^2(x_i), \dots, T_P^r(x_i)), (I_P^1(x_i), I_P^2(x_i), \dots, I_P^r(x_i)), (F_P^1(x_i), F_P^2(x_i), \dots, F_P^r(x_i))) \rangle : x \in X \}$  and  $T_P^1(x_i), T_P^2(x_i), \dots, T_P^r(x_i) \in [0, 1]$ ,  $I_P^1(x_i), I_P^2(x_i), \dots, I_P^r(x_i) \in [0, 1]$  and  $F_P^1(x_i), F_P^2(x_i), \dots, F_P^r(x_i) \in [0, 1]$  such that  $0 \leq \sup T_P^j(x_i) + \sup I_P^j(x_i) + \sup F_P^j(x_i) \leq 3$  and  $Q = \{ \langle x, (T_Q^1(x_i), T_Q^2(x_i), \dots, T_Q^r(x_i)), (I_Q^1(x_i), I_Q^2(x_i), \dots, I_Q^r(x_i)), (F_Q^1(x_i), F_Q^2(x_i), \dots, F_Q^r(x_i))) \rangle : x \in X \}$  and  $(T_Q^1(x_i), T_Q^2(x_i), \dots, T_Q^r(x_i)) \in [0, 1]$ ,  $(I_Q^1(x_i), I_Q^2(x_i), \dots, I_Q^r(x_i)) \in [0, 1]$  and  $(F_Q^1(x_i), F_Q^2(x_i), \dots, F_Q^r(x_i)) \in [0, 1]$  such that  $0 \leq \sup T_Q^j(x_i) + \sup I_Q^j(x_i) + \sup F_Q^j(x_i) \leq 3$ , for  $j = 1, 2, \dots, r$ , for any  $x \in X$ . Now refined tangent similarity function between  $P$  and  $Q$  can be presented as:

$T_{NRS}(P, Q) =$

$$\frac{1}{r} \sum_{i=1}^r \left[ \frac{1}{n} \sum_{j=1}^n \left\langle 1 - \tan \left( \frac{\pi}{12} \left( \left| T_P^j(x_i) - T_Q^j(x_i) \right| + \left| I_P^j(x_i) - I_Q^j(x_i) \right| + \left| F_P^j(x_i) - F_Q^j(x_i) \right| \right) \right) \right\rangle \right] \quad (1)$$

In the paper, we redefine the above in more simplified form as

$$T_{NRS}(P, Q) = 1 - \frac{1}{nr} \sum_{j=1}^r \sum_{i=1}^n \tan \left( \frac{\pi}{12} \left( \left| T_P^j(x_i) - T_Q^j(x_i) \right| + \left| I_P^j(x_i) - I_Q^j(x_i) \right| + \left| F_P^j(x_i) - F_Q^j(x_i) \right| \right) \right) \quad (2)$$

#### Proposition 3.1

The defined refined tangent similarity measure  $T_{NRS}(P, Q)$  between NRSs  $P$  and  $Q$  satisfies the following properties:

1.  $0 \leq T_{NRS}(P, Q) \leq 1$
2.  $T_{NRS}(P, Q) = 1$  iff  $P = Q$
3.  $T_{NRS}(Q, P) = T_{NRS}(P, Q)$
4. If  $R$  is a NRS in  $X$  and  $P \subset Q \subset R$  then

$T_{NRS}(P, R) \leq T_{NRS}(P, Q)$  and  $T_{NRS}(P, R) \leq T_{NRS}(Q, R)$

For proofs of the propositions in 3.1, see the paper [31].

### DECISION MAKING UNDER SINGLE VALUED REFINED NEUTROSOPHIC SET BASED ON TANGENT SIMILARITY MEASURE

Let  $E_1, E_2, \dots, E_m$  be a discrete set of experts,  $A_1, A_2, \dots, A_n$  be the set of attributes and  $C_1, C_2, \dots, C_k$  be the set of candidates. The step by step process of decision making under refined neutrosophic tangent similarity measure are presented as follows.

#### Step 1: Determination the relation between experts ( $E_i$ ) and attributes ( $A_j$ )

Each expert  $E_i$  ( $i = 1, 2, \dots, m$ ) with the attribute  $A_j$  ( $j = 1, 2, \dots, n$ ) is presented in the tale 1 as follows.



**Table 1: Relation between experts and attributes in terms of NRSs**

	$A_1$	$A_2$	...	$A_n$
$E_1$	$\langle T_{11}^1, I_{11}^1, F_{11}^1 \rangle$	$\langle T_{12}^1, I_{12}^1, F_{12}^1 \rangle$	...	$\langle T_{1n}^1, I_{1n}^1, F_{1n}^1 \rangle$
	$\langle T_{11}^2, I_{11}^2, F_{11}^2 \rangle$	$\langle T_{12}^2, I_{12}^2, F_{12}^2 \rangle$	...	$\langle T_{1n}^2, I_{1n}^2, F_{1n}^2 \rangle$
	$\langle T_{11}^r, I_{11}^r, F_{11}^r \rangle$	$\langle T_{12}^r, I_{12}^r, F_{12}^r \rangle$	...	$\langle T_{1n}^r, I_{1n}^r, F_{1n}^r \rangle$
$E_2$	$\langle T_{21}^1, I_{21}^1, F_{21}^1 \rangle$	$\langle T_{22}^1, I_{22}^1, F_{22}^1 \rangle$	...	$\langle T_{2n}^1, I_{2n}^1, F_{2n}^1 \rangle$
	$\langle T_{21}^2, I_{21}^2, F_{21}^2 \rangle$	$\langle T_{22}^2, I_{22}^2, F_{22}^2 \rangle$	...	$\langle T_{2n}^2, I_{2n}^2, F_{2n}^2 \rangle$
	$\langle T_{21}^r, I_{21}^r, F_{21}^r \rangle$	$\langle T_{22}^r, I_{22}^r, F_{22}^r \rangle$	...	$\langle T_{2n}^r, I_{2n}^r, F_{2n}^r \rangle$
...	...	...	...	...
$E_m$	$\langle T_{m1}^1, I_{m1}^1, F_{m1}^1 \rangle$	$\langle T_{m2}^1, I_{m2}^1, F_{m2}^1 \rangle$	...	$\langle T_{mn}^1, I_{mn}^1, F_{mn}^1 \rangle$
	$\langle T_{m1}^2, I_{m1}^2, F_{m1}^2 \rangle$	$\langle T_{m2}^2, I_{m2}^2, F_{m2}^2 \rangle$	...	$\langle T_{mn}^2, I_{mn}^2, F_{mn}^2 \rangle$
	$\langle T_{m1}^r, I_{m1}^r, F_{m1}^r \rangle$	$\langle T_{m2}^r, I_{m2}^r, F_{m2}^r \rangle$	...	$\langle T_{mn}^r, I_{mn}^r, F_{mn}^r \rangle$

**Step 2: Determination the relation between attributes ( $A_j$ ) and candidates ( $C_t$ )**

The relation between attributes  $A_j$  ( $j = 1, 2, \dots, n$ ) and candidates (alternatives)  $C_t$  ( $t = 1, 2, \dots, k$ ) in terms of SVNS is presented in the table 2.

**Table 2: The relation between attributes and candidates in terms of SVNS**

	$C_1$	$C_2$	...	$C_k$
$A_1$	$\langle T_{11}, I_{11}, F_{11} \rangle$	$\langle T_{12}, I_{12}, F_{12} \rangle$	...	$\langle T_{1k}, I_{1k}, F_{1k} \rangle$
$A_2$	$\langle T_{21}, I_{21}, F_{21} \rangle$	$\langle T_{22}, I_{22}, F_{22} \rangle$	...	$\langle T_{2k}, I_{2k}, F_{2k} \rangle$
...	...	...	...	...
$A_n$	$\langle T_{n1}, I_{n1}, F_{n1} \rangle$	$\langle T_{n2}, I_{n2}, F_{n2} \rangle$	...	$\langle T_{nk}, I_{nk}, F_{nk} \rangle$

**Step 3: Determination of the relation between experts and candidates**

Using equation (2), the tangent similarity measure between the table 1 and the table 2 is presented in table 3.

Table 3: The tangent similarity measure between experts and candidates

	$C_1$	$C_2$	...	$C_k$
$E_1$	$\alpha_{11}$	$\alpha_{12}$	...	$\alpha_{1k}$
$E_2$	$\alpha_{21}$	$\alpha_{22}$	...	$\alpha_{2k}$
...	...	...	...	...
$E_m$	$\alpha_{m1}$	$\alpha_{m2}$	...	$\alpha_{mk}$

The average of each column is presented as follows:

$$\Omega_t = \frac{\sum_{i=1}^m \alpha_{it}}{m} \text{ where } t = 1, 2, \dots, k.$$



**Step 4: Ranking of the candidates**

Ranking of candidates is prepared based on the descending order average of each column  $\Omega_i$  of the table 3.

**Step 5:** End.

**EXAMPLE ON TEACHER SELECTION PROCESS**

Suppose that a university is going to recruit an assistant professor in mathematics. A selection board has been formed consisting of four experts ( $E_1, E_2, E_3, E_4$ ). After primary selection five candidates ( $C_1, C_2, C_3, C_4, C_5$ ) appear before the board. Six attributes (criteria) (demonstration ( $A_1$ ), pedagogical knowledge ( $A_2$ ), action research ( $A_3$ ), emotional stability ( $A_4$ ), and knowledge of child psychology ( $A_5$ ), social quality ( $A_6$ )) obtained from experts' opinions are the parameters for selection. Selection process is divided into three phases represented by neutrosophic refined sets. The relation between experts and attributes has been presented in the table 4.

**Table 4: The relation between experts and criteria**

	Demonstration ( $A_1$ )	Pedagogical knowledge ( $A_2$ )	Action Research ( $A_3$ )	Emotional stability ( $A_4$ )	Knowledge of child psychology ( $A_5$ )	Social quality ( $A_6$ )
$E_1$	(0.8, 0.1, 0.1) (0.6, 0.2, 0.3) (0.5, 0.4, 0.2)	(0.7, 0.2, 0.1) (0.7, 0.4, 0.3) (0.8, 0.1, 0.1)	(0.6, 0.1, 0.3) (0.7, 0.1, 0.4) (0.7, 0.1, 0.1)	(0.7, 0.1, 0.2) (0.5, 0.2, 0.1) (0.6, 0.1, 0.1)	(0.6, 0.4, 0.1) (0.8, 0.2, 0.2) (0.7, 0.2, 0.3)	(0.3, 0.5, 0.4) (0.1, 0.6, 0.1) (0.1, 0.5, 0.2)
$E_2$	(0.8, 0.2, 0.1) (0.5, 0.3, 0.3) (0.6, 0.1, 0.3)	(0.9, 0.1, 0.1) (0.6, 0.4, 0.2) (0.8, 0.2, 0.1)	(0.3, 0.4, 0.2) (0.4, 0.2, 0.2) (0.3, 0.3, 0.3)	(0.3, 0.5, 0.4) (0.4, 0.4, 0.4) (0.4, 0.3, 0.2)	(0.3, 0.3, 0.3) (0.4, 0.2, 0.1) (0.5, 0.2, 0.1)	(0.1, 0.6, 0.1) (0.5, 0.2, 0.4) (0.6, 0.4, 0.1)
$E_3$	(0.7, 0.1, 0.2) (0.4, 0.3, 0.5) (0.5, 0.3, 0.3)	(0.4, 0.1, 0.2) (0.6, 0.1, 0.1) (0.6, 0.3, 0.2)	(0.5, 0.4, 0.1) (0.4, 0.4, 0.1) (0.6, 0.3, 0.2)	(0.5, 0.1, 0.1) (0.4, 0.2, 0.2) (0.4, 0.1, 0.1)	(0.4, 0.1, 0.3) (0.3, 0.2, 0.5) (0.4, 0.1, 0.2)	(0.3, 0.1, 0.1) (0.1, 0.2, 0.3) (0.2, 0.1, 0.1)
$E_4$	(0.6, 0.3, 0.1) (0.7, 0.1, 0.1) (0.8, 0.1, 0.1)	(0.5, 0.0, 0.1) (0.2, 0.2, 0.1) (0.3, 0.1, 0.2)	(0.4, 0.2, 0.1) (0.3, 0.1, 0.1) (0.4, 0.2, 0.2)	(0.4, 0.4, 0.2) (0.4, 0.1, 0.2) (0.5, 0.2, 0.1)	(0.6, 0.3, 0.2) (0.6, 0.2, 0.2) (0.5, 0.1, 0.1)	(0.2, 0.5, 0.4) (0.3, 0.4, 0.2) (0.3, 0.4, 0.1)

The relation between attributes  $A_j$  ( $j = 1, 2, \dots, n$ ) and candidates (alternatives)  $C_t$  ( $t = 1, 2, \dots, k$ ) in terms of SVNs is presented in the table 5.

**Table 5: The relation between criterion and candidates**

	$C_1$	$C_2$	$C_3$	$C_4$	$C_5$
( $A_1$ )	(0.6, 0.4, 0.1)	(0.7, 0.2, 0.3)	(0.8, 0.0, 0.1)	(0.4, 0.5, 0.2)	(0.6, 0.3, 0.4)
( $A_2$ )	(0.2, 0.1, 0.1)	(0.7, 0.3, 0.1)	(0.9, 0.2, 0.3)	(0.8, 0.2, 0.0)	(0.3, 0.3, 0.4)
( $A_3$ )	(0.4, 0.4, 0.4)	(0.8, 0.2, 0.2)	(0.6, 0.1, 0.4)	(0.7, 0.2, 0.1)	(0.7, 0.1, 0.3)
( $A_4$ )	(0.6, 0.6, 0.2)	(0.8, 0.2, 0.1)	(0.2, 0.1, 0.7)	(0.1, 0.6, 0.3)	(0.6, 0.3, 0.2)
( $A_5$ )	(0.6, 0.1, 0.2)	(0.7, 0.1, 0.2)	(0.5, 0.2, 0.1)	(0.5, 0.5, 0.2)	(0.8, 0.2, 0.0)
( $A_6$ )	(0.7, 0.2, 0.1)	(0.8, 0.2, 0.2)	(0.3, 0.2, 0.1)	(0.6, 0.3, 0.1)	(0.7, 0.0, 0.1)

Using equation (2), the tangent similarity measure between the table 4 and the table 5 is presented in table 6.

**Table 6: The tangent similarity measure between experts and candidates**

	$C_1$	$C_2$	$C_3$	$C_4$	$C_5$
$E_1$	0.9456	0.8851	0.8646	0.8403	0.8533
$E_2$	0.8635	0.8541	0.8252	0.8650	0.8347
$E_3$	0.8606	0.8665	0.8454	0.8322	0.8440



$E_4$	0.8840	0.8604	0.8601	0.8381	0.8468
$\Omega_t$	<b>0.8884</b>	<b>0.8665</b>	<b>0.8488</b>	<b>0.8439</b>	<b>0.8447</b>

The average of each column is presented by  $\Omega_t$  (see the table 6). The merit list on the basis of average has been shown in the table 7.

**Table 7: Merit List**

Rank	Candidates
1	B <sub>1</sub>
2	B <sub>2</sub>
3	B <sub>3</sub>
4	B <sub>5</sub>
5	B <sub>4</sub>

From the merit list, it is observed that B<sub>1</sub> is the most eligible candidate for the post.

## CONCLUSION

In the present study, we have developed multi criteria group decision making model in neutrosophic refined set and presented its application in teacher selection. The tangent similarity function has been presented in more simplified form. The ranking of the candidates has been prepared on the basis of decending order avarage of each column of tangent similarity measure table. The proposed approach can be also applied to other group decision making problems under refined neutrosophic set environment. The proposed approach can be also extended to neutrosophic hybrid envirnment.

## REFERENCES

1. F. Smarandache, A unifying field in logics: neutrosophic logic, neutrosophy, neutrosophic set, neutrosophic probability, and neutrosophic statistics, Rehoboth: American Research Press (1998).
2. L.A. Zadeh, Fuzzy sets, information and Control 8(3) (1965) 338-3534.
3. K. T. Atanassov, Intuitionistic fuzzy sets, Fuzzy Sets and Systems 20 (1986) 87-96.
4. H. Wang, F. Smarandache, Y. Q. Zhang, R. Sunderraman, Single valued neutrosophic sets, Multispace and Multi structure 4 (2010) 410-413.
5. J. Ye, Improved cosine similarity measures of simplified neutrosophic sets for medical diagnoses, Artificial Intelligence in Medicine 63(3) (2015) 171-179.
6. P. Biswas, S. Pramanik, B.C. Giri, TOPSIS method for multi-attribute group decision making under single-valued neutrosophic environment, Neural computing and Applications (2015). DOI: 10.1007/s00521-015-1891-2.
7. S. Pramanik, P. Biswas, B.C. Giri, Hybrid Vector similarity measures and their applications to multi-attribute decision making under neutrosophic environment, Neural Computing and Applications (2015), DOI 10.1007/s00521-015-2125-3.
8. S. Pramanik, S.N. Chackrabarti, A study on problems of construction workers in West Bengal based on neutrosophic cognitive maps, International Journal of Innovative Research in Science, Engineering and Technology 2(11) (2013) 6387-6394.
9. K. Mondal, S. Pramanik, A study on problems of Hijras in West Bengal based on neutrosophic cognitive maps, Neutrosophic Sets and Systems 5(2014) 21-26.
10. S. Pramanik, T.K. Roy, Neutrosophic game theoretic approach to Indo-Pak conflict over Jammu-Kashmir, Neutrosophic Sets and Systems 2 (2014) 82-101.
11. S. Broumi, F. Smarandache, Several similarity measures of neutrosophic sets, Neutrosophic Sets and Systems 1(2013) 54-62.
12. S. Broumi, F. Smarandache, Correlation coefficient of interval neutrosophic set, In Neutrosophic Theory and Its Applications, Collected Papers I, Bruxelles: EuropaNova (2014) 67-73.



## Global Journal of Engineering Science and Research Management

13. S. Broumi, F. Smarandache, Cosine similarity measure of interval valued neutrosophic sets, Neutrosophic Sets and Systems 5 (2014) 15-20.
14. A. Bhattacharya, On a measure of divergence of two multinomial population, Sankhya Series A 7 (1946) 401- 406.
15. P. Majumder, S. K. Samanta, On similarity and entropy of neutrosophic sets, Journal of Intelligent and Fuzzy Systems 26(2014) 1245–1252.
16. J. Ye, Q. S. Zhang, Single valued neutrosophic similarity measures for multiple attribute decision making, Neutrosophic Sets and Systems 2(2014) 48-54.
17. J. Ye, Clustering methods using distance-based similarity measures of single-valued neutrosophic sets, Journal of Intelligent Systems 23(4) (2014) 379-389.
18. K. Mondal, S. Pramanik, Multi-criteria decision making approach for teacher recruitment in higher education under simplified neutrosophic environment, Neutrosophic Sets and Systems 6 (2014) 28-34.
19. P. Biswas, S. Pramanik, B. C. Giri, Cosine similarity measure based multi-attribute decision-making with trapezoidal fuzzy neutrosophic numbers, Neutrosophic sets and Systems 8 (2015) 48-58.
20. S. Pramanik, K. Mondal, Cosine similarity measure of rough neutrosophic sets and its application in medical diagnosis, Global Journal of Advanced Research 2(1) (2015) 212-220.
21. K. Mondal, S. Pramanik, Decision making based on some similarity measures under interval rough neutrosophic environment, Neutrosophic Sets and Systems 10 (2015) 46-57.
22. S. Pramanik, K. Mondal, Some rough neutrosophic similarity measures and their application to multi attribute decision making, Global Journal of Engineering Science and Research Management 2 (7) (2015) 61-74.
23. R. R. Yager, On the theory of bags (multi sets), International Journal of General Systems 13(1986) 23-37.
24. S. Sebastian, T. V. Ramakrishnan, Multi fuzzy sets: an extension of fuzzy sets, Fuzzy Information Engine 3(1) (2011) 35-43.
25. T. K. Shinoj, S. J. John, Intuitionistic fuzzy multi-sets and its application in medical diagnosis, World Academy of Science, Engine Technology 61 (2012) 1178-1181.
26. F. Smarandache, n-Valued refined neutrosophic logic and its applications in physics, Progress in Physics 4 (2013) 143-146.
27. S. Broumi, F. Smarandache, Neutrosophic refined similarity measure based on cosine function, Neutrosophic Sets and Systems 6 (2014) 42-48.
28. S. Ye, J. Ye, Dice Similarity measure between single valued neutrosophic multisets and its application in medical diagnosis, Neutrosophic Sets and Systems 6 (2014) 50-55.
29. S. Pramanik, K. Mondal, **Weighted fuzzy similarity measure based on tangent function and its application to medical diagnosis, International Journal of Innovative Research in Science, Engineering Technology 4(2) (2015) 158-164.**
30. K. Mondal, S. Pramanik, Intuitionistic Fuzzy similarity measure based on tangent function and its application to multi-attribute decision making, Global Journal of Advanced Research 2(2) (2015) 464-471.
31. K. Mondal, S. Pramanik, Neutrosophic refined similarity measure based on tangent function and its application to multi attribute decision making, Journal of New theory 8 (2015) 41-50.
32. K. Mondal, S. Pramanik, Neutrosophic refined similarity measure based on cotangent function and its application to multi-attribute decision making, Global Journal of Advanced Research 2(2) (2015) 486-494.