



A Study of Neutrosophic Cubic Hesitant Fuzzy Hybrid Geometric Aggregation Operators and its Application to Multi Expert Decision Making System

¹Ateeq Ur Rehman, ¹Muhammad Gulistan*, ¹Zahid Khan, ^{2,3}Fuad S. Al-Duais

¹Department of Mathematics and Statistics, Hazara University, Mansehra 21130 Pakistan; ati.mathematics@gmail.com, gulistanmath@hu.edu.pk, gulistanm21@yahoo.com, zahidkhan@hu.edu.pk

²Department of Mathematics, College of Science and Humanities in Al-Aflaj, Prince Sattam bin Abdulaziz University, Al-Aflaj 11942, Saudi Arabia, f.alduais@psau.edu.sa

³Administration Department, Administrative Science College, Thamar University, Thamar, Yemen

*Correspondence: gulistanmath@hu.edu.pk, gulistanm21@yahoo.com

Abstract: Hesitancy is an imperative part of belief system. In order to counter the hesitancy in neutrosophic cubic set (NCS), the notion of neutrosophic cubic hesitant fuzzy set (NCHFS) is presented. NCHFS couple NCS with hesitant fuzzy set (HFS). Operational laws in NCHFS are developed with examples. To meet the challenges of decision making problems, neutrosophic cubic hesitant fuzzy geometric (NCHFG) aggregation operators, neutrosophic cubic hesitant fuzzy Einstein geometric (NCHFEG) aggregation operators, neutrosophic cubic hesitant fuzzy hybrid geometric (NCHFHEG) aggregation operators are developed in the current study. At the end a multi expert decision making (MEDM) process is proposed and furnished upon numerical data of a company as applications.

Keywords: Neutrosophic Cubic Fuzzy Hesitant Set (NCHFS), Neutrosophic Cubic Hesitant Fuzzy Weighted geometric (NCHFVG) operator, Neutrosophic Cubic Hesitant Fuzzy Einstein Geometric (NCHFEG) operator, Neutrosophic Cubic Hesitant Fuzzy Einstein Hybrid Geometric (NCHFHEG) Operator. multi expert decision making (MEDM)

1. Introduction

We are in different mental states of acceptance, hesitancy and refusal while taking decisions in life. Many methods in MADM ignore the uncertainty and hence yields the results which are unreliable. The role of expert in decision making (DM) is vital. The participation of more than one expert in a DM process reduce the uncertainty. Zadeh proposed the notion of fuzzy set (FS) [1] as a function from a given set of objects to $[0,1]$ called membership. Later Zadeh extended the idea to interval valued fuzzy set (IVFS) [2]. An IVFS a function from a given set of objects to the subintervals of $[0,1]$. The FS theory has many applications in artificial intelligence, robotics, computer networks, engineering and DM [3,4]. Different researchers [5-8] established similarity measures and other important concepts and successfully apply their models to medical diagnosis and selection criteria. R.A. Krohling and V.C. Campanharo, M. Xia and Z. Xu, M.K. Mehlawat and P.A. Gupta established different useful techniques to sort out MADM problems [9-11]. K. Atanassov introduced non-membership degree and presented the idea of intuitionistic fuzzy set (IFS) [12] which consist of both membership and non-membership degree within $[0,1]$. An extension of IFS was proposed and

named as interval value intuitionistic fuzzy set (IVIFS) [13]. IVIFS contains membership and non-membership in the form of subintervals of $[0,1]$. This characteristic of intuitionistic fuzzy set made it more applicable than previous versions and attracted researchers [14-16] to apply it to the fields of science, engineering and daily life problems. Jun et al., combined IVIFS and FS and proposed cubic set (CS). The CS is the generalization of IFS and IVIFS. CS become vital tool to deal the vague data. Several researchers explored algebraic aspects and apparently define ideal theory in CS [17-20]. F. Smarandache initiated the concept of indeterminacy and describe the notion of neutrosophic set (NS) [21]. NS consist of three components truth, indeterminacy and falsehood and all are independent. This characteristic of neutrosophic set enabled researchers to work with inconsistent and vague data more effectively. Wang *et al* proposed single valued neutrosophic set (SNVS) [22] by restricting components of NS to $[0,1]$. The NS was further extended to interval neutrosophic set (INS) [23]. After the appearance of NS, researchers put their contributions in theoretical as well as technological developments of the set [24-27]. Several researchers use neutrosophic and interval valued neutrosophic environments to construct MADM methods [28-32]. Zhan *et al.*, define aggregation operators and furnished some applications in MADM [33]. Torra define hesitant fuzzy set (HFS) [34] in contrast of FS. HFS on X is a function that maps every object of X into a subset of $[0, 1]$. Jun *et al.*, presented the notion of NCS [35] which consist of both INS and NS. These characteristics of NCS make it a powerful tool to deal the vague and inconsistent data more efficiently. Soon after its exploration it attracted the researcher to work in many fields like medicine, algebra, engineering and DM. Later the idea of cubic hesitant fuzzy set was introduced by Tahir *et al.*, [36]. Ye [37] establish similarity measure in neutrosophic hesitant fuzzy sets (NHFS) and established MADM method using these measures. Liu et. Al [38] proposed hybrid geometric aggregation operators in interval neutrosophic hesitant fuzzy sets (INHFS) and discuss its applications in MADM. Zhu et al. [39] proposed the method of β -normalization to enlarge a HFE, which is a useful technique in case of different cardinalities.

The remaining of the paper is formulated as follows. In section 2, we reviewed some basic definitions used later on. Section 3 deals with NCHFS, algebraic and Einstein operational laws in NCHFS. In section 4 we introduced aggregation operators in NCHFS. Section 5 concern with establishing a MEDM method based on NCHFG operators and use this method in MEDM problem.

2. Preliminaries

Definition 2.1: [1] A fuzzy set (FS) on a nonempty set W is a mapping $\Gamma: W \rightarrow [0,1]$.

Definition 2.2: [12] The cubic set (CS) on a nonempty set Z is defined by $\mu = \langle x; I(x), \delta(x)/x \in X \rangle$, where $I(x)$ is an IVIFS on Z and $\delta(x)$ is an FS on Z .

Definition 2.3: [22] A neutrosophic set associated with a crisp set S , is a set of the form $\mu = \langle e; \xi_T(e), \xi_I(e), \xi_F(e)/e \in S \rangle$ where $\xi_T, \xi_I, \xi_F: S \rightarrow [0,1]$ respectively called a truth membership function, a non-membership function and a false membership function.

Definition 2.4: [34] A hesitant fuzzy set on a crisp set W is a mapping which assigns a set of values in $[0,1]$, to each element of W .

Definition 2.5: [35] A neutrosophic cubic set in a nonempty set E is defined as a pair (B, μ) where $B = \langle x; B_T(e), B_I(e), B_F(e)/e \in E \rangle$ is an INS and $\mu = \langle x; \mu_T(e), \mu_I(e), \mu_F(e)/e \in X \rangle$ is a NS.

Definition 2.6: [39] A neutrosophic hesitant fuzzy set a nonempty set E is described as $\mu = \langle x; \mu_T(e), \mu_I(e), \mu_F(e)/e \in E \rangle$ where $\mu_T(e), \mu_I(e), \mu_F(e)$ are three HFSs such that $\mu_T(e) + \mu_I(e) + \mu_F(e) \leq 3$.

Definition2.7: [40] The object $\zeta = \langle x; \xi_T(x), \xi_I(x), \xi_F(x)/x \in X \rangle$, is called an INHFS on X , where $\xi_T(x), \xi_I(x), \xi_F(x)$ are IHFSs.

Zhu et al. proposed the following β -normalization method to enlarge a hesitant fuzzy element, which is a useful technique in case of different cardinalities.

Definition 2.8:[41] Let m^+ and m^- be the maximum and minimum elements of an hesitant fuzzy set H and $\zeta(0 \leq \zeta \leq 1)$ an optimized parameter. We call $m = \zeta m^+ + (1 - \zeta)m^-$ an added element.

3. NCHFS and operational Laws in NCHFS

Definition3.1: Let X be a nonempty set. A neutrosophic cubic hesitant fuzzy set in X is a pair $\alpha = \langle A, \lambda \rangle$ where $A = \langle x; A_T(x), A_I(x), A_F(x)/x \in X \rangle$ is an interval-valued neutrosophic hesitant set in X and $\lambda = \langle x; \lambda_T(x), \lambda_I(x), \lambda_F(x)/x \in X \rangle$ is a neutrosophic hesitant set in X . Furthermore $A_T = \{[A_{j_T}^L, A_{j_T}^U]; j = 1, \dots, l\}$, $A_I = \{[A_{j_I}^L, A_{j_I}^U]; j = 1, \dots, m\}$, $A_F = \{[A_{j_F}^L, A_{j_F}^U]; j = 1, \dots, n\}$ are some interval values in $[0,1]$ and $\lambda_T = \{\lambda_{j_T}; j = 1, \dots, r\}$, $\lambda_I = \{\lambda_{j_I}; j = 1, \dots, s\}$, $\lambda_F = \{\lambda_{j_F}; j = 1, \dots, t\}$ are some values in $[0,1]$.

Example 3.2: Let $X = \{u, v, w\}$ The pair $\alpha = \langle A, \lambda \rangle$ with

$$A_T(u) = \{[0.1, 0.5], [0.2, 0.7]\}, \lambda_T(u) = \{0.3, 0.5, 0.7\}, A_I(u) = \{[0.2, 0.4], [0.3, 0.6]\}, \lambda_I(u) = \{0.1, 0.4, 0.7\}, A_F(u) = \{[0.1, 0.4], [0.0, 0.3], [0.6, 0.8]\}, \lambda_F(u) = \{0.4, 0.6\}$$

$$A_T(v) = \{[0.1, 0.5], [0.2, 0.7]\}, \lambda_T(v) = \{0.3, 0.5\}, A_I(v) = \{[0.2, 0.3], [0.1, 0.6]\}, \lambda_I(v) = \{0.7, 0.8\}, A_F(v) = \{[0.1, 0.4], [0, 0.3]\}, \lambda_F(v) = \{0.4, 0.6\}$$

$$A_T(w) = \{[0.1, 0.5], [0.2, 0.7]\}, \lambda_T(w) = \{0.3, 0.5\}, A_I(w) = \{[0.2, 0.3], [0.1, 0.6]\}, \lambda_I(w) = \{0.7, 0.8\}, A_F(z) = \{[0.1, 0.4], [0, 0.3]\}, \lambda_F(w) = \{0.4, 0.6\}$$

is a NCHFS.

Definition 3.3: The sum of two NCHFSs $\alpha = \langle A, \lambda \rangle, \beta = \langle B, \mu \rangle$ is defined as

$$\alpha \oplus \beta = \left\langle x, \left\{ \left[A_{j_T}^L + B_{j_T}^L - A_{j_T}^L B_{j_T}^L, A_{j_T}^U + B_{j_T}^U - A_{j_T}^U B_{j_T}^U \right], \left[A_{j_I}^L + B_{j_I}^L - A_{j_I}^L B_{j_I}^L, A_{j_I}^U + B_{j_I}^U - A_{j_I}^U B_{j_I}^U \right], \left[A_{j_F}^L + B_{j_F}^L - A_{j_F}^L B_{j_F}^L, A_{j_F}^U + B_{j_F}^U - A_{j_F}^U B_{j_F}^U \right] \right\}, \left\{ \lambda_{j_T} + \mu_{j_T} - \lambda_{j_T} \mu_{j_T}, \lambda_{j_I} + \mu_{j_I} - \lambda_{j_I} \mu_{j_I}, \lambda_{j_F} \right\} \right\rangle,$$

Moreover the β -normalization is used in case of different cardinalities.

Example3.4:If

$$\alpha = \left\langle \{[0.1, 0.5], [0.2, 0.7]\}, \{[0.2, 0.3], [0.1, 0.6]\}, \{[0.1, 0.4], [0.0, 0.3]\}, \{0.1, 0.2\}, \{0.3, 0.5, 0.7\}, \{0.4, 0.8\} \right\rangle,$$

$$\text{and } \beta = \left\langle \{[0.4, 0.5], [0.3, 0.4]\}, \{[0.1, 0.3], [0.2, 0.5]\}, \{[0.1, 0.4], [0.7, 0.8]\}, \{0.3, 0.4, 0.5\}, \{0.7, 0.8\}, \{0.4, 0.6\} \right\rangle,$$

then using above definition and β -normalization with parameter $\xi = 0.5$ we have

$$\alpha \oplus \beta = \left\langle \{[0.46, 0.75], [0.44, 0.82]\}, \{[0.28, 0.51], [0.28, 0.8]\}, \{[0.01, 0.16], [0, 0.24]\}, \{0.03, 0.06, 0.1\}, \{0.21, 0.375, 0.56\}, \{0.64, 0.92\} \right\rangle.$$

Definition 3.5: The Product of two NCHFSs $\alpha = \langle A, \lambda \rangle, \beta = \langle B, \mu \rangle$ is defined by

$$\alpha \otimes \beta = \left\langle x, \left\{ \left[A_{j_T}^L B_{j_T}^L, A_{j_T}^U B_{j_T}^U \right], \left[A_{j_I}^L B_{j_I}^L, A_{j_I}^U B_{j_I}^U \right], \left[A_{j_F}^L + B_{j_F}^L - A_{j_F}^L B_{j_F}^L, A_{j_F}^U + B_{j_F}^U - A_{j_F}^U B_{j_F}^U \right] \right\}, \left\{ \lambda_{j_T} \mu_{j_T}, \lambda_{j_I} \mu_{j_I}, \lambda_{j_F} + \mu_{j_F} - \lambda_{j_F} \mu_{j_F} \right\} \right\rangle.$$

Moreover the β -normalization is used in case of different cardinalities.

Example3.6:If

$$\alpha = \left\langle \{[0.1, 0.5], [0.2, 0.7]\}, \{[0.2, 0.3], [0.1, 0.6]\}, \{[0.1, 0.4], [0.0, 0.3]\}, \{0.1, 0.2\}, \{0.3, 0.5, 0.7\}, \{0.4, 0.8\} \right\rangle, \text{ and}$$

$$\beta = \left\langle \{[0.4, 0.5], [0.3, 0.4]\}, \{[0.1, 0.3], [0.2, 0.5]\}, \{[0.1, 0.4], [0.7, 0.8]\}, \{0.3, 0.4, 0.5\}, \{0.7, 0.8\}, \{0.4, 0.6\} \right\rangle, \text{ then using}$$

above definition and β -normalization with parameter $\xi = 0.5$ we have

$$\alpha \otimes \beta = \left\langle \left\{ [0.04, 0.25], [0.06, 0.28] \right\}, \left\{ [0.02, 0.09], [0.02, 0.3] \right\}, \left\{ [0.19, 0.64], [0.7, 0.86] \right\}, \right. \\ \left. \{0.37, 0.49, 0.6\}, \{0.79, 0.875, 0.94\}, \{0.16, 0.48\} \right\rangle.$$

Definition 3.7: The scalar multiplication of a scalar q with a NCHFS $\alpha = \langle A, \lambda \rangle$ is defined by

$$q\alpha = \left\langle \left\{ \left[1 - \left(1 - A_{j_T}^L \right)^q, 1 - \left(1 - A_{j_T}^U \right)^q \right] \right\}, \left\{ \left[1 - \left(1 - A_{j_I}^L \right)^q, 1 - \left(1 - A_{j_I}^U \right)^q \right] \right\}, \left\{ \left[\left(A_{j_F}^L \right)^q, \left(A_{j_F}^U \right)^q \right] \right\}, \right. \\ \left. \left\{ 1 - \left(1 - \lambda_{j_T} \right)^q \right\}, \left\{ 1 - \left(1 - \lambda_{j_I} \right)^q \right\}, \left\{ \left(\lambda_{j_F} \right)^q \right\} \right\rangle.$$

Example3.8:If

$$\alpha = \left\langle \{[0.1, 0.5], [0.2, 0.7]\}, \{[0.2, 0.3], [0.1, 0.6]\}, \{[0.1, 0.4], [0.0, 0.3]\}, \{0.1, 0.2\}, \{0.3, 0.5\}, \{0.4, 0.8\} \right\rangle,$$

then using above definition with $q=3$ we have

$$3\alpha = \left\langle \left\{ [0.271, 0.875], [0.488, 0.973] \right\}, \left\{ [0.488, 0.657], [0.271, 0.936] \right\}, \left\{ [0.001, 0.64], [0.27] \right\}, \right. \\ \left. \{0.001, 0.008\}, \{0.27, 0.125\}, \{0.784, 0.992\} \right\rangle.$$

Definition 3.9: For NCHFS $\alpha = \langle A, \lambda \rangle$ and a scalar q

$$\alpha^q = \left\langle x, \left\{ \left[\left(A_{p_T}^L \right)^q, \left(A_{p_T}^U \right)^q \right] \right\}, \left\{ \left[\left(A_{p_I}^L \right)^q, \left(A_{p_I}^U \right)^q \right] \right\}, \left\{ \left[1 - \left(1 - A_{p_F}^L \right)^q, 1 - \left(1 - A_{p_F}^U \right)^q \right] \right\}, \right. \\ \left. \left\{ \left(\lambda_{p_T} \right)^q \right\}, \left\{ \left(\lambda_{p_I} \right)^q \right\}, \left\{ 1 - \left(1 - \lambda_{p_F} \right)^q \right\} \right\rangle$$

where $\alpha^q = \alpha \otimes \alpha \otimes \dots \otimes \alpha (q - \text{times})$ moreover α^q is a NCHF value for every $q > 0$.

Definition 3.10: The Einstein sum of two NCHFSs $\alpha = \langle A, \lambda \rangle, \beta = \langle B, \mu \rangle$ is defined by

$$\alpha \oplus_E \beta = \left\langle \left\{ \left[\frac{A_{j_T}^L + B_{j_T}^L}{1 + A_{j_T}^L B_{j_T}^L}, \frac{A_{j_T}^U + B_{j_T}^U}{1 + A_{j_T}^U B_{j_T}^U} \right] \right\}, \left\{ \left[\frac{A_{j_I}^L + B_{j_I}^L}{1 + A_{j_I}^L B_{j_I}^L}, \frac{A_{j_I}^U + B_{j_I}^U}{1 + A_{j_I}^U B_{j_I}^U} \right] \right\}, \left\{ \left[\frac{A_{j_F}^L B_{j_F}^L}{1 + (1 - A_{j_F}^L)(1 - B_{j_F}^L)}, \frac{A_{j_F}^U B_{j_F}^U}{1 + (1 - A_{j_F}^U)(1 - B_{j_F}^U)} \right] \right\}, \right. \\ \left. \left\{ \frac{\lambda_{j_T} + \mu_{j_T}}{1 + \lambda_{j_T} \mu_{j_T}} \right\}, \left\{ \frac{\lambda_{j_I} + \mu_{j_I}}{1 + \lambda_{j_I} \mu_{j_I}} \right\}, \left\{ \frac{\lambda_{j_F} \mu_{j_F}}{1 + (1 - \lambda_{j_F})(1 - \mu_{j_F})} \right\} \right\rangle.$$

Moreover the β -normalization is used in case of different cardinalities.

Definition 3.11: The Einstein product of two NCHFSs $\alpha = \langle A, \lambda \rangle, \beta = \langle B, \mu \rangle$ is defined by

$$\alpha \otimes_E \beta = \left\langle \left\{ \left[\frac{A_{j_T}^L B_{j_T}^L}{1 + (1 - A_{j_T}^L)(1 - B_{j_T}^L)}, \frac{A_{j_T}^U B_{j_T}^U}{1 + (1 - A_{j_T}^U)(1 - B_{j_T}^U)} \right] \right\}, \left\{ \left[\frac{A_{j_I}^L B_{j_I}^L}{1 + (1 - A_{j_I}^L)(1 - B_{j_I}^L)}, \frac{A_{j_I}^U B_{j_I}^U}{1 + (1 - A_{j_I}^U)(1 - B_{j_I}^U)} \right] \right\}, \right. \\ \left. \left\{ \left[\frac{A_{j_F}^L + B_{j_F}^L}{1 + A_{j_F}^L B_{j_F}^L}, \frac{A_{j_F}^U + B_{j_F}^U}{1 + A_{j_F}^U B_{j_F}^U} \right] \right\}, \left\{ \frac{\lambda_{j_T} \mu_{j_T}}{1 + (1 - \lambda_{j_T})(1 - \mu_{j_T})} \right\}, \left\{ \frac{\lambda_{j_I} \mu_{j_I}}{1 + (1 - \lambda_{j_I})(1 - \mu_{j_I})} \right\}, \left\{ \frac{\lambda_{j_F} + \mu_{j_F}}{1 + \lambda_{j_F} \mu_{j_F}} \right\} \right\rangle.$$

Moreover the β -normalization is used in case of different cardinalities.

Definition 3.12: The Einstein scalar multiplication of a scalar q with a NCHFS $\alpha = \langle A, \lambda \rangle$ is defined by

$$q_E \alpha = \left\langle \left\{ \left[\frac{(1+A_{j_T}^L)^q - (1-A_{j_T}^L)^q}{(1+A_{j_T}^L)^q + (1-A_{j_T}^L)^q}, \frac{(1+A_{j_T}^U)^q - (1-A_{j_T}^U)^q}{(1+A_{j_T}^U)^q + (1-A_{j_T}^U)^q} \right], \left[\frac{(1+A_{j_I}^L)^q - (1-A_{j_I}^L)^q}{(1+A_{j_I}^L)^q + (1-A_{j_I}^L)^q}, \frac{(1+A_{j_I}^U)^q - (1-A_{j_I}^U)^q}{(1+A_{j_I}^U)^q + (1-A_{j_I}^U)^q} \right] \right\}, \left\{ \frac{(1+\lambda_{j_T})^q - (1-\lambda_{j_T})^q}{(1+\lambda_{j_T})^q + (1-\lambda_{j_T})^q}, \frac{(1+\lambda_{j_I})^q - (1-\lambda_{j_I})^q}{(1+\lambda_{j_I})^q + (1-\lambda_{j_I})^q}, \frac{2(\lambda_{j_F})^q}{(2-\lambda_{j_F})^q + (\lambda_{j_F})^q} \right\} \right\rangle$$

Theorem 3.13: For a scalar q and a NCHFS $\alpha = \langle A, \lambda \rangle$ we have

$$\alpha^{E^q} = \left\langle \left\{ \left[\frac{2(A_{j_T}^L)^q}{(2-A_{j_T}^L)^q + (A_{j_T}^L)^q}, \frac{2(A_{j_T}^U)^q}{(2-A_{j_T}^U)^q + (A_{j_T}^U)^q} \right], \left[\frac{2(A_{j_I}^L)^q}{(2-A_{j_I}^L)^q + (A_{j_I}^L)^q}, \frac{2(A_{j_I}^U)^q}{(2-A_{j_I}^U)^q + (A_{j_I}^U)^q} \right] \right\}, \left\{ \frac{(1+\lambda_{j_T})^q - (1-\lambda_{j_T})^q}{(1+\lambda_{j_T})^q + (1-\lambda_{j_T})^q}, \frac{(1+\lambda_{j_I})^q - (1-\lambda_{j_I})^q}{(1+\lambda_{j_I})^q + (1-\lambda_{j_I})^q}, \frac{2(\lambda_{j_F})^q}{(2-\lambda_{j_F})^q + (\lambda_{j_F})^q} \right\} \right\rangle$$

where $\alpha^{E^q} = \alpha \otimes_E \alpha \otimes_E \dots \otimes_E \alpha$ (q -times) moreover α^{E^q} is a NCHF value for every $q > 0$.

Proof: Using induction on q . for $q=1$ we have

$$\alpha^{E^1} = \left\langle \left\{ \left[\frac{2(A_{j_T}^L)}{(2-A_{j_T}^L) + (A_{j_T}^L)}, \frac{2(A_{j_T}^U)}{(2-A_{j_T}^U) + (A_{j_T}^U)} \right], \left[\frac{2(A_{j_I}^L)}{(2-A_{j_I}^L) + (A_{j_I}^L)}, \frac{2(A_{j_I}^U)}{(2-A_{j_I}^U) + (A_{j_I}^U)} \right] \right\}, \left\{ \frac{(1+\lambda_{j_T}) - (1-\lambda_{j_T})}{(1+\lambda_{j_T}) + (1-\lambda_{j_T})}, \frac{(1+\lambda_{j_I}) - (1-\lambda_{j_I})}{(1+\lambda_{j_I}) + (1-\lambda_{j_I})}, \frac{2(\lambda_{j_F})}{(2-\lambda_{j_F}) + (\lambda_{j_F})} \right\} \right\rangle$$

Assuming that result is true for $q=m$.

$$\alpha^{E^m} = \left\langle \left\{ \left[\frac{2(A_{j_T}^L)^m}{(2-A_{j_T}^L)^m + (A_{j_T}^L)^m}, \frac{2(A_{j_T}^U)^m}{(2-A_{j_T}^U)^m + (A_{j_T}^U)^m} \right], \left[\frac{2(A_{j_I}^L)^m}{(2-A_{j_I}^L)^m + (A_{j_I}^L)^m}, \frac{2(A_{j_I}^U)^m}{(2-A_{j_I}^U)^m + (A_{j_I}^U)^m} \right] \right\}, \left\{ \frac{(1+\lambda_{j_T})^m - (1-\lambda_{j_T})^m}{(1+\lambda_{j_T})^m + (1-\lambda_{j_T})^m}, \frac{(1+\lambda_{j_I})^m - (1-\lambda_{j_I})^m}{(1+\lambda_{j_I})^m + (1-\lambda_{j_I})^m}, \frac{2(\lambda_{j_F})^m}{(2-\lambda_{j_F})^m + (\lambda_{j_F})^m} \right\} \right\rangle$$

α^{E^m} is neutrosophic cubic hesitant fuzzy value. Using assumption, we have

$$\alpha^{E^m} \otimes \alpha^{E^1} = \left\{ \left[\frac{2(A_{j_T}^L)^m}{(2-A_{j_T}^L)^m + (A_{j_T}^L)^m}, \frac{2(A_{j_T}^U)^m}{(2-A_{j_T}^U)^m + (A_{j_T}^U)^m} \right], \left[\frac{2(A_{j_I}^L)^m}{(2-A_{j_I}^L)^m + (A_{j_I}^L)^m}, \frac{2(A_{j_I}^U)^m}{(2-A_{j_I}^U)^m + (A_{j_I}^U)^m} \right], \left[\frac{(1+A_{j_F}^L)^m - (1-A_{j_T}^L)^m}{(1+A_{j_F}^L)^m + (1-A_{j_T}^L)^m}, \frac{(1+A_{j_F}^U)^m - (1-A_{j_T}^U)^m}{(1+A_{j_F}^U)^m + (1-A_{j_T}^U)^m} \right], \left\{ \frac{2(\lambda_{j_T})^m}{(2-\lambda_{j_T})^m + (\lambda_{j_T})^m} \right\}, \left\{ \frac{2(\lambda_{j_I})^m}{(2-\lambda_{j_I})^m + (\lambda_{j_I})^m} \right\}, \left\{ \frac{(1+\lambda_{j_F})^m - (1-\lambda_{j_F})^m}{(1+\lambda_{j_F})^m + (1-\lambda_{j_F})^m} \right\} \right\} \otimes_E \left\{ \left[\frac{2(A_{j_T}^L)}{(2-A_{j_T}^L) + (A_{j_T}^L)}, \frac{2(A_{j_T}^U)}{(2-A_{j_T}^U) + (A_{j_T}^U)} \right], \left[\frac{2(A_{j_I}^L)}{(2-A_{j_I}^L) + (A_{j_I}^L)}, \frac{2(A_{j_I}^U)}{(2-A_{j_I}^U) + (A_{j_I}^U)} \right], \left[\frac{(1+A_{j_F}^L) - (1-A_{j_T}^L)}{(1+A_{j_F}^L) + (1-A_{j_T}^L)}, \frac{(1+A_{j_F}^U) - (1-A_{j_T}^U)}{(1+A_{j_F}^U) + (1-A_{j_T}^U)} \right], \left\{ \frac{2(\lambda_{j_T})}{(2-\lambda_{j_T}) + (\lambda_{j_T})} \right\}, \left\{ \frac{2(\lambda_{j_I})}{(2-\lambda_{j_I}) + (\lambda_{j_I})} \right\}, \left\{ \frac{(1+\lambda_{j_F}) - (1-\lambda_{j_F})}{(1+\lambda_{j_F}) + (1-\lambda_{j_F})} \right\} \right\}$$

$$= \left\{ \left[\frac{4(A_{j_T}^L)^{m+1}}{\left(\frac{2(A_{j_T}^L)^m}{(2-A_{j_T}^L)^m + (A_{j_T}^L)^m} \right) \left(\frac{2(A_{j_T}^L)^1}{(2-A_{j_T}^L)^1 + (A_{j_T}^L)^1} \right)}, \frac{4(A_{j_T}^U)^{m+1}}{\left(\frac{2(A_{j_T}^U)^m}{(2-A_{j_T}^U)^m + (A_{j_T}^U)^m} \right) \left(\frac{2(A_{j_T}^U)^1}{(2-A_{j_T}^U)^1 + (A_{j_T}^U)^1} \right)} \right], \left[\frac{4(A_{j_I}^L)^{m+1}}{\left(\frac{2(A_{j_I}^L)^m}{(2-A_{j_I}^L)^m + (A_{j_I}^L)^m} \right) \left(\frac{2(A_{j_I}^L)^1}{(2-A_{j_I}^L)^1 + (A_{j_I}^L)^1} \right)}, \frac{4(A_{j_I}^U)^{m+1}}{\left(\frac{2(A_{j_I}^U)^m}{(2-A_{j_I}^U)^m + (A_{j_I}^U)^m} \right) \left(\frac{2(A_{j_I}^U)^1}{(2-A_{j_I}^U)^1 + (A_{j_I}^U)^1} \right)} \right], \left[\frac{\left(\frac{(1+A_{j_F}^L)^m - (1-A_{j_T}^L)^m}{(1+A_{j_F}^L)^m + (1-A_{j_T}^L)^m} \right) + \left(\frac{(1+A_{j_F}^L)^1 - (1-A_{j_T}^L)^1}{(1+A_{j_F}^L)^1 + (1-A_{j_T}^L)^1} \right)}{1 + \left(\frac{(1+A_{j_F}^L)^m - (1-A_{j_T}^L)^m}{(1+A_{j_F}^L)^m + (1-A_{j_T}^L)^m} \right) \left(\frac{(1+A_{j_F}^L)^1 - (1-A_{j_T}^L)^1}{(1+A_{j_F}^L)^1 + (1-A_{j_T}^L)^1} \right)}, \frac{\left(\frac{(1+A_{j_F}^U)^m - (1-A_{j_T}^U)^m}{(1+A_{j_F}^U)^m + (1-A_{j_T}^U)^m} \right) + \left(\frac{(1+A_{j_F}^U)^1 - (1-A_{j_T}^U)^1}{(1+A_{j_F}^U)^1 + (1-A_{j_T}^U)^1} \right)}{1 + \left(\frac{(1+A_{j_F}^U)^m - (1-A_{j_T}^U)^m}{(1+A_{j_F}^U)^m + (1-A_{j_T}^U)^m} \right) \left(\frac{(1+A_{j_F}^U)^1 - (1-A_{j_T}^U)^1}{(1+A_{j_F}^U)^1 + (1-A_{j_T}^U)^1} \right)} \right], \left\{ \frac{4(\lambda_{j_T})^{m+1}}{\left(\frac{2(\lambda_{j_T})^m}{(2-\lambda_{j_T})^m + (\lambda_{j_T})^m} \right) \left(\frac{2(\lambda_{j_T})^1}{(2-\lambda_{j_T})^1 + (\lambda_{j_T})^1} \right)}, \left\{ \frac{(1+\lambda_{j_F})^m - (1-\lambda_{j_F})^m}{(1+\lambda_{j_F})^m + (1-\lambda_{j_F})^m} + \frac{(1+\lambda_{j_F})^1 - (1-\lambda_{j_F})^1}{(1+\lambda_{j_F})^1 + (1-\lambda_{j_F})^1} \right\}}{1 + \left(\frac{(1+\lambda_{j_F})^m - (1-\lambda_{j_F})^m}{(1+\lambda_{j_F})^m + (1-\lambda_{j_F})^m} \right) \left(\frac{(1+\lambda_{j_F})^1 - (1-\lambda_{j_F})^1}{(1+\lambda_{j_F})^1 + (1-\lambda_{j_F})^1} \right)} \right\} \right\}$$

$$\begin{aligned}
 & \left\{ \left[\frac{4(A_{jT}^L)^{m+1}}{\left((2-A_{jT}^L)^m + (A_{jT}^L)^m \right) \left((2-A_{jT}^L)^1 + (A_{jT}^L)^1 \right)} \right], \left[\frac{4(A_{jT}^U)^{m+1}}{\left((2-A_{jT}^U)^m + (A_{jT}^U)^m \right) \left((2-A_{jT}^U)^1 + (A_{jT}^U)^1 \right)} \right] \right\}, \\
 & \left\{ \left[\frac{4(A_{jI}^L)^{m+1}}{\left((2-A_{jI}^L)^m + (A_{jI}^L)^m \right) \left((2-A_{jI}^L)^1 + (A_{jI}^L)^1 \right)} \right], \left[\frac{4(A_{jI}^U)^{m+1}}{\left((2-A_{jI}^U)^m + (A_{jI}^U)^m \right) \left((2-A_{jI}^U)^1 + (A_{jI}^U)^1 \right)} \right] \right\}, \\
 & \left\{ \left[\frac{\left((1+A_{jF}^L)^m - (1-A_{jF}^L)^m \right) \left((1+A_{jF}^L)^1 + (1-A_{jF}^L)^1 \right) + \left((1+A_{jF}^L)^m + (1-A_{jF}^L)^m \right) \left((1+A_{jF}^L)^1 - (1-A_{jF}^L)^1 \right)}{\left((1+A_{jF}^L)^m + (1-A_{jF}^L)^m \right) \left((1+A_{jF}^L)^1 + (1-A_{jF}^L)^1 \right)} \right], \left[\frac{\left((1+A_{jF}^U)^m - (1-A_{jF}^U)^m \right) \left((1+A_{jF}^U)^1 + (1-A_{jF}^U)^1 \right) + \left((1+A_{jF}^U)^m + (1-A_{jF}^U)^m \right) \left((1+A_{jF}^U)^1 - (1-A_{jF}^U)^1 \right)}{\left((1+A_{jF}^U)^m + (1-A_{jF}^U)^m \right) \left((1+A_{jF}^U)^1 + (1-A_{jF}^U)^1 \right)} \right] \right\}, \\
 & \left\{ \left[\frac{4(\lambda_{jT})^{m+1}}{\left((2-\lambda_{jT})^m + (\lambda_{jT})^m \right) \left((2-\lambda_{jT})^1 + (\lambda_{jT})^1 \right)} \right], \left[\frac{4(\lambda_{jI})^{m+1}}{\left((2-\lambda_{jI})^m + (\lambda_{jI})^m \right) \left((2-\lambda_{jI})^1 + (\lambda_{jI})^1 \right)} \right] \right\}, \\
 & \left\{ \left[\frac{\left((1+\lambda_{jF})^m - (1-\lambda_{jF})^m \right) \left((1+\lambda_{jF})^1 + (1-\lambda_{jF})^1 \right) + \left((1+\lambda_{jF})^m + (1-\lambda_{jF})^m \right) \left((1+\lambda_{jF})^1 - (1-\lambda_{jF})^1 \right)}{\left((1+\lambda_{jF})^m + (1-\lambda_{jF})^m \right) \left((1+\lambda_{jF})^1 + (1-\lambda_{jF})^1 \right)} \right], \left[\frac{\left((1+\lambda_{jF})^m - (1-\lambda_{jF})^m \right) \left((1+\lambda_{jF})^1 + (1-\lambda_{jF})^1 \right) + \left((1+\lambda_{jF})^m + (1-\lambda_{jF})^m \right) \left((1+\lambda_{jF})^1 - (1-\lambda_{jF})^1 \right)}{\left((1+\lambda_{jF})^m + (1-\lambda_{jF})^m \right) \left((1+\lambda_{jF})^1 + (1-\lambda_{jF})^1 \right)} \right] \right\}
 \end{aligned}$$

$$\begin{aligned}
 &= \left\langle \left\{ \left[\frac{4(A_{j_T}^L)^{m+1}}{2((2-A_{j_T}^L)^{m+1} + (A_{j_T}^L)^{m+1})}, \frac{4(A_{j_T}^U)^{m+1}}{2((2-A_{j_T}^U)^{m+1} + (A_{j_T}^U)^{m+1})} \right], \right. \right. \\
 &\quad \left. \left[\frac{4(A_{j_I}^L)^{m+1}}{2((2-A_{j_I}^L)^{m+1} + (A_{j_I}^L)^{m+1})}, \frac{4(A_{j_I}^U)^{m+1}}{2((2-A_{j_I}^U)^{m+1} + (A_{j_I}^U)^{m+1})} \right], \right. \\
 &\quad \left. \left[\frac{(1+A_{j_F}^L)^{m+1} - (1-A_{j_T}^L)^{m+1}}{(1+A_{j_F}^L)^{m+1} + (1-A_{j_T}^L)^{m+1}}, \frac{(1+A_{j_F}^U)^{m+1} - (1-A_{j_T}^U)^{m+1}}{(1+A_{j_F}^U)^{m+1} + (1-A_{j_T}^U)^{m+1}} \right], \right. \\
 &\quad \left. \left. \left. \left. \left. \frac{4(\lambda_{j_T})^{m+1}}{2((2-\lambda_{j_T})^{m+1} + (\lambda_{j_T})^{m+1})} \right\}, \left. \frac{4(\lambda_{j_I})^{m+1}}{2((2-\lambda_{j_I})^{m+1} + (\lambda_{j_I})^{m+1})} \right\}, \right. \right. \right. \\
 &\quad \left. \left. \left. \left. \left. \frac{(1+\lambda_{j_F})^{m+1} - (1-\lambda_{j_F})^{m+1}}{(1+\lambda_{j_F})^{m+1} + (1-\lambda_{j_F})^{m+1}} \right\} \right. \right. \right. \\
 &= \left\langle \left\{ \left[\frac{2(A_{j_T}^L)^{m+1}}{(2-A_{j_T}^L)^{m+1} + (A_{j_T}^L)^{m+1}}, \frac{2(A_{j_T}^U)^{m+1}}{(2-A_{j_T}^U)^{m+1} + (A_{j_T}^U)^{m+1}} \right], \left[\frac{2(A_{j_I}^L)^{m+1}}{(2-A_{j_I}^L)^{m+1} + (A_{j_I}^L)^{m+1}}, \frac{2(A_{j_I}^U)^{m+1}}{(2-A_{j_I}^U)^{m+1} + (A_{j_I}^U)^{m+1}} \right] \right\} \right. \\
 &\quad \left. \left\{ \left[\frac{(1+A_{j_F}^L)^{m+1} - (1-A_{j_T}^L)^{m+1}}{(1+A_{j_F}^L)^{m+1} + (1-A_{j_T}^L)^{m+1}}, \frac{(1+A_{j_F}^U)^{m+1} - (1-A_{j_T}^U)^{m+1}}{(1+A_{j_F}^U)^{m+1} + (1-A_{j_T}^U)^{m+1}} \right], \left. \left. \left. \left. \left. \frac{2(\lambda_{j_T})^{m+1}}{(2-\lambda_{j_T})^{m+1} + (\lambda_{j_T})^{m+1}} \right\}, \left. \frac{2(\lambda_{j_I})^{m+1}}{(2-\lambda_{j_I})^{m+1} + (\lambda_{j_I})^{m+1}} \right\}, \right. \right. \right. \right. \right. \\
 &\quad \left. \left. \left. \left. \left. \frac{(1+\lambda_{j_F})^{m+1} - (1-\lambda_{j_F})^{m+1}}{(1+\lambda_{j_F})^{m+1} + (1-\lambda_{j_F})^{m+1}} \right\} \right. \right. \right. \right. \right.
 \end{aligned}$$

Definition 3.14: The Score, accuracy and certainty of a $NCHFS\alpha = \langle A, \lambda \rangle$ where $A = \langle A_T, A_I, A_F \rangle, A_T = \{[A_{j_T}^L, A_{j_T}^U]; j = 1, \dots, l\}, A_I = \{[A_{j_I}^L, A_{j_I}^U]; j = 1, \dots, m\}, A_F = \{[A_{j_F}^L, A_{j_F}^U]; i = 1, \dots, n\}$ and $\lambda = \langle \lambda_T, \lambda_I, \lambda_F \rangle, \lambda_T = \{\lambda_{j_T}; j = 1, \dots, r\}, \lambda_I = \{\lambda_{j_I}; j = 1, \dots, s\}, \lambda_F = \{\lambda_{j_F}; j = 1, \dots, t\}$ are defined as:

$$S(\alpha) = \frac{1}{2} \left\{ \frac{1}{6} \left(\frac{1}{l} \sum_{j=1}^l (A_{j_T}^L + A_{j_T}^U) + \frac{1}{m} \sum_{j=1}^m (A_{j_I}^L + A_{j_I}^U) + \frac{1}{n} \sum_{j=1}^n (2 - (A_{j_T}^L + A_{j_T}^U)) \right) + \frac{1}{3} \left(\frac{1}{r} \sum_{j=1}^r \lambda_{j_T} + \frac{1}{s} \sum_{j=1}^s \lambda_{j_I} + \frac{1}{t} \sum_{j=1}^t (1 - \lambda_{j_F}) \right) \right\},$$

$$H(\alpha) = \frac{1}{9} \left\{ \frac{1}{l} \sum_{j=1}^l (A_{j_T}^L + A_{j_T}^U) + \frac{1}{m} \sum_{j=1}^m (A_{j_I}^L + A_{j_I}^U) + \frac{1}{n} \sum_{j=1}^n (A_{j_F}^U + A_{j_F}^L) + \frac{1}{r} \sum_{j=1}^r \lambda_{j_T} + \frac{1}{s} \sum_{j=1}^s \lambda_{j_I} + \frac{1}{t} \sum_{j=1}^t \lambda_{j_F} \right\},$$

$$C(\alpha) = \frac{1}{3} \left\{ \frac{1}{l} \sum_{j=1}^l (A_{j_T}^L + A_{j_T}^U) + \frac{1}{r} \sum_{j=1}^r \lambda_{j_T} \right\}.$$

If $\alpha = \langle \{[0.1, 0.5], [0.2, 0.7]\}, \{[0.2, 0.3], [0.1, 0.6]\}, \{[0.1, 0.4], [0, 0.3]\}, \{0.1, 0.2\}, \{0.3, 0.5\}, \{0.4, 0.8\} \rangle$, and

$\beta = \langle \{[0.4, 0.5], [0.3, 0.4]\}, \{[0.1, 0.3], [0.2, 0.5]\}, \{[0.1, 0.4], [0.7, 0.8]\}, \{0.3, 0.5\}, \{0.7, 0.8\}, \{0.4, 0.6\} \rangle$, then $S(\alpha) = 0.404167, S(\beta) = 0.470833, H(\alpha) = 0.3222, H(\beta) = 0.4444, C(\alpha) = 0.15, C(\beta) = 0.4$.




Figure 1: Scores, Accuracy and Certainty of above NCHFSs

Definition 3.15: Let $\alpha = \langle A, \lambda \rangle, \beta = \langle B, \mu \rangle$ are two NCHFSs. We say that $\alpha > \beta$ if $S(\alpha) > S(\beta)$. If $S(\alpha) = S(\beta)$, then $\alpha > \beta$ if $A(\alpha) > A(\beta)$. If $A(\alpha) = A(\beta)$, then $\alpha > \beta$ if $C(\alpha) > C(\beta)$. If $S(\alpha) = S(\beta), A(\alpha) = A(\beta), C(\alpha) > C(\beta)$, then $\alpha = \beta$.

In the next section we define aggregation operators on neutrosophic cubic hesitant fuzzy set and prove some elegant results.

4. Aggregation Operators

Definition 4.1: The Neutrosophic cubic hesitant fuzzy weighted geometric operator is defined as

$$NCHWG(\alpha_1, \dots, \alpha_n) = \bigotimes_{j=1}^n \alpha_j^{w_j}, \text{ where } \alpha_{j(k)} \text{ are neutrosophic cubic hesitant fuzzy values taken in}$$

descending order with corresponding weight vector $w = (w_1, \dots, w_n)^t$.

Definition 4.2: Neutrosophic cubic hesitant fuzzy order weighted geometric operator is defined as:

$$NCHFOWG(\alpha_1, \dots, \alpha_n) = \bigotimes_{j=1}^n (\alpha_{(k)_j})^{w_j} \text{ where } \alpha_{j(k)} \text{ are neutrosophic cubic hesitant fuzzy values taken}$$

in descending order with corresponding weight vector $w = (w_1, \dots, w_n)^t$.

Definition 4.3: The Neutrosophic cubic hesitant fuzzy Einstein weighted geometric operator is defined as:

$$NCEHWG(\alpha_1, \dots, \alpha_n) = \bigotimes_{j=1}^n (\alpha_j^E)^{w_j} \text{ where } \alpha_{j(k)} \text{ are neutrosophic cubic hesitant fuzzy values taken in}$$

descending order with corresponding weight vector $w = (w_1, \dots, w_n)^t$

Definition 4.4: Neutrosophic cubic hesitant fuzzy Einstein ordered weighted geometric operator is defined as:

$$NCHEOWG(\alpha_1, \dots, \alpha_n) = \bigotimes_{j=1}^n (\alpha_{(k)_j}^E)^{w_j} \text{ where } \alpha_{j(k)} \text{ are neutrosophic cubic hesitant fuzzy values taken in}$$

descending order with corresponding weight vector $w = (w_1, \dots, w_n)^t$

Theorem 4.5: Let $\{\alpha_{(k)} = \langle A_{(k)}, \lambda_{(k)} \rangle\}$ the set of neutrosophic cubic hesitant with corresponding weight vector $w = (w_1, \dots)^t$ fuzzy values, then

$$NCHFVG(\alpha_1, \dots, \alpha_m) = \left\langle \left\{ \left[\prod_{k=1}^m (A_{P_{T(k)}}^L)^{w_k}, \prod_{k=1}^m (A_{P_{T(k)}}^U)^{w_k} \right], \left[\prod_{k=1}^m (A_{P_{I(k)}}^L)^{w_k}, \prod_{k=1}^m (A_{P_{I(k)}}^U)^{w_k} \right] \right\}, \right. \\ \left. \left\{ \left[1 - \prod_{k=1}^m (1 - A_{P_{F(k)}}^L)^{w_k}, 1 - \prod_{k=1}^m (1 - A_{P_{F(k)}}^U)^{w_k} \right], \left\{ \prod_{k=1}^m (\lambda_{P_{T(k)}})^{w_k} \right\}, \right. \right. \\ \left. \left. \left\{ \prod_{k=1}^m (\lambda_{P_{I(k)}})^{w_k} \right\}, \left\{ 1 - \prod_{k=1}^m (1 - \lambda_{P_{F(k)}})^{w_k} \right\} \right\} \right\rangle$$

Proof: Using induction for $m=2$

$$NCHFVG(\alpha_1, \alpha_2) = \bigotimes_{k=1}^2 \alpha_k^{w_k} \\ = \left\langle \left\{ \left[\prod_{k=1}^2 (A_{P_{T(k)}}^L)^{w_k}, \prod_{k=1}^2 (A_{P_{T(k)}}^U)^{w_k} \right], \left[\prod_{k=1}^2 (A_{P_{I(k)}}^L)^{w_k}, \prod_{k=1}^2 (A_{P_{I(k)}}^U)^{w_k} \right] \right\}, \right. \\ \left. \left\{ \left[1 - \prod_{k=1}^2 (1 - A_{P_{F(k)}}^L)^{w_k}, 1 - \prod_{k=1}^2 (1 - A_{P_{F(k)}}^U)^{w_k} \right], \left\{ \prod_{k=1}^2 (\lambda_{P_{T(k)}})^{w_k} \right\}, \right. \right. \\ \left. \left. \left\{ \prod_{k=1}^2 (\lambda_{P_{I(k)}})^{w_k} \right\}, \left\{ 1 - \prod_{k=1}^2 (1 - \lambda_{P_{F(k)}})^{w_k} \right\} \right\} \right\rangle$$

For $m = q$ we have

$$NCHFVG(\alpha_1, \dots, \alpha_q) = \left\langle \left\{ \left[\prod_{k=1}^q (A_{P_{T(k)}}^L)^{w_k}, \prod_{k=1}^q (A_{P_{T(k)}}^U)^{w_k} \right], \left[\prod_{k=1}^q (A_{P_{I(k)}}^L)^{w_k}, \prod_{k=1}^q (A_{P_{I(k)}}^U)^{w_k} \right] \right\}, \right. \\ \left. \left\{ \left[1 - \prod_{k=1}^q (1 - A_{P_{F(k)}}^L)^{w_k}, 1 - \prod_{k=1}^q (1 - A_{P_{F(k)}}^U)^{w_k} \right], \left\{ \prod_{k=1}^q (\lambda_{P_{T(k)}})^{w_k} \right\}, \right. \right. \\ \left. \left. \left\{ \prod_{k=1}^q (\lambda_{P_{I(k)}})^{w_k} \right\}, \left\{ 1 - \prod_{k=1}^q (1 - \lambda_{P_{F(k)}})^{w_k} \right\} \right\} \right\rangle$$

we prove for $m=q+1$

$$\bigotimes_{k=1}^q (\alpha_k)^{w_k} \otimes (\alpha_{q+1})^{w_{q+1}} = \left\langle \left\{ \left[\prod_{k=1}^q (A_{P_{T(k)}}^L)^{w_k}, \prod_{k=1}^q (A_{P_{T(k)}}^U)^{w_k} \right], \left[\prod_{k=1}^q (A_{P_{I(k)}}^L)^{w_k}, \prod_{k=1}^q (A_{P_{I(k)}}^U)^{w_k} \right] \right\}, \right. \\ \left. \left\{ \left[1 - \prod_{k=1}^q (1 - A_{P_{F(k)}}^L)^{w_k}, 1 - \prod_{k=1}^q (1 - A_{P_{F(k)}}^U)^{w_k} \right], \left\{ \prod_{k=1}^q (\lambda_{P_{T(k)}})^{w_k} \right\}, \left\{ \prod_{k=1}^q (\lambda_{P_{I(k)}})^{w_k} \right\}, \right. \right. \\ \left. \left. \left\{ 1 - \prod_{k=1}^q (1 - \lambda_{P_{F(k)}})^{w_k} \right\} \right\} \right\rangle \otimes \left\langle \left\{ \left[(A_{P_{T(q+1)}}^L)^{w_{q+1}}, (A_{P_{T(q+1)}}^U)^{w_{q+1}} \right], \left[(A_{P_{I(q+1)}}^L)^{w_{q+1}}, (A_{P_{I(q+1)}}^U)^{w_{q+1}} \right] \right\}, \right. \\ \left. \left\{ \left[1 - (1 - A_{P_{F(q+1)}}^L)^{w_{q+1}}, 1 - (1 - A_{P_{F(q+1)}}^U)^{w_{q+1}} \right], \left\{ (\lambda_{P_{T(q+1)}})^{w_{q+1}} \right\}, \right. \right. \\ \left. \left. \left\{ (\lambda_{P_{I(q+1)}})^{w_{q+1}} \right\}, \left\{ 1 - (1 - \lambda_{P_{F(q+1)}})^{w_{q+1}} \right\} \right\} \right\rangle$$

$$= \left\{ \left[\left[\prod_{k=1}^q \left(A_{p_T(k)}^L \right)^{w_k} \left(A_{p_T(q+1)}^L \right)^{w_{q+1}} \right], \left[\prod_{k=1}^q \left(A_{p_T(k)}^U \right)^{w_k} \left(A_{p_T(k)}^U \right)^{w_{q+1}} \right] \right], \left[\left[\prod_{k=1}^q \left(A_{p_I(k)}^L \right)^{w_k} \left(A_{p_I(q+1)}^L \right)^{w_{q+1}} \right], \left[\prod_{k=1}^q \left(A_{p_I(k)}^U \right)^{w_k} \left(A_{p_I(q+1)}^U \right)^{w_{q+1}} \right] \right] \right\},$$

$$\left\{ \left[\left[1 - \prod_{k=1}^q \left(1 - A_{p_F(k)}^L \right)^{w_k} + 1 - \left(1 - A_{p_F(q+1)}^L \right)^{w_{q+1}} - \left(\left[\prod_{k=1}^q \left(1 - A_{p_F(k)}^L \right)^{w_k} \right] \left[1 - \left(1 - A_{p_F(q+1)}^L \right)^{w_{q+1}} \right] \right) \right], \right. \right.$$

$$\left. \left[\left[1 - \prod_{k=1}^q \left(1 - A_{p_F(k)}^U \right)^{w_k} + 1 - \left(1 - A_{p_F(q+1)}^U \right)^{w_{q+1}} - \left(\left[\prod_{k=1}^q \left(1 - A_{p_F(k)}^U \right)^{w_k} \right] \left[1 - \left(1 - A_{p_F(q+1)}^U \right)^{w_{q+1}} \right] \right) \right] \right] \right\},$$

$$\left\{ \left[\prod_{k=1}^q \left(\lambda_{p_T(k)} \right)^{w_k} \left(\lambda_{p_T(q+1)} \right)^{w_{q+1}} \right], \left[\prod_{k=1}^q \left(\lambda_{p_I(k)} \right)^{w_k} \left(\lambda_{p_I(q+1)} \right)^{w_{q+1}} \right] \right\},$$

$$\left\{ \left[1 - \prod_{k=1}^q \left(1 - \lambda_{p_F(k)} \right)^{w_k} + 1 - \left(1 - \lambda_{p_F(q+1)} \right)^{w_{q+1}} - \left(\left[\prod_{k=1}^q \left(1 - \lambda_{p_F(k)} \right)^{w_k} \right] \left[1 - \left(1 - \lambda_{p_F(q+1)} \right)^{w_{q+1}} \right] \right) \right] \right\}$$

$$\left\{ \left[\prod_{k=1}^{q+1} \left(A_{p_T(k)}^L \right)^{w_k} \right], \left[\prod_{k=1}^{q+1} \left(A_{p_T(k)}^U \right)^{w_k} \right] \right\}, \left\{ \left[\prod_{k=1}^{q+1} \left(A_{p_I(k)}^L \right)^{w_k} \right], \left[\prod_{k=1}^{q+1} \left(A_{p_I(k)}^U \right)^{w_k} \right] \right\},$$

$$\left\{ \left[\left[2 - \prod_{k=1}^{q+1} \left(1 - A_{p_F(k)}^L \right)^{w_k} - 1 + \prod_{k=1}^q \left(1 - A_{p_F(k)}^L \right) + \left(1 - A_{p_F(q+1)}^L \right)^{w_{q+1}} - \left(\left[\prod_{k=1}^q \left(1 - A_{p_F(k)}^L \right)^{w_k} \right] \left[1 - A_{p_F(q+1)}^L \right] \right) \right], \right. \right.$$

$$\left. \left[\left[2 - \prod_{k=1}^{q+1} \left(1 - A_{p_F(k)}^U \right)^{w_k} - 1 + \prod_{k=1}^q \left(1 - A_{p_F(k)}^U \right) + \left(1 - A_{p_F(q+1)}^U \right)^{w_{q+1}} - \left(\left[\prod_{k=1}^q \left(1 - A_{p_F(k)}^U \right)^{w_k} \right] \left[1 - A_{p_F(q+1)}^U \right] \right) \right] \right] \right\},$$

$$\left\{ \left[\prod_{k=1}^{q+1} \left(\lambda_{p_T(k)} \right)^{w_k} \right], \left[\prod_{k=1}^{q+1} \left(\lambda_{p_I(k)} \right)^{w_k} \right] \right\},$$

$$\left\{ \left[\left[2 - \prod_{k=1}^{q+1} \left(1 - \lambda_{p_F(k)} \right)^{w_k} - 1 + \prod_{k=1}^q \left(1 - \lambda_{p_F(k)} \right) + \left(1 - \lambda_{p_F(k)} \right)^{w_{q+1}} \right], \right. \right.$$

$$\left. \left[\left[\prod_{k=1}^q \left(1 - \lambda_{p_F(k)} \right)^{w_k} \right] \left[1 - \lambda_{p_F(k)} \right] \right] \right\}$$

$$= \left(\left[\left[\prod_{k=1}^{q+1} \left(A_{p_{T(k)}}^L \right)^{w_k}, \prod_{k=1}^{q+1} \left(A_{p_{T(k)}}^U \right)^{w_k} \right], \left[\prod_{k=1}^{q+1} \left(A_{p_{I(k)}}^L \right)^{w_k}, \prod_{k=1}^{q+1} \left(A_{p_{I(k)}}^U \right)^{w_k} \right] \right], \left[\prod_{k=1}^{q+1} \left(1 - A_{p_{F(k)}}^L \right)^{w_k}, \prod_{k=1}^{q+1} \left(1 - A_{p_{F(k)}}^U \right)^{w_k} \right], \left[\prod_{k=1}^{q+1} \left(\lambda_{p_{T(k)}} \right)^{w_k} \right], \left[\prod_{k=1}^{q+1} \left(\lambda_{p_{I(k)}} \right)^{w_k} \right], \left[\prod_{k=1}^{q+1} \left(1 - \lambda_{p_{F(k)}} \right)^{w_k} \right] \right)$$

Theorem 4.6: Let $\{\alpha_{(k)} = \langle A_{(k)}, \lambda_{(k)} \rangle\}$ the set of neutrosophic cubic hesitant fuzzy values, then

i) Idempotency: If $\alpha_k = \alpha, k = 1, \dots, m$ then $NCHFVG(\alpha_1, \dots, \alpha_m) = \alpha$.

ii) Monotonicity: If $S(\alpha_q) \geq S(\alpha_l)$ then $NCHFVG(\alpha_q) \leq NCHFVG(\alpha_l)$.

Theorem 4.7: Let $\{\alpha_{(k)} = \langle A_{(k)}, \lambda_{(k)} \rangle\}$ the set of neutrosophic cubic hesitant with corresponding weight vector $w = (w_1, \dots)^t$ fuzzy values, then

$$NCHFEGW(\alpha_1, \dots, \alpha_m) = \left\{ \left[\frac{2 \prod_{k=1}^m \left(A_{T(k)}^L \right)^{w_k}}{2 \prod_{k=1}^m \left(A_{T(k)}^U \right)^{w_k}}, \frac{2 \prod_{k=1}^m \left(A_{T(k)}^U \right)^{w_k}}{2 \prod_{k=1}^m \left(A_{T(k)}^L \right)^{w_k}} \right], \left[\frac{2 \prod_{k=1}^m \left(A_{I(k)}^L \right)^{w_k}}{2 \prod_{k=1}^m \left(A_{I(k)}^U \right)^{w_k}}, \frac{2 \prod_{k=1}^m \left(A_{I(k)}^U \right)^{w_k}}{2 \prod_{k=1}^m \left(A_{I(k)}^L \right)^{w_k}} \right], \right. \\
 \left. \left[\frac{2 \prod_{k=1}^m \left(2 - A_{T(k)}^L \right)^{w_k} + \prod_{k=1}^m \left(A_{T(k)}^L \right)^{w_k}}{2 \prod_{k=1}^m \left(2 - A_{T(k)}^U \right)^{w_k} + \prod_{k=1}^m \left(A_{T(k)}^U \right)^{w_k}}, \frac{2 \prod_{k=1}^m \left(2 - A_{I(k)}^L \right)^{w_k} + \prod_{k=1}^m \left(A_{I(k)}^L \right)^{w_k}}{2 \prod_{k=1}^m \left(2 - A_{I(k)}^U \right)^{w_k} + \prod_{k=1}^m \left(A_{I(k)}^U \right)^{w_k}} \right], \right. \\
 \left. \left[\frac{\prod_{k=1}^m \left(1 + A_{F(k)}^L \right)^{w_k} - \prod_{k=1}^m \left(1 - A_{F(k)}^L \right)^{w_k}}{\prod_{k=1}^m \left(1 + A_{F(k)}^U \right)^{w_k} - \prod_{k=1}^m \left(1 - A_{F(k)}^U \right)^{w_k}}, \frac{\prod_{k=1}^m \left(1 + A_{F(k)}^U \right)^{w_k} - \prod_{k=1}^m \left(1 - A_{F(k)}^U \right)^{w_k}}{\prod_{k=1}^m \left(1 + A_{F(k)}^L \right)^{w_k} - \prod_{k=1}^m \left(1 - A_{F(k)}^L \right)^{w_k}} \right], \right. \\
 \left. \left[\frac{2 \prod_{k=1}^m \left(\lambda_{PT(k)} \right)^{w_k}}{2 \prod_{k=1}^m \left(\lambda_{PI(k)} \right)^{w_k}}, \frac{2 \prod_{k=1}^m \left(\lambda_{PI(k)} \right)^{w_k}}{2 \prod_{k=1}^m \left(\lambda_{PT(k)} \right)^{w_k}} \right], \right. \\
 \left. \left[\frac{\prod_{k=1}^m \left(1 + \lambda_{PF(k)} \right)^{w_k} - \prod_{k=1}^m \left(1 - \lambda_{PF(k)} \right)^{w_k}}{\prod_{k=1}^m \left(1 + \lambda_{PI(k)} \right)^{w_k} + \prod_{k=1}^m \left(\lambda_{PI(k)} \right)^{w_k}}, \frac{\prod_{k=1}^m \left(1 + \lambda_{PI(k)} \right)^{w_k} + \prod_{k=1}^m \left(\lambda_{PI(k)} \right)^{w_k}}{\prod_{k=1}^m \left(1 + \lambda_{PF(k)} \right)^{w_k} - \prod_{k=1}^m \left(1 - \lambda_{PF(k)} \right)^{w_k}} \right] \right\}$$

Proof: we use induction.

$$\alpha^{E^{w_1}} = \left\{ \left[\frac{2 \left(A_{j_T}^L \right)^{w_1}}{\left(2 - A_{j_T}^L \right)^{w_1} + \left(A_{j_T}^L \right)^{w_1}}, \frac{2 \left(A_{j_T}^U \right)^{w_1}}{\left(2 - A_{j_T}^U \right)^{w_1} + \left(A_{j_T}^U \right)^{w_1}} \right], \left[\frac{2 \left(A_{j_I}^L \right)^{w_1}}{\left(2 - A_{j_I}^L \right)^{w_1} + \left(A_{j_I}^L \right)^{w_1}}, \frac{2 \left(A_{j_I}^U \right)^{w_1}}{\left(2 - A_{j_I}^U \right)^{w_1} + \left(A_{j_I}^U \right)^{w_1}} \right], \right. \\
 \left. \left[\frac{\left(1 + A_{j_F}^L \right)^{w_1} - \left(1 - A_{j_F}^L \right)^{w_1}}{\left(1 + A_{j_F}^U \right)^{w_1} - \left(1 - A_{j_F}^U \right)^{w_1}}, \frac{\left(1 + A_{j_F}^U \right)^{w_1} - \left(1 - A_{j_F}^U \right)^{w_1}}{\left(1 + A_{j_F}^L \right)^{w_1} - \left(1 - A_{j_F}^L \right)^{w_1}} \right], \left[\frac{2 \left(\lambda_{j_T} \right)^{w_1}}{\left(2 - \lambda_{j_T} \right)^{w_1} + \left(\lambda_{j_T} \right)^{w_1}}, \frac{2 \left(\lambda_{j_I} \right)^{w_1}}{\left(2 - \lambda_{j_I} \right)^{w_1} + \left(\lambda_{j_I} \right)^{w_1}} \right], \right. \\
 \left. \left[\frac{\left(1 + \lambda_{j_F} \right)^{w_1} - \left(1 - \lambda_{j_F} \right)^{w_1}}{\left(1 + \lambda_{j_F} \right)^{w_1} + \left(1 - \lambda_{j_F} \right)^{w_1}} \right] \right\}$$

$$\text{and } \alpha^{E^{w_2}} = \left\langle \left\{ \left[\frac{2(A_{j_T}^L)^{w_2}}{(2-A_{j_T}^L)^{w_2} + (A_{j_T}^L)^{w_2}}, \frac{2(A_{j_T}^U)^{w_2}}{(2-A_{j_T}^U)^{w_2} + (A_{j_T}^U)^{w_2}} \right], \left[\frac{2(A_{j_I}^L)^{w_2}}{(2-A_{j_I}^L)^{w_2} + (A_{j_I}^L)^{w_2}}, \frac{2(A_{j_I}^U)^{w_2}}{(2-A_{j_I}^U)^{w_2} + (A_{j_I}^U)^{w_2}} \right] \right\}, \left\{ \left[\frac{(1+A_{j_F}^L)^{w_2} - (1-A_{j_T}^L)^{w_2}}{(1+A_{j_F}^L)^{w_2} + (1-A_{j_T}^L)^{w_2}}, \frac{(1+A_{j_F}^U)^{w_2} - (1-A_{j_T}^U)^{w_2}}{(1+A_{j_F}^U)^{w_2} + (1-A_{j_T}^U)^{w_2}} \right], \left[\frac{2(\lambda_{j_T})^{w_2}}{(2-\lambda_{j_T})^{w_2} + (\lambda_{j_T})^{w_2}} \right], \left[\frac{2(\lambda_{j_I})^{w_2}}{(2-\lambda_{j_I})^{w_2} + (\lambda_{j_I})^{w_2}} \right], \left[\frac{(1+\lambda_{j_F})^{w_2} - (1-\lambda_{j_F})^{w_2}}{(1+\lambda_{j_F})^{w_2} + (1-\lambda_{j_F})^{w_2}} \right] \right\} \right\rangle$$

$$NCHFEG(\alpha_1, \alpha_2) = \bigotimes_{k=1}^2 \alpha_k^{E^{w_k}}$$

$$= \left\langle \left\{ \left[\frac{2 \prod_{k=1}^2 (A_{p_{T(k)}}^L)^{w_k}}{\prod_{k=1}^2 (2-A_{p_{T(k)}}^L)^{w_k} + \prod_{k=1}^2 (A_{p_{T(k)}}^L)^{w_k}}, \frac{2 \prod_{k=1}^2 (A_{p_{T(k)}}^U)^{w_k}}{\prod_{k=1}^2 (2-A_{p_{T(k)}}^U)^{w_k} + \prod_{k=1}^2 (A_{p_{T(k)}}^U)^{w_k}} \right], \left[\frac{2 \prod_{k=1}^2 (A_{p_{I(k)}}^L)^{w_k}}{\prod_{k=1}^2 (2-A_{p_{I(k)}}^L)^{w_k} + \prod_{k=1}^2 (A_{p_{I(k)}}^L)^{w_k}}, \frac{2 \prod_{k=1}^2 (A_{p_{I(k)}}^U)^{w_k}}{\prod_{k=1}^2 (2-A_{p_{I(k)}}^U)^{w_k} + \prod_{k=1}^2 (A_{p_{I(k)}}^U)^{w_k}} \right] \right\}, \left\{ \left[\frac{\prod_{k=1}^2 (1+A_{p_{F(k)}}^L)^{w_k} - \prod_{k=1}^2 (1-A_{p_{F(k)}}^L)^{w_k}}{\prod_{k=1}^2 (1+A_{p_{F(k)}}^L)^{w_k} + \prod_{k=1}^2 (1-A_{p_{F(k)}}^L)^{w_k}}, \frac{\prod_{k=1}^2 (1+A_{p_{F(k)}}^U)^{w_k} - \prod_{k=1}^2 (1-A_{p_{F(k)}}^U)^{w_k}}{\prod_{k=1}^2 (1+A_{p_{F(k)}}^U)^{w_k} + \prod_{k=1}^2 (1-A_{p_{F(k)}}^U)^{w_k}} \right], \left[\frac{2 \prod_{k=1}^2 (\lambda_{p_{T(k)}})^{w_k}}{\prod_{k=1}^2 (2-\lambda_{p_{T(k)}})^{w_k} + \prod_{k=1}^2 (\lambda_{p_{T(k)}})^{w_k}} \right], \left[\frac{2 \prod_{k=1}^2 (\lambda_{p_{I(k)}})^{w_k}}{\prod_{k=1}^2 (2-\lambda_{p_{I(k)}})^{w_k} + \prod_{k=1}^2 (\lambda_{p_{I(k)}})^{w_k}} \right], \left[\frac{\prod_{k=1}^2 (1+\lambda_{p_{F(k)}})^{w_k} - \prod_{k=1}^2 (1-\lambda_{p_{F(k)}})^{w_k}}{\prod_{k=1}^2 (1+\lambda_{p_{F(k)}})^{w_k} + \prod_{k=1}^2 (1-\lambda_{p_{F(k)}})^{w_k}} \right] \right\} \right\rangle$$

For m=q we have

$$NCHFEG(\alpha_1, \dots, \alpha_q) = \left\langle \left\{ \left[\frac{2 \prod_{k=1}^q (A_{p_{T(k)}}^L)^{w_k}}{\prod_{k=1}^q (2-A_{p_{T(k)}}^L)^{w_k} + \prod_{k=1}^q (A_{p_{T(k)}}^L)^{w_k}}, \frac{2 \prod_{k=1}^q (A_{p_{T(k)}}^U)^{w_k}}{\prod_{k=1}^q (2-A_{p_{T(k)}}^U)^{w_k} + \prod_{k=1}^q (A_{p_{T(k)}}^U)^{w_k}} \right], \left[\frac{2 \prod_{k=1}^q (A_{p_{I(k)}}^L)^{w_k}}{\prod_{k=1}^q (2-A_{p_{I(k)}}^L)^{w_k} + \prod_{k=1}^q (A_{p_{I(k)}}^L)^{w_k}}, \frac{2 \prod_{k=1}^q (A_{p_{I(k)}}^U)^{w_k}}{\prod_{k=1}^q (2-A_{p_{I(k)}}^U)^{w_k} + \prod_{k=1}^q (A_{p_{I(k)}}^U)^{w_k}} \right] \right\}, \left\{ \left[\frac{\prod_{k=1}^q (1+A_{p_{F(k)}}^L)^{w_k} - \prod_{k=1}^q (1-A_{p_{F(k)}}^L)^{w_k}}{\prod_{k=1}^q (1+A_{p_{F(k)}}^L)^{w_k} + \prod_{k=1}^q (1-A_{p_{F(k)}}^L)^{w_k}}, \frac{\prod_{k=1}^q (1+A_{p_{F(k)}}^U)^{w_k} - \prod_{k=1}^q (1-A_{p_{F(k)}}^U)^{w_k}}{\prod_{k=1}^q (1+A_{p_{F(k)}}^U)^{w_k} + \prod_{k=1}^q (1-A_{p_{F(k)}}^U)^{w_k}} \right], \left[\frac{2 \prod_{k=1}^q (\lambda_{p_{T(k)}})^{w_k}}{\prod_{k=1}^q (2-\lambda_{p_{T(k)}})^{w_k} + \prod_{k=1}^q (\lambda_{p_{T(k)}})^{w_k}} \right], \left[\frac{2 \prod_{k=1}^q (\lambda_{p_{I(k)}})^{w_k}}{\prod_{k=1}^q (2-\lambda_{p_{I(k)}})^{w_k} + \prod_{k=1}^q (\lambda_{p_{I(k)}})^{w_k}} \right], \left[\frac{\prod_{k=1}^q (1+\lambda_{p_{F(k)}})^{w_k} - \prod_{k=1}^q (1-\lambda_{p_{F(k)}})^{w_k}}{\prod_{k=1}^q (1+\lambda_{p_{F(k)}})^{w_k} + \prod_{k=1}^q (1-\lambda_{p_{F(k)}})^{w_k}} \right] \right\} \right\rangle$$

Using assumption, we have

$$NCHFEGW(\alpha_1, \dots, \alpha_{q+1}) = \bigotimes_E \left(\alpha_k^E \right)^{w_k} \bigotimes_E \left(\alpha_{q+1}^E \right)^{w_{q+1}}$$

$$= \left\{ \left[\frac{2 \prod_{k=1}^q \left(A_{p_T(k)}^L \right)^{w_k}}{\prod_{k=1}^q \left(2 - A_{p_T(k)}^L \right)^{w_k} + \prod_{k=1}^q \left(A_{p_T(k)}^L \right)^{w_k}}, \frac{2 \prod_{k=1}^q \left(A_{p_T(k)}^U \right)^{w_k}}{\prod_{k=1}^q \left(2 - A_{p_T(k)}^U \right)^{w_k} + \prod_{k=1}^q \left(A_{p_T(k)}^U \right)^{w_k}} \right], \right. \\ \left. \left[\frac{2 \prod_{k=1}^q \left(A_{p_I(k)}^L \right)^{w_k}}{\prod_{k=1}^q \left(2 - A_{p_I(k)}^L \right)^{w_k} + \prod_{k=1}^q \left(A_{p_I(k)}^L \right)^{w_k}}, \frac{2 \prod_{k=1}^q \left(A_{p_I(k)}^U \right)^{w_k}}{\prod_{k=1}^q \left(2 - A_{p_I(k)}^U \right)^{w_k} + \prod_{k=1}^q \left(A_{p_I(k)}^U \right)^{w_k}} \right], \right. \\ \left. \left[\frac{\prod_{k=1}^q \left(1 + A_{p_F(k)}^L \right)^{w_k} - \prod_{k=1}^q \left(1 - A_{p_F(k)}^L \right)^{w_k}}{\prod_{k=1}^q \left(1 + A_{p_F(k)}^L \right)^{w_k} + \prod_{k=1}^q \left(1 - A_{p_F(k)}^L \right)^{w_k}}, \frac{\prod_{k=1}^q \left(1 + A_{p_F(k)}^U \right)^{w_k} - \prod_{k=1}^q \left(1 - A_{p_F(k)}^U \right)^{w_k}}{\prod_{k=1}^q \left(1 + A_{p_F(k)}^U \right)^{w_k} + \prod_{k=1}^q \left(1 - A_{p_F(k)}^U \right)^{w_k}} \right] \right\} \\ \otimes_E \left\{ \frac{2 \prod_{k=1}^q \left(\lambda_{p_T(k)} \right)^{w_k}}{\prod_{k=1}^q \left(2 - \lambda_{p_T(k)} \right)^{w_k} + \prod_{k=1}^q \left(\lambda_{p_T(k)} \right)^{w_k}}, \right. \\ \left. \frac{2 \prod_{k=1}^q \left(\lambda_{p_I(k)} \right)^{w_k}}{\prod_{k=1}^q \left(2 - \lambda_{p_I(k)} \right)^{w_k} + \prod_{k=1}^q \left(\lambda_{p_I(k)} \right)^{w_k}}, \right. \\ \left. \frac{\prod_{k=1}^q \left(1 + \lambda_{p_F(k)} \right)^{w_k} - \prod_{k=1}^q \left(1 - \lambda_{p_F(k)} \right)^{w_k}}{\prod_{k=1}^q \left(1 + \lambda_{p_F(k)} \right)^{w_k} + \prod_{k=1}^q \left(1 - \lambda_{p_F(k)} \right)^{w_k}} \right\} \\ \left. \left[\frac{2 \left(A_{p_T}^L \right)^{w_{q+1}}}{\left(2 - A_{p_T}^L \right)^{w_{q+1}} + \left(A_{p_T}^L \right)^{w_{q+1}}}, \frac{2 \left(A_{p_T}^U \right)^{w_{q+1}}}{\left(2 - A_{p_T}^U \right)^{w_{q+1}} + \left(A_{p_T}^U \right)^{w_{q+1}}}, \right. \right. \\ \left. \left[\frac{2 \left(A_{p_I}^L \right)^{w_{q+1}}}{\left(2 - A_{p_I}^L \right)^{w_{q+1}} + \left(A_{p_I}^L \right)^{w_{q+1}}}, \frac{2 \left(A_{p_I}^U \right)^{w_{q+1}}}{\left(2 - A_{p_I}^U \right)^{w_{q+1}} + \left(A_{p_I}^U \right)^{w_{q+1}}}, \right. \right. \\ \left. \left[\frac{\left(1 + A_{p_F}^L \right)^{w_{q+1}} - \left(1 - A_{p_F}^L \right)^{w_{q+1}}}{\left(1 + A_{p_F}^L \right)^{w_{q+1}} + \left(1 - A_{p_F}^L \right)^{w_{q+1}}}, \frac{\left(1 + A_{p_F}^U \right)^{w_{q+1}} - \left(1 - A_{p_F}^U \right)^{w_{q+1}}}{\left(1 + A_{p_F}^U \right)^{w_{q+1}} + \left(1 - A_{p_F}^U \right)^{w_{q+1}}}, \right. \right. \\ \left. \left[\frac{2 \left(\lambda_{p_T} \right)^{w_{q+1}}}{\left(2 - \lambda_{p_T} \right)^{w_{q+1}} + \left(\lambda_{p_T} \right)^{w_{q+1}}}, \right. \right. \\ \left. \left[\frac{2 \left(\lambda_{p_I} \right)^{w_{q+1}}}{\left(2 - \lambda_{p_I} \right)^{w_{q+1}} + \left(\lambda_{p_I} \right)^{w_{q+1}}}, \right. \right. \\ \left. \left[\frac{\left(1 + \lambda_{p_F} \right)^{w_{q+1}} - \left(1 - \lambda_{p_F} \right)^{w_{q+1}}}{\left(1 + \lambda_{p_F} \right)^{w_{q+1}} + \left(1 - \lambda_{p_F} \right)^{w_{q+1}}} \right] \right] \right\}$$

$$= \left\{ \left[\frac{2 \prod_{k=1}^{q+1} (A_{pT(k)}^L)^{w_k}}{\prod_{k=1}^{q+1} (2 - A_{pT(k)}^L) + \prod_{k=1}^{q+1} (A_{pT(k)}^L)^{w_k}}, \frac{2 \prod_{k=1}^{q+1} (A_{pT(k)}^U)^{w_k}}{\prod_{k=1}^{q+1} (2 - A_{pT(k)}^U) + \prod_{k=1}^{q+1} (A_{pT(k)}^U)^{w_k}} \right], \right. \\ \left. \left[\frac{2 \prod_{k=1}^{q+1} (A_{pI(k)}^L)^{w_k}}{\prod_{k=1}^{q+1} (2 - A_{pI(k)}^L) + \prod_{k=1}^{q+1} (A_{pI(k)}^L)^{w_k}}, \frac{2 \prod_{k=1}^{q+1} (A_{pI(k)}^U)^{w_k}}{\prod_{k=1}^{q+1} (2 - A_{pI(k)}^U) + \prod_{k=1}^{q+1} (A_{pI(k)}^U)^{w_k}} \right], \right. \\ \left. \left[\frac{\prod_{k=1}^{q+1} (1 + A_{pF(k)}^L)^{w_k} - \prod_{k=1}^{q+1} (1 - A_{pF(k)}^L)^{w_k}}{\prod_{k=1}^{q+1} (1 + A_{pF(k)}^L) + \prod_{k=1}^{q+1} (1 - A_{pF(k)}^L)^{w_k}}, \frac{\prod_{k=1}^{q+1} (1 + A_{pF(k)}^U)^{w_k} - \prod_{k=1}^{q+1} (1 - A_{pF(k)}^U)^{w_k}}{\prod_{k=1}^{q+1} (1 + A_{pF(k)}^U) + \prod_{k=1}^{q+1} (1 - A_{pF(k)}^U)^{w_k}} \right], \right. \\ \left. \left[\frac{2 \prod_{k=1}^{q+1} (\lambda_{pT(k)})^{w_k}}{\prod_{k=1}^{q+1} (2 - \lambda_{pT(k)}) + \prod_{k=1}^{q+1} (\lambda_{pT(k)})^{w_k}} \right], \right. \\ \left. \left[\frac{2 \prod_{k=1}^{q+1} (\lambda_{pI(k)})^{w_k}}{\prod_{k=1}^{q+1} (2 - \lambda_{pI(k)}) + \prod_{k=1}^{q+1} (\lambda_{pI(k)})^{w_k}} \right], \right. \\ \left. \left[\frac{\prod_{k=1}^{q+1} (1 + \lambda_{pF(k)})^{w_k} - \prod_{k=1}^{q+1} (1 - \lambda_{pF(k)})^{w_k}}{\prod_{k=1}^{q+1} (1 + \lambda_{pF(k)}) + \prod_{k=1}^{q+1} (\lambda_{pF(k)})^{w_k}} \right] \right\}$$

Definition4.8: Neutrosophic cubic hesitant fuzzy hybrid geometric operator (NCHFHG) is a mapping defined

as $NCHFHG(\alpha_1, \dots, \alpha_m) = \bigotimes_{j=1}^m (\alpha_{\sigma(j)})^{w_j}$, where $\alpha_{\sigma(j)} = (\alpha_j)^{mw_j}$ is the j th largest value, m is the

balancing coefficient and $w = (w_1, \dots, w_m)^t$ is the weighting vector.

Theorem 4.9: Let $\{\alpha_{(k)} = \langle A_{(k)}, \lambda_{(k)} \rangle\}$ the set of neutrosophic cubic hesitant fuzzy values with corresponding weight vector $w = (w_1, \dots)^t$, then

$$NCHFHG(\alpha_1, \dots, \alpha_m) = \left\{ \left[\left[\bigotimes_{k=1}^m (A_{jT\sigma(k)}^L)^{w_k}, \bigotimes_{k=1}^m (A_{jT\sigma(k)}^U)^{w_k} \right], \left[\bigotimes_{k=1}^m (A_{jI\sigma(k)}^L)^{w_k}, \bigotimes_{k=1}^m (A_{jI\sigma(k)}^U)^{w_k} \right] \right\}, \\ \left\{ \left[1 - \bigotimes_{k=1}^m (1 - A_{jF\sigma(k)}^L)^{w_k}, 1 - \bigotimes_{k=1}^m (1 - A_{jF\sigma(k)}^U)^{w_k} \right], \left[\bigotimes_{k=1}^m (\lambda_{jT\sigma(k)})^{w_k} \right], \right. \\ \left. \left[\bigotimes_{k=1}^m (\lambda_{jI\sigma(k)})^{w_k} \right], \left[1 - \bigotimes_{k=1}^m (1 - \lambda_{jF\sigma(k)})^{w_k} \right] \right\}$$

Furthermore $NCHFHG(\alpha_1, \alpha_2, \dots, \alpha_m)$ is also a neutrosophic cubic hesitant fuzzy value.

Proof: Using induction for $m=2$

$$NCHFHG(\alpha_1, \alpha_2) = \bigotimes_{k=1}^2 (\alpha_{\sigma(k)})^{w_k}$$

$$\begin{aligned}
 &= \left\langle \left\{ \left[\left(A_{jT\sigma(1)}^L \right)^{w_1}, \left(A_{jT\sigma(1)}^U \right)^{w_1} \right] \right\}, \left\{ \left[\left(A_{jI\sigma(1)}^L \right)^{w_1}, \left(A_{jI\sigma(1)}^U \right)^{w_1} \right] \right\}, \right. \\
 &\quad \left. \left\{ 1 - \left(1 - A_{jF\sigma(1)}^L \right)^{w_1}, 1 - \left(1 - A_{jF\sigma(1)}^U \right)^{w_1} \right\}, \left\{ \left(\lambda_{jT\sigma(1)} \right)^{w_1} \right\}, \right. \\
 &\quad \left. \left\{ \left(\lambda_{jI\sigma(1)} \right)^{w_1} \right\}, \left\{ 1 - \left(1 - \lambda_{jF\sigma(1)} \right)^{w_1} \right\} \right\rangle \otimes \left\langle \left\{ \left[\left(A_{jT\sigma(2)}^L \right)^{w_2}, \left(A_{jT\sigma(2)}^U \right)^{w_2} \right] \right\}, \left\{ \left[\left(A_{jI\sigma(2)}^L \right)^{w_2}, \left(A_{jI\sigma(2)}^U \right)^{w_2} \right] \right\}, \right. \\
 &\quad \left. \left\{ 1 - \left(1 - A_{jF\sigma(2)}^L \right)^{w_2}, 1 - \left(1 - A_{jF\sigma(2)}^U \right)^{w_2} \right\}, \left\{ \left(\lambda_{jT\sigma(2)} \right)^{w_2} \right\}, \right. \\
 &\quad \left. \left\{ \left(\lambda_{jI\sigma(2)} \right)^{w_2} \right\}, \left\{ 1 - \left(1 - \lambda_{jF\sigma(2)} \right)^{w_2} \right\} \right\rangle \\
 &= \left\langle \left\{ \left[\left(\bigotimes_{k=1}^2 A_{jT\sigma(k)}^L \right)^{w_k}, \left(\bigotimes_{k=1}^2 A_{jT\sigma(k)}^U \right)^{w_k} \right] \right\}, \left\{ \left[\left(\bigotimes_{k=1}^2 A_{jI\sigma(k)}^L \right)^{w_k}, \left(\bigotimes_{k=1}^2 A_{jI\sigma(k)}^U \right)^{w_k} \right] \right\}, \right. \\
 &\quad \left. \left\{ 1 - \left(\bigotimes_{k=1}^2 \left(1 - A_{jF\sigma(k)}^L \right)^{w_k} \right), 1 - \left(\bigotimes_{k=1}^2 \left(1 - A_{jF\sigma(k)}^U \right)^{w_k} \right) \right\}, \left\{ \left(\bigotimes_{k=1}^2 \lambda_{jT\sigma(k)} \right)^{w_k} \right\}, \right. \\
 &\quad \left. \left\{ \left(\bigotimes_{k=1}^2 \lambda_{jI\sigma(k)} \right)^{w_k} \right\}, \left\{ 1 - \left(\bigotimes_{k=1}^2 \left(1 - \lambda_{jF\sigma(k)} \right)^{w_k} \right) \right\} \right\rangle
 \end{aligned}$$

For $m = q$ we have

$$NCHFHG(\alpha_1, \dots, \alpha_q) = \left\langle \left\{ \left[\left(\bigotimes_{k=1}^q A_{jT\sigma(k)}^L \right)^{w_k}, \left(\bigotimes_{k=1}^q A_{jT\sigma(k)}^U \right)^{w_k} \right] \right\}, \left\{ \left[\left(\bigotimes_{k=1}^q A_{jI\sigma(k)}^L \right)^{w_k}, \left(\bigotimes_{k=1}^q A_{jI\sigma(k)}^U \right)^{w_k} \right] \right\}, \right. \\
 \left. \left\{ 1 - \left(\bigotimes_{k=1}^q \left(1 - A_{jF\sigma(k)}^L \right)^{w_k} \right), 1 - \left(\bigotimes_{k=1}^q \left(1 - A_{jF\sigma(k)}^U \right)^{w_k} \right) \right\}, \left\{ \left(\bigotimes_{k=1}^q \lambda_{jT\sigma(k)} \right)^{w_k} \right\}, \right. \\
 \left. \left\{ \left(\bigotimes_{k=1}^q \lambda_{jI\sigma(k)} \right)^{w_k} \right\}, \left\{ 1 - \left(\bigotimes_{k=1}^q \left(1 - \lambda_{jF\sigma(k)} \right)^{w_k} \right) \right\} \right\rangle$$

we prove for $m = q + 1$

$$\begin{aligned}
 \left(\bigotimes_{k=1}^q \alpha_{\sigma(k)} \right)^{w_k} \otimes \left(\alpha_{\sigma(q+1)} \right)^{w_{q+1}} &= \left\langle \left\{ \left[\left(\bigotimes_{k=1}^q A_{jT\sigma(k)}^L \right)^{w_k}, \left(\bigotimes_{k=1}^q A_{jT\sigma(k)}^U \right)^{w_k} \right] \right\}, \left\{ \left[\left(\bigotimes_{k=1}^q A_{jI\sigma(k)}^L \right)^{w_k}, \left(\bigotimes_{k=1}^q A_{jI\sigma(k)}^U \right)^{w_k} \right] \right\}, \right. \\
 &\quad \left. \left\{ 1 - \left(\bigotimes_{k=1}^q \left(1 - A_{jF\sigma(k)}^L \right)^{w_k} \right), 1 - \left(\bigotimes_{k=1}^q \left(1 - A_{jF\sigma(k)}^U \right)^{w_k} \right) \right\}, \left\{ \left(\bigotimes_{k=1}^q \lambda_{jT\sigma(k)} \right)^{w_k} \right\}, \right. \\
 &\quad \left. \left\{ \left(\bigotimes_{k=1}^q \lambda_{jI\sigma(k)} \right)^{w_k} \right\}, \left\{ 1 - \left(\bigotimes_{k=1}^q \left(1 - \lambda_{jF\sigma(k)} \right)^{w_k} \right) \right\} \right\rangle \otimes \left\langle \left\{ \left[\left(A_{jT\sigma(q+1)}^L \right)^{w_{q+1}}, \left(A_{jT\sigma(q+1)}^U \right)^{w_{q+1}} \right] \right\}, \left\{ \left[\left(A_{jI\sigma(q+1)}^L \right)^{w_{q+1}}, \left(A_{jI\sigma(q+1)}^U \right)^{w_{q+1}} \right] \right\}, \right. \\
 &\quad \left. \left\{ 1 - \left(1 - A_{jF\sigma(q+1)}^L \right)^{w_{q+1}}, 1 - \left(1 - A_{jF\sigma(q+1)}^U \right)^{w_{q+1}} \right\}, \left\{ \left(\lambda_{jT\sigma(q+1)} \right)^{w_{q+1}} \right\}, \right. \\
 &\quad \left. \left\{ \left(\lambda_{jI\sigma(q+1)} \right)^{w_{q+1}} \right\}, \left\{ 1 - \left(1 - \lambda_{jF\sigma(q+1)} \right)^{w_{q+1}} \right\} \right\rangle
 \end{aligned}$$

Theorem 4.10: With $w_j = \frac{1}{m}$, $NCHFHG$ becomes $NCHFVG$.

Proof:
$$NCHFHG(\alpha_1, \dots, \alpha_m) = (\alpha_{\sigma(1)})^{w_1} \otimes \dots \otimes (\alpha_{\sigma(m)})^{w_m}$$

$$= (\alpha_1)^{w_1} \otimes \dots \otimes (\alpha_m)^{w_m}$$

$$= NCHFVG(\alpha_1, \dots, \alpha_m)$$

Definition 4.11: Neutrosophic cubic hesitant fuzzy Einstein hybrid geometric operator(NCHFHEG) is a

mapping defined as $NCHFHEG(\alpha_1, \dots, \alpha_m) = \bigotimes_E^m \left(\alpha_{\sigma(j)} \right)^{w_j}$ where $\alpha_{\sigma(j)} = (\alpha_j)^{m w_j}$ is the j th largest value, m is the balancing coefficient and $w = (w_1, \dots, w_m)^t$ is the weighting vector.

Theorem 4.12: Let $\{\alpha_{(k)} = \langle A_{(k)}, \lambda_{(k)} \rangle\}$ the set of neutrosophic cubic hesitant fuzzy values with corresponding weight vector $w = (w_1, \dots)^t$ then

$$NCHFHEG(\alpha_1, \alpha_2, \dots, \alpha_m) = \left\{ \frac{\left[\frac{2 \otimes_{k=1}^m \left(A_{jT\sigma(k)}^L \right)^{w_k}}{m \otimes_{k=1}^m \left(2 - A_{jT\sigma(k)}^L \right)^{w_k} + \otimes_{k=1}^m \left(A_{jT\sigma(k)}^L \right)^{w_k}}, \frac{2 \otimes_{k=1}^m \left(A_{jT\sigma(k)}^U \right)^{w_k}}{m \otimes_{k=1}^m \left(2 - A_{jT\sigma(k)}^U \right)^{w_k} + \otimes_{k=1}^m \left(A_{jT\sigma(k)}^U \right)^{w_k}} \right\}, \left\{ \frac{2 \otimes_{k=1}^m \left(A_{jI\sigma(k)}^L \right)^{w_k}}{m \otimes_{k=1}^m \left(2 - A_{jI\sigma(k)}^L \right)^{w_k} + \otimes_{k=1}^m \left(A_{jI\sigma(k)}^L \right)^{w_k}}, \frac{2 \otimes_{k=1}^m \left(A_{jI\sigma(k)}^U \right)^{w_k}}{m \otimes_{k=1}^m \left(2 - A_{jI\sigma(k)}^U \right)^{w_k} + \otimes_{k=1}^m \left(A_{jI\sigma(k)}^U \right)^{w_k}} \right\}, \left\{ \frac{m \otimes_{k=1}^m \left(1 + A_{jF\sigma(k)}^L \right)^{w_k} - m \otimes_{k=1}^m \left(1 - A_{jF\sigma(k)}^L \right)^{w_k}}{m \otimes_{k=1}^m \left(1 + A_{jF\sigma(k)}^L \right)^{w_k} + \otimes_{k=1}^m \left(1 - A_{jF\sigma(k)}^L \right)^{w_k}}, \frac{m \otimes_{k=1}^m \left(1 + A_{jF\sigma(k)}^U \right)^{w_k} - m \otimes_{k=1}^m \left(1 - A_{jF\sigma(k)}^U \right)^{w_k}}{m \otimes_{k=1}^m \left(1 + A_{jF\sigma(k)}^U \right)^{w_k} + \otimes_{k=1}^m \left(1 - A_{jF\sigma(k)}^U \right)^{w_k}} \right\}, \left\{ \frac{2 \otimes_{k=1}^m \left(1 + \lambda_{jT\sigma(k)} \right)^{w_k}}{m \otimes_{k=1}^m \left(2 - \lambda_{jT\sigma(k)} \right)^{w_k} + \otimes_{k=1}^m \left(1 + \lambda_{jT\sigma(k)} \right)^{w_k}}, \frac{2 \otimes_{k=1}^m \left(1 + \lambda_{jI\sigma(k)} \right)^{w_k}}{m \otimes_{k=1}^m \left(2 - \lambda_{jI\sigma(k)} \right)^{w_k} + \otimes_{k=1}^m \left(1 + \lambda_{jI\sigma(k)} \right)^{w_k}} \right\}, \left\{ \frac{m \otimes_{k=1}^m \left(1 + \lambda_{jF\sigma(k)} \right)^{w_k} - m \otimes_{k=1}^m \left(1 - \lambda_{jF\sigma(k)} \right)^{w_k}}{m \otimes_{k=1}^m \left(1 + \lambda_{jF\sigma(k)} \right)^{w_k} + \otimes_{k=1}^m \left(1 - \lambda_{jF\sigma(k)} \right)^{w_k}} \right\}$$

Proof: Using induction and from Theorem 3.12

$$\alpha^{E^{w_1}} = \left\langle \left[\left[\frac{2(A_{j_T}^L)^{w_1}}{(2-A_{j_T}^L)^{w_1} + (A_{j_T}^L)^{w_1}}, \frac{2(A_{j_T}^U)^{w_1}}{(2-A_{j_T}^U)^{w_1} + (A_{j_T}^U)^{w_1}} \right], \left[\frac{2(A_{j_I}^L)^{w_1}}{(2-A_{j_I}^L)^{w_1} + (A_{j_I}^L)^{w_1}}, \frac{2(A_{j_I}^U)^{w_1}}{(2-A_{j_I}^U)^{w_1} + (A_{j_I}^U)^{w_1}} \right] \right], \left[\left[\frac{(1+A_{j_F}^L)^{w_1} - (1-A_{j_F}^L)^{w_1}}{(1+A_{j_F}^L)^{w_1} + (1-A_{j_F}^L)^{w_1}}, \frac{(1+A_{j_F}^U)^{w_1} - (1-A_{j_F}^U)^{w_1}}{(1+A_{j_F}^U)^{w_1} + (1-A_{j_F}^U)^{w_1}} \right], \left[\frac{2(\lambda_{j_T})^{w_1}}{(2-\lambda_{j_T})^{w_1} + (\lambda_{j_T})^{w_1}} \right], \left[\frac{2(\lambda_{j_I})^{w_1}}{(2-\lambda_{j_I})^{w_1} + (\lambda_{j_I})^{w_1}} \right], \left[\frac{2(\lambda_{j_F})^{w_1}}{(2-\lambda_{j_F})^{w_1} + (\lambda_{j_F})^{w_1}} \right], \left[\frac{(1+\lambda_{j_F})^{w_1} - (1-\lambda_{j_F})^{w_1}}{(1+\lambda_{j_F})^{w_1} + (1-\lambda_{j_F})^{w_1}} \right] \right] \right\rangle$$

$$\text{and } \alpha^{E^{w_2}} = \left\langle \left[\left[\frac{2(A_{j_T}^L)^{w_2}}{(2-A_{j_T}^L)^{w_2} + (A_{j_T}^L)^{w_2}}, \frac{2(A_{j_T}^U)^{w_2}}{(2-A_{j_T}^U)^{w_2} + (A_{j_T}^U)^{w_2}} \right], \left[\frac{2(A_{j_I}^L)^{w_2}}{(2-A_{j_I}^L)^{w_2} + (A_{j_I}^L)^{w_2}}, \frac{2(A_{j_I}^U)^{w_2}}{(2-A_{j_I}^U)^{w_2} + (A_{j_I}^U)^{w_2}} \right] \right], \left[\left[\frac{(1+A_{j_F}^L)^{w_2} - (1-A_{j_F}^L)^{w_2}}{(1+A_{j_F}^L)^{w_2} + (1-A_{j_F}^L)^{w_2}}, \frac{(1+A_{j_F}^U)^{w_2} - (1-A_{j_F}^U)^{w_2}}{(1+A_{j_F}^U)^{w_2} + (1-A_{j_F}^U)^{w_2}} \right], \left[\frac{2(\lambda_{j_T})^{w_2}}{(2-\lambda_{j_T})^{w_2} + (\lambda_{j_T})^{w_2}} \right], \left[\frac{2(\lambda_{j_I})^{w_2}}{(2-\lambda_{j_I})^{w_2} + (\lambda_{j_I})^{w_2}} \right], \left[\frac{2(\lambda_{j_F})^{w_2}}{(2-\lambda_{j_F})^{w_2} + (\lambda_{j_F})^{w_2}} \right], \left[\frac{(1+\lambda_{j_F})^{w_2} - (1-\lambda_{j_F})^{w_2}}{(1+\lambda_{j_F})^{w_2} + (1-\lambda_{j_F})^{w_2}} \right] \right] \right\rangle$$

$$NCHFEG(\alpha_1, \alpha_2) = \bigotimes_{k=1}^2 \alpha_k^{E^{w_k}}$$

$$= \left\langle \left[\left[\frac{2 \bigotimes_{k=1}^2 \left(A_{j_T \sigma(k)}^L \right)^{w_k}}{2 \bigotimes_{k=1}^2 \left(2 - A_{j_T \sigma(k)}^L \right)^{w_k} + \bigotimes_{k=1}^2 \left(A_{j_T \sigma(k)}^L \right)^{w_k}}, \frac{2 \bigotimes_{k=1}^2 \left(A_{j_T \sigma(k)}^U \right)^{w_k}}{2 \bigotimes_{k=1}^2 \left(2 - A_{j_T \sigma(k)}^U \right)^{w_k} + \bigotimes_{k=1}^2 \left(A_{j_T \sigma(k)}^U \right)^{w_k}} \right], \left[\frac{2 \bigotimes_{k=1}^2 \left(A_{j_I \sigma(k)}^L \right)^{w_k}}{2 \bigotimes_{k=1}^2 \left(2 - A_{j_I \sigma(k)}^L \right)^{w_k} + \bigotimes_{k=1}^2 \left(A_{j_I \sigma(k)}^L \right)^{w_k}}, \frac{2 \bigotimes_{k=1}^2 \left(A_{j_I \sigma(k)}^U \right)^{w_k}}{2 \bigotimes_{k=1}^2 \left(2 - A_{j_I \sigma(k)}^U \right)^{w_k} + \bigotimes_{k=1}^2 \left(A_{j_I \sigma(k)}^U \right)^{w_k}} \right] \right], \left[\left[\frac{2 \bigotimes_{k=1}^2 \left(1 + A_{j_F \sigma(k)}^L \right)^{w_k} - 2 \bigotimes_{k=1}^2 \left(1 - A_{j_F \sigma(k)}^L \right)^{w_k}}{2 \bigotimes_{k=1}^2 \left(1 + A_{j_F \sigma(k)}^L \right)^{w_k} + \bigotimes_{k=1}^2 \left(1 - A_{j_F \sigma(k)}^L \right)^{w_k}}, \frac{2 \bigotimes_{k=1}^2 \left(1 + A_{j_F \sigma(k)}^U \right)^{w_k} - 2 \bigotimes_{k=1}^2 \left(1 - A_{j_F \sigma(k)}^U \right)^{w_k}}{2 \bigotimes_{k=1}^2 \left(1 + A_{j_F \sigma(k)}^U \right)^{w_k} + \bigotimes_{k=1}^2 \left(1 - A_{j_F \sigma(k)}^U \right)^{w_k}} \right], \left[\frac{2 \bigotimes_{k=1}^2 \left(\lambda_{j_T \sigma(k)} \right)^{w_k}}{2 \bigotimes_{k=1}^2 \left(2 - \lambda_{j_T \sigma(k)} \right)^{w_k} + \bigotimes_{k=1}^2 \left(\lambda_{j_T \sigma(k)} \right)^{w_k}} \right], \left[\frac{2 \bigotimes_{k=1}^2 \left(\lambda_{j_I \sigma(k)} \right)^{w_k}}{2 \bigotimes_{k=1}^2 \left(2 - \lambda_{j_I \sigma(k)} \right)^{w_k} + \bigotimes_{k=1}^2 \left(\lambda_{j_I \sigma(k)} \right)^{w_k}} \right], \left[\frac{2 \bigotimes_{k=1}^2 \left(\lambda_{j_F \sigma(k)} \right)^{w_k}}{2 \bigotimes_{k=1}^2 \left(2 - \lambda_{j_F \sigma(k)} \right)^{w_k} + \bigotimes_{k=1}^2 \left(\lambda_{j_F \sigma(k)} \right)^{w_k}} \right], \left[\frac{2 \bigotimes_{k=1}^2 \left(1 + \lambda_{j_F \sigma(k)} \right)^{w_k} - 2 \bigotimes_{k=1}^2 \left(1 - \lambda_{j_F \sigma(k)} \right)^{w_k}}{2 \bigotimes_{k=1}^2 \left(1 + \lambda_{j_F \sigma(k)} \right)^{w_k} + \bigotimes_{k=1}^2 \left(1 - \lambda_{j_F \sigma(k)} \right)^{w_k}} \right] \right] \right\rangle$$

For $m = q$ we have

$$NCHFEG(\alpha_1, \alpha_2, \dots, \alpha_q) = \left\{ \left[\begin{array}{c} \left[\begin{array}{c} 2 \otimes_{k=1}^q \left(A_{j_T \sigma(k)}^L \right)^{w_k} \\ \left[\begin{array}{c} \frac{q}{k=1} \left(2 - A_{j_T \sigma(k)}^L \right)^{w_k} + \frac{q}{k=1} \left(A_{j_T \sigma(k)}^L \right)^{w_k} \\ \frac{q}{k=1} \left(2 - A_{j_T \sigma(k)}^U \right)^{w_k} + \frac{q}{k=1} \left(A_{j_T \sigma(k)}^U \right)^{w_k} \end{array} \right] \\ \left[\begin{array}{c} 2 \otimes_{k=1}^q \left(A_{j_I \sigma(k)}^L \right)^{w_k} \\ \frac{q}{k=1} \left(2 - A_{j_I \sigma(k)}^L \right)^{w_k} + \frac{q}{k=1} \left(A_{j_I \sigma(k)}^L \right)^{w_k} \end{array} \right] \\ \left[\begin{array}{c} 2 \otimes_{k=1}^m \left(A_{j_I \sigma(k)}^U \right)^{w_k} \\ \frac{q}{k=1} \left(2 - A_{j_I \sigma(k)}^U \right)^{w_k} + \frac{q}{k=1} \left(A_{j_I \sigma(k)}^U \right)^{w_k} \end{array} \right] \\ \left[\begin{array}{c} \frac{q}{k=1} \left(1 + A_{j_F \sigma(k)}^L \right)^{w_k} - \frac{q}{k=1} \left(1 - A_{j_F \sigma(k)}^L \right)^{w_k} \\ \frac{q}{k=1} \left(1 + A_{j_F \sigma(k)}^L \right)^{w_k} + \frac{q}{k=1} \left(1 - A_{j_F \sigma(k)}^L \right)^{w_k} \end{array} \right] \\ \left[\begin{array}{c} \frac{q}{k=1} \left(1 + A_{j_F \sigma(k)}^U \right)^{w_k} - \frac{q}{k=1} \left(1 - A_{j_F \sigma(k)}^U \right)^{w_k} \\ \frac{q}{k=1} \left(1 + A_{j_F \sigma(k)}^U \right)^{w_k} + \frac{q}{k=1} \left(1 - A_{j_F \sigma(k)}^U \right)^{w_k} \end{array} \right] \\ \left[\begin{array}{c} 2 \otimes_{k=1}^q \left(\lambda_{j_T \sigma(k)} \right)^{w_k} \\ \frac{q}{k=1} \left(2 - \lambda_{j_T \sigma(k)} \right)^{w_k} + \frac{q}{k=1} \left(\lambda_{j_T \sigma(k)} \right)^{w_k} \end{array} \right] \\ \left[\begin{array}{c} 2 \otimes_{k=1}^q \left(\lambda_{j_I \sigma(k)} \right)^{w_k} \\ \frac{q}{k=1} \left(2 - \lambda_{j_I \sigma(k)} \right)^{w_k} + \frac{q}{k=1} \left(\lambda_{j_I \sigma(k)} \right)^{w_k} \end{array} \right] \\ \left[\begin{array}{c} \frac{q}{k=1} \left(1 + \lambda_{j_F \sigma(k)} \right)^{w_k} - \frac{q}{k=1} \left(1 - \lambda_{j_F \sigma(k)} \right)^{w_k} \\ \frac{q}{k=1} \left(1 + \lambda_{j_F \sigma(k)} \right)^{w_k} + \frac{q}{k=1} \left(1 - \lambda_{j_F \sigma(k)} \right)^{w_k} \end{array} \right] \\ \left[\begin{array}{c} \frac{q}{k=1} \left(1 + \lambda_{j_F \sigma(k)} \right)^{w_k} - \frac{q}{k=1} \left(1 - \lambda_{j_F \sigma(k)} \right)^{w_k} \\ \frac{q}{k=1} \left(1 + \lambda_{j_F \sigma(k)} \right)^{w_k} + \frac{q}{k=1} \left(1 - \lambda_{j_F \sigma(k)} \right)^{w_k} \end{array} \right] \end{array} \right] \right\}$$

Using assumption, we have

$$\left(\bigotimes_{k=1}^q \left(\alpha_{\sigma(k)}^E \right)^{w_k} \right) \otimes_E \left(\alpha_{q+1}^E \right)^{w_{q+1}} = \left[\begin{array}{c} \left\{ \frac{2 \bigotimes_{k=1}^q \left(A_{J_T \sigma(k)}^L \right)^{w_k}}{\left(\frac{q \bigotimes_{k=1}^q \left(2 - A_{J_T \sigma(k)}^L \right)^{w_k} + \bigotimes_{k=1}^q \left(A_{J_T \sigma(k)}^L \right)^{w_k}}{2 \bigotimes_{k=1}^q \left(A_{J_T \sigma(k)}^U \right)^{w_k}} \right\}} \\ \left\{ \frac{q \bigotimes_{k=1}^q \left(2 - A_{J_T \sigma(k)}^U \right)^{w_k} + \bigotimes_{k=1}^q \left(A_{J_T \sigma(k)}^U \right)^{w_k}}{2 \bigotimes_{k=1}^q \left(A_{J_T \sigma(k)}^L \right)^{w_k}} \right\}} \\ \left\{ \frac{q \bigotimes_{k=1}^q \left(2 - A_{J_T \sigma(k)}^U \right)^{w_k} + \bigotimes_{k=1}^q \left(A_{J_T \sigma(k)}^U \right)^{w_k}}{2 \bigotimes_{k=1}^q \left(A_{J_T \sigma(k)}^L \right)^{w_k}} \right\}} \\ \left\{ \frac{q \bigotimes_{k=1}^q \left(2 - A_{J_I \sigma(k)}^L \right)^{w_k} + \bigotimes_{k=1}^q \left(A_{J_I \sigma(k)}^L \right)^{w_k}}{2 \bigotimes_{k=1}^q \left(A_{J_I \sigma(k)}^U \right)^{w_k}} \right\}} \\ \left\{ \frac{q \bigotimes_{k=1}^q \left(2 - A_{J_I \sigma(k)}^L \right)^{w_k} + \bigotimes_{k=1}^q \left(A_{J_I \sigma(k)}^L \right)^{w_k}}{2 \bigotimes_{k=1}^q \left(A_{J_I \sigma(k)}^U \right)^{w_k}} \right\}} \\ \left\{ \frac{q \bigotimes_{k=1}^q \left(2 - A_{J_I \sigma(k)}^U \right)^{w_k} + \bigotimes_{k=1}^q \left(A_{J_I \sigma(k)}^U \right)^{w_k}}{2 \bigotimes_{k=1}^q \left(A_{J_I \sigma(k)}^L \right)^{w_k}} \right\}} \\ \left\{ \frac{q \bigotimes_{k=1}^q \left(1 + A_{J_F \sigma(k)}^L \right)^{w_k} - \bigotimes_{k=1}^q \left(1 - A_{J_F \sigma(k)}^L \right)^{w_k}}{q \bigotimes_{k=1}^q \left(1 + A_{J_F \sigma(k)}^L \right)^{w_k} + \bigotimes_{k=1}^q \left(1 - A_{J_F \sigma(k)}^L \right)^{w_k}} \right\}} \\ \left\{ \frac{q \bigotimes_{k=1}^q \left(1 + A_{J_F \sigma(k)}^L \right)^{w_k} + \bigotimes_{k=1}^q \left(1 - A_{J_F \sigma(k)}^L \right)^{w_k}}{q \bigotimes_{k=1}^q \left(1 + A_{J_F \sigma(k)}^U \right)^{w_k} - \bigotimes_{k=1}^q \left(1 - A_{J_F \sigma(k)}^U \right)^{w_k}} \right\}} \\ \left\{ \frac{q \bigotimes_{k=1}^q \left(1 + A_{J_F \sigma(k)}^U \right)^{w_k} - \bigotimes_{k=1}^q \left(1 - A_{J_F \sigma(k)}^U \right)^{w_k}}{q \bigotimes_{k=1}^q \left(1 + A_{J_F \sigma(k)}^L \right)^{w_k} + \bigotimes_{k=1}^q \left(1 - A_{J_F \sigma(k)}^L \right)^{w_k}} \right\}} \\ \left\{ \frac{q \bigotimes_{k=1}^q \left(1 + A_{J_F \sigma(k)}^U \right)^{w_k} + \bigotimes_{k=1}^q \left(1 - A_{J_F \sigma(k)}^U \right)^{w_k}}{2 \bigotimes_{k=1}^q \left(\lambda_{j_T \sigma(k)} \right)^{w_k}} \right\}} \\ \left\{ \frac{q \bigotimes_{k=1}^q \left(2 - \lambda_{j_T \sigma(k)} \right)^{w_k} + \bigotimes_{k=1}^q \left(\lambda_{j_T \sigma(k)} \right)^{w_k}}{2 \bigotimes_{k=1}^q \left(\lambda_{j_I \sigma(k)} \right)^{w_k}} \right\}} \\ \left\{ \frac{q \bigotimes_{k=1}^q \left(2 - \lambda_{j_I \sigma(k)} \right)^{w_k} + \bigotimes_{k=1}^q \left(\lambda_{j_I \sigma(k)} \right)^{w_k}}{2 \bigotimes_{k=1}^q \left(\lambda_{j_I \sigma(k)} \right)^{w_k}} \right\}} \\ \left\{ \frac{q \bigotimes_{k=1}^q \left(2 - \lambda_{j_I \sigma(k)} \right)^{w_k} + \bigotimes_{k=1}^q \left(\lambda_{j_I \sigma(k)} \right)^{w_k}}{q \bigotimes_{k=1}^q \left(1 + \lambda_{j_F \sigma(k)} \right)^{w_k} - \bigotimes_{k=1}^q \left(1 - \lambda_{j_F \sigma(k)} \right)^{w_k}} \right\}} \\ \left\{ \frac{q \bigotimes_{k=1}^q \left(1 + \lambda_{j_F \sigma(k)} \right)^{w_k} + \bigotimes_{k=1}^q \left(1 - \lambda_{j_F \sigma(k)} \right)^{w_k}}{q \bigotimes_{k=1}^q \left(1 + \lambda_{j_F \sigma(k)} \right)^{w_k} + \bigotimes_{k=1}^q \left(1 - \lambda_{j_F \sigma(k)} \right)^{w_k}} \right\}} \end{array} \right] \otimes_E \left[\begin{array}{c} \left\{ \frac{2 \left(A_{j_T}^L \right)^{w_{q+1}}}{\left(2 - A_{j_T}^L \right)^{w_{q+1}} + \left(A_{j_T}^L \right)^{w_{q+1}}} \right\}} \\ \left\{ \frac{2 \left(A_{j_T}^U \right)^{w_{q+1}}}{\left(2 - A_{j_T}^U \right)^{w_{q+1}} + \left(A_{j_T}^U \right)^{w_{q+1}}} \right\}} \\ \left\{ \frac{2 \left(A_{j_I}^L \right)^{w_{q+1}}}{\left(2 - A_{j_I}^L \right)^{w_{q+1}} + \left(A_{j_I}^L \right)^{w_{q+1}}} \right\}} \\ \left\{ \frac{2 \left(A_{j_I}^U \right)^{w_{q+1}}}{\left(2 - A_{j_I}^U \right)^{w_{q+1}} + \left(A_{j_I}^U \right)^{w_{q+1}}} \right\}} \\ \left\{ \frac{\left(1 + A_{j_F}^L \right)^{w_{q+1}} - \left(1 - A_{j_F}^L \right)^{w_{q+1}}}{\left(1 + A_{j_F}^L \right)^{w_{q+1}} + \left(1 - A_{j_F}^L \right)^{w_{q+1}}} \right\}} \\ \left\{ \frac{\left(1 + A_{j_F}^L \right)^{w_{q+1}} + \left(1 - A_{j_F}^L \right)^{w_{q+1}}}{\left(1 + A_{j_F}^U \right)^{w_{q+1}} - \left(1 - A_{j_F}^U \right)^{w_{q+1}}} \right\}} \\ \left\{ \frac{\left(1 + A_{j_F}^U \right)^{w_{q+1}} - \left(1 - A_{j_F}^U \right)^{w_{q+1}}}{\left(1 + A_{j_F}^L \right)^{w_{q+1}} + \left(1 - A_{j_F}^L \right)^{w_{q+1}}} \right\}} \\ \left\{ \frac{2 \left(\lambda_{j_T} \right)^{w_{q+1}}}{\left(2 - \lambda_{j_T} \right)^{w_{q+1}} + \left(\lambda_{j_T} \right)^{w_{q+1}}} \right\}} \\ \left\{ \frac{2 \left(\lambda_{j_I} \right)^{w_{q+1}}}{\left(2 - \lambda_{j_I} \right)^{w_{q+1}} + \left(\lambda_{j_I} \right)^{w_{q+1}}} \right\}} \\ \left\{ \frac{\left(1 + \lambda_{j_F} \right)^{w_{q+1}} - \left(1 - \lambda_{j_F} \right)^{w_{q+1}}}{\left(1 + \lambda_{j_F} \right)^{w_{q+1}} + \left(1 - \lambda_{j_F} \right)^{w_{q+1}}} \right\}} \end{array} \right]$$

$$D^{(1)} = \left(\begin{array}{ccc} \left\langle \begin{array}{l} \{[0.1, 0.5], [0.2, 0.7]\}, \\ \{[0.2, 0.3], [0.1, 0.6]\}, \\ \{[0.1, 0.5], [0.2, 0.3]\}, \\ \{0.4, 0.6\}, \{0.3, 0.5\}, \{0.3, 0.4\} \end{array} \right\rangle & \left\langle \begin{array}{l} \{[0.3, 0.4], [0.2, 0.9]\}, \\ \{[0.2, 0.6], [0.3, 0.6]\}, \\ \{[0.3, 0.4], [0, 0.1]\}, \\ \{0.5, 0.7\}, \{0.3, 0.4\}, \{0.4, 0.5\} \end{array} \right\rangle & \left\langle \begin{array}{l} \{[0.3, 0.7], [0.2, 0.4]\}, \\ \{[0.2, 0.5], [0.1, 0.6]\}, \\ \{[0.2, 0.4], [0, 0.1]\}, \\ \{0.5, 0.6\}, \{0.2, 0.4\}, \{0.2, 0.3\} \end{array} \right\rangle \\ \left\langle \begin{array}{l} \{[0.2, 0.5], [0.3, 0.7]\}, \\ \{[0.3, 0.4], [0.1, 0.5]\}, \\ \{[0.1, 0.3], [0, 0.2]\}, \\ \{0.7, 0.9\}, \{0.3, 0.4\}, \{0.1, 0.2\} \end{array} \right\rangle & \left\langle \begin{array}{l} \{[0.1, 0.3], [0.2, 0.5]\}, \\ \{[0.2, 0.3], [0.1, 0.6]\}, \\ \{[0.1, 0.4], [0, 0.3]\}, \\ \{0.4, 0.5\}, \{0.3, 0.7\}, \{0.3, 0.5\} \end{array} \right\rangle & \left\langle \begin{array}{l} \{[0.1, 0.4], [0.2, 0.6]\}, \\ \{[0.1, 0.3], [0, 0.2]\}, \\ \{[0.2, 0.4], [0, 0.3]\}, \\ \{0.4, 0.5\}, \{0.3, 0.6\}, \{0.1, 0.3\} \end{array} \right\rangle \\ \left\langle \begin{array}{l} \{[0.4, 0.5], [0.3, 0.4]\}, \\ \{[0.1, 0.3], [0.2, 0.5]\}, \\ \{[0.1, 0.4], [0.7, 0.8]\}, \\ \{0.4, 0.5\}, \{0.3, 0.4\}, \{0.2, 0.4\} \end{array} \right\rangle & \left\langle \begin{array}{l} \{[0.3, 0.5], [0.1, 0.4]\}, \\ \{[0.2, 0.3], [0.2, 0.6]\}, \\ \{[0.1, 0.5], [0.6, 0.8]\}, \\ \{0.5, 0.6\}, \{0.1, 0.4\}, \{0.2, 0.3\} \end{array} \right\rangle & \left\langle \begin{array}{l} \{[0.2, 0.5], [0.1, 0.4]\}, \\ \{[0.2, 0.5], [0.2, 0.6]\}, \\ \{[0.1, 0.4], [0.6, 0.7]\}, \\ \{0.4, 0.6\}, \{0.3, 0.4\}, \{0.4, 0.5\} \end{array} \right\rangle \end{array} \right)$$

Explanation of decision matrix entries:

In case of $d_{11}^{(1)}$, $\{[0.1, 0.5], [0.2, 0.7]\}$ is interval hesitant degree of preference to attribute F_1 corresponding to attribute K_1 , $\{[0.2, 0.3], [0.1, 0.6]\}$ is interval hesitant degree of indeterminacy (preference/ non-preference) for attribute F_1 corresponding to attribute K_1 , $\{[0.1, 0.5], [0.2, 0.3]\}$ is interval hesitant degree of non-preference for attribute F_1 corresponding to attribute K_1 , $\{0.4, 0.6\}$ is hesitant degree of preference for attribute F_1 corresponding to attribute K_1 , $\{0.3, 0.5\}$ is hesitant degree of indeterminacy (preference/ non-preference) for attribute F_1 corresponding to attribute K_1 , $\{0.3, 0.4\}$ is hesitant degree of non-preference for attribute F_1 corresponding to attribute K_1 , given by the first expert.

Decision Matrix for second expert

$$D^{(2)} = \left(\begin{array}{ccc} \left\langle \begin{array}{l} \{[0.1, 0.3], [0.2, 0.5]\}, \\ \{[0.2, 0.4], [0.1, 0.5]\}, \\ \{[0.1, 0.4], [0, 0.3]\}, \\ \{0.4, 0.8\}, \{0.3, 0.4\}, \{0.3, 0.4\} \end{array} \right\rangle & \left\langle \begin{array}{l} \{[0.2, 0.4], [0.2, 0.7]\}, \\ \{[0.1, 0.3], [0, 0.2]\}, \\ \{[0.3, 0.5], [0, 0.3]\}, \\ \{0.7, 0.8\}, \{0.3, 0.6\}, \{0.2, 0.3\} \end{array} \right\rangle & \left\langle \begin{array}{l} \{[0.2, 0.5], [0.2, 0.7]\}, \\ \{[0, 0.4], [0.1, 0.3]\}, \\ \{[0.3, 0.5], [0, 0.4]\}, \\ \{0.6, 0.7\}, \{0.2, 0.5\}, \{0.4, 0.5\} \end{array} \right\rangle \\ \left\langle \begin{array}{l} \{[0.1, 0.5], [0.3, 0.7]\}, \\ \{[0.1, 0.3], [0.1, 0.2]\}, \\ \{[0.2, 0.4], [0, 0.3]\}, \\ \{0.6, 0.7\}, \{0.3, 0.6\}, \{0.4, 0.5\} \end{array} \right\rangle & \left\langle \begin{array}{l} \{[0.3, 0.6], [0.2, 0.7]\}, \\ \{[0.1, 0.2], [0.2, 0.5]\}, \\ \{[0, 0.3], [0.1, 0.4]\}, \\ \{0.7, 0.8\}, \{0.3, 0.4\}, \{0.2, 0.3\} \end{array} \right\rangle & \left\langle \begin{array}{l} \{[0.4, 0.5], [0.4, 0.7]\}, \\ \{[0.2, 0.3], [0.1, 0.6]\}, \\ \{[0.3, 0.4], [0, 0.3]\}, \\ \{0.5, 0.6\}, \{0.3, 0.5\}, \{0.3, 0.4\} \end{array} \right\rangle \\ \left\langle \begin{array}{l} \{[0.2, 0.5], [0.1, 0.4]\}, \\ \{[0.2, 0.4], [0.2, 0.5]\}, \\ \{[0.1, 0.3], [0.8, 0.9]\}, \\ \{0.7, 0.8\}, \{0.6, 0.8\}, \{0.3, 0.4\} \end{array} \right\rangle & \left\langle \begin{array}{l} \{[0.2, 0.3], [0.4, 0.5]\}, \\ \{[0.1, 0.4], [0.2, 0.5]\}, \\ \{[0.7, 0.8], [0.1, 0.2]\}, \\ \{0.7, 0.9\}, \{0.7, 0.8\}, \{0.2, 0.5\} \end{array} \right\rangle & \left\langle \begin{array}{l} \{[0.5, 0.7], [0.3, 0.4]\}, \\ \{[0.1, 0.3], [0.2, 0.5]\}, \\ \{[0.1, 0.4], [0.2, 0.6]\}, \\ \{0.6, 0.8\}, \{0.1, 0.4\}, \{0.2, 0.3\} \end{array} \right\rangle \end{array} \right)$$

Step 2

using weight vector (0.5,0.5) for decision makers, the aggregated matrix is

$$D = \left(\begin{array}{ccc} \left\langle \left\{ \left[[0.1, 0.387298], [0.2, 0.591608] \right], \right. \right. & \left\langle \left\{ \left[[0.244949, 0.4], [0.2, 0.793725] \right], \right. \right. & \left\langle \left\{ \left[[0.244949, 0.591608], [0.2, 0.52915] \right], \right. \right. \\ \left. \left[[0.2, 0.34641], [0.1, 0.547723] \right], \right. & \left. \left[[0.141421, 0.424264], [0, 0.34641] \right], \right. & \left. \left[[0, 0.447214], [0.1, 0.424264] \right], \right. \\ \left. \left[[0.1, 0.452277], [0.105573, 0.3] \right], \right. & \left. \left[[0.3, 0.452277], [0, 0.206275] \right], \right. & \left. \left[[0.251669, 0.452277], [0, 0.265153] \right], \right. \\ \left. \left\{ 0.4, 0.69282 \right\}, \right. & \left. \left\{ 0.591608, 0.748331 \right\}, \right. & \left. \left\{ 0.547723, 0.648074 \right\}, \right. \\ \left. \left\{ 0.3, 0.447214 \right\}, \right. & \left. \left\{ 0.3, 0.489898 \right\}, \right. & \left. \left\{ 0.2, 0.4 \right\}, \right. \\ \left. \left\{ 0.3, 0.4 \right\} \right. & \left. \left\{ 0.30718, 0.408392 \right\} \right. & \left. \left\{ 0.30718, 0.408392 \right\} \right. \\ \\ \left\langle \left\{ \left[[0.141421, 0.5], [0.3, 0.7] \right], \right. \right. & \left\langle \left\{ \left[[0.173205, 0.424264], [0.2, 0.591608] \right], \right. \right. & \left\langle \left\{ \left[[0.2, 0.447214], [0.282843, 0.648074] \right], \right. \right. \\ \left. \left[[0.173205, 0.34641], [0.1, 0.316228] \right], \right. & \left. \left[[0.141421, 0.244949], [0.141421, 0.547723] \right], \right. & \left. \left[[0.141421, 0.3], [0, 0.34641] \right], \right. \\ \left. \left[[0.151472, 0.351926], [0, 0.251669] \right], \right. & \left. \left[[0.051317, 0.351926], [0.051317, 0.351926] \right], \right. & \left. \left[[0.251669, 0.4], [0, 0.3] \right], \right. \\ \left. \left\{ 0.648074, 0.793725 \right\}, \right. & \left. \left\{ 0.52915, 0.632456 \right\}, \right. & \left. \left\{ 0.447214, 0.547723 \right\}, \right. \\ \left. \left\{ 0.3, 0.489898 \right\}, \right. & \left. \left\{ 0.3, 0.52915 \right\}, \right. & \left. \left\{ 0.3, 0.547723 \right\}, \right. \\ \left. \left\{ 0.265153, 0.367544 \right\} \right. & \left. \left\{ 0.251669, 0.408392 \right\} \right. & \left. \left\{ 0.206275, 0.351926 \right\} \right. \\ \\ \left\langle \left\{ \left[[0.282843, 0.5], [0.173205, 0.4] \right], \right. \right. & \left\langle \left\{ \left[[0.244949, 0.387298], [0.2, 0.447214] \right], \right. \right. & \left\langle \left\{ \left[[0.316228, 0.591608], [0.173205, 0.4] \right], \right. \right. \\ \left. \left[[0.141421, 0.34641], [0.2, 0.5] \right], \right. & \left. \left[[0.141421, 0.34641], [0.2, 0.547723] \right], \right. & \left. \left[[0.141421, 0.387298], [0.2, 0.547723] \right], \right. \\ \left. \left[[0.1, 0.351926], [0.755051, 0.858579] \right], \right. & \left. \left[[0.480385, 0.683772], [0.4, 0.6] \right], \right. & \left. \left[[0.1, 0.4], [0.434315, 0.65359] \right], \right. \\ \left. \left\{ 0.52915, 0.632456 \right\}, \right. & \left. \left\{ 0.591608, 0.734847 \right\}, \right. & \left. \left\{ 0.489898, 0.69282 \right\}, \right. \\ \left. \left\{ 0.424264, 0.565685 \right\}, \right. & \left. \left\{ 0.264575, 0.565685 \right\}, \right. & \left. \left\{ 0.173205, 0.4 \right\}, \right. \\ \left. \left\{ 0.251669, 0.4 \right\} \right. & \left. \left\{ 0.2, 0.408392 \right\} \right. & \left. \left\{ 0.30718, 0.408392 \right\} \right. \end{array} \right)$$

Step 3

using weight vector (0.3,0.4,0.3) for attributes and NCHWG we have the following decision vector

$$d = \left(\begin{array}{c} \left\langle \left\{ \left[[0.18722, 0.4455], [0.2, 0.643505] \right], \right. \right. \\ \left. \left[[0, 0.405589], [0, 0.422372] \right], \right. \\ \left. \left[[0.229912, 0.452277], [0.032918, 0.25311] \right], \right. \\ \left. \left\{ 0.514043, 0.700344 \right\}, \left\{ 0.26564, 0.463822 \right\}, \left\{ 0.305034, 0.405887 \right\} \right. \\ \\ \left\langle \left\{ \left[[0.170172, 0.452793], [0.250618, 0.639485] \right], \right. \right. \\ \left. \left[[0.15029, 0.288831], [0, 0.404857] \right], \right. \\ \left. \left[[0.14557, 0.366739], [0.020852, 0.30752] \right], \right. \\ \left. \left\{ 0.534655, 0.648457 \right\}, \left\{ 0.3, 0.522434 \right\}, \left\{ 0.242474, 0.379688 \right\} \right. \\ \\ \left\langle \left\{ \left[[0.276117, 0.47481], [0.183463, 0.418256] \right], \right. \right. \\ \left. \left[[0.141421, 0.358201], [0.2, 0.532946] \right], \right. \\ \left. \left[[0.277533, 0.524738], [0.549437, 0.71955] \right], \right. \\ \left. \left\{ 0.540653, 0.690198 \right\}, \left\{ 0.268459, 0.509824 \right\}, \left\{ 0.248982, 0.405887 \right\} \right. \end{array} \right)$$

Step 4

Using Score function, we rank the alternatives as $S(F_1) = 0.491743, S(F_2) = 0.511797, S(F_3) = 0.467604$. Hence most desirable alternative is F_2 .




Figure 2: Ranking based on score function

Concluding Remarks

Decision making is one of the crucial problems in real life. For decision making different tools has been established. Torra's hesitant fuzzy set has been used in many practical problems due to flexibility of choosing membership grades. On the other side Jun's neutrosophic cubic set is capable of dealing truth, falsity and indeterminacy membership grades, but the element of hesitancy is missing in truth and falsity membership grades of neutrosophic cubic set. We have discussed the role of hesitancy in truth and falsity membership grades of neutrosophic cubic set. We define neutrosophic cubic hesitant fuzzy set and some basic operations like addition, multiplication, Einstein addition and multiplication in neutrosophic cubic hesitant fuzzy sets. Then we prove some elegant results. In section 4 geometric aggregation operators are defined. Using these aggregation operators an example is constructed.

Acknowledgement: The Deanship of Scientific Research at Prince Sattam bin Abdul-Aziz University has supported this study and all thanks and appreciation to them by the authors of this study.

References

1. Zadeh. L. A. Fuzzy sets, *Inform. Control*, 1965, **8**, pp. 338-353.
2. L. A. Zadeh, Outlines of new approach to the analysis of complex system and decision processes interval valued fuzzy sets, *IEEE Trans.*, 1968.
3. I. B. Turksen, Interval valued strict preferences with Zadeh triplet, *Fuzzy sets and Systems*, (78),1996, 183-195.
4. S. J. Chen and C.L. Hwang, *Fuzzy Multiple Attribute Decision-Making, Methods and Applications; Lecture Notes in Economics and Mathematical Systems; Springer-Verlag: Berlin/Heidelberg, Germany*, (375), 1992.
5. T.H. Chang and T.C. Wang, Using the fuzzy multi-criteria decision making approach for measuring the possibility of successful knowledge management, *Inf. Sci.*, (179), 2009, 355--370.
6. S. Pramanik and K. Mondal, Weighted fuzzy similarity measure based on tangent function and its application to medical diagnosis, *Int. J. Innovat. Res. Sci. Engg. Tech.*, (4), 2015, 158--164.
7. C.T. Chen, Extension of the TOPSIS for group decision-making under fuzzy environment, *Fuzzy Sets Syst.*, (114),

- 2000, 1--9.
8. G. Zhang and J. Lu, An integrated group decision-making method dealing with fuzzy preferences for alternatives and individual judgments for selection criteria, *Group Decis. Negot.*, (12), 2003, 501--515.
 9. R.A. Krohling and V.C. Campanharo, Fuzzy TOPSIS for group decision making: A case study for accidents with oil spill in the sea., *Expert Syst. Appl.*, (38), 2011, 4190--4197.
 10. M. Xia and Z. Xu, A novel method for fuzzy multi-criteria decision making, *Int. J. Inf. Technol. Decis. Mak.*, (13), 2014, 497--519.
 11. M.K. Mehlawat and P.A. Gupta, A new fuzzy group multi-criteria decision making method with an application to the critical path selection, *Int. J. Adv. Manuf. Technol.*, (83), 2016, 1281--1296.
 12. K. Atanassov, Intuitionistic fuzzy sets, *Fuzzy sets and Systems*, (20), 1986, 87-96.
 13. K. Atanassov and G. Gargov, Interval intuitionistic fuzzy sets, *Fuzzy sets and Systems*, (31), 1989, 343-349.
 14. Z. Xu, Some similarity measures of intuitionistic fuzzy sets and their applications to multiple attribute decision making, *Fuzzy Optim. Decis. Mak.*, (6), 2007, 109--121.
 15. K. Mondal and S. Pramanik, Intuitionistic fuzzy multi criteria group decision making approach to quality-brick selection problem, *J. Appl. Quant. Methods*, (9), 2014, 35--50.
 16. P.P. Dey, S. Pramanik and B.C. Giri, Multi-criteria group decision making in intuitionistic fuzzy environment based on grey relational analysis for weaver selection in Khadi institution, *J. Appl. Quant. Methods*, (10), 2015, 1--14.
 17. Y. B. Jun, C. S. Kim and K. O. Yang, Cubic sets, *Ann. Fuzzy Math. Inform.*, (1), 2012, 83-98.
 18. Y. B. Jun, C. S. Kim and M. S. Kang, Cubic subalgebras and ideals of BCK/BCI-algebras, *Far East. J. Math. Sci. (FJMS)*, (44), 2010, 239-250.
 19. Y. B. Jun, C. S. Kim and J. G. Kang, Cubic q-ideals of BCI-algebras, *Ann. Fuzzy Math. Inform.*, (1), 2011, 25-34.
 20. Y. B. Jun, K. J. Lee and M. S. Kang, Cubic structures applied to ideals of BCI-algebras, *Comput. Math. Appl.*, (62), 2011, 3334-3342.
 21. F. Smarandache, A unifying field in logics neutrosophy: neutrosophic probability, set and logic., Rehoboth: American Research Press, 1999.
 22. H. Wang, F. Smarandache, Y. Q. Zhang and R. Sunderraman, Single valued neutrosophic sets, *Multispace Multistructure*, 2010, 410 -- 413.
 23. H. Wang, F. Smarandache, Y. Q. Zhang and R. Sunderraman, Interval neutrosophic sets and logics, Hexis, Phoenix, USA, 2005.
 24. A. Kharal, A neutrosophic multi-criteria decision making method, *New Math. Nat. Comput.*, (10), 2014, 143--162.
 25. J. Ye, Similarity measures between interval neutrosophic sets and their multi criteria decision-making method, *J. Intell. Fuzzy Syst.*, (26), 2014, 165--172.
 26. J. Ye, Multiple attribute group decision-making method with completely unknown weights based on similarity measures under single valued neutrosophic environment, *J. Intell. Fuzzy Syst.*, (27), 2014, 2927--2935.
 27. K. Mondal and S. Pramanik, Multi-criteria group decision making approach for teacher recruitment in higher education under simplified neutrosophic environment, *Neutrosophic Sets Syst.*, (6), 2014, 28--34.
 28. P. Biswas, S. Pramanik, and B.C. Giri, Entropy based grey relational analysis method for multi-attribute decision making under single valued neutrosophic assessments, *Neutrosophic Sets Syst.*, (2), 2014, 102--110.
 29. P. Biswas, and S. Pramanik, and B.C. Giri, A new methodology for neutrosophic multi-attribute decision-making with unknown weight information, *Neutrosophic Sets Syst.*, (3), 2014, 44--54.
 30. P. Biswas, S. Pramanik and B.C. Giri, Cosine similarity measure based multi-attribute decision-making with trapezoidal fuzzy neutrosophic numbers, *Neutrosophic Sets Syst.*, (8), 2014, 46--56.

31. K. Mondal and S. Pramanik, Neutrosophic decision making model for clay-brick selection in construction field based on grey relational analysis, *Neutrosophic Sets Syst.*, (9), 2015, 64--71.
32. B. Li, J. Wang, L. Yang and X. Li, A Novel Generalized Simplified Neutrosophic Number Einstein Aggregation Operator, *International Journal of Applied mathematics*, (48),2018, 67-72.
33. J. Zhan, M. Khan, M. Gulistan, Applications of neutrosophic cubic sets in multi-criteria decision-making, *Int. J. Uncertain. Quantif.*, (7), 2017, 377--394.
34. V. Torra, Hesitant fuzzy sets, *International Journal of Intelligent Systems*, VOL. 25, 529--539 (2010)
35. Y. B. Jun, F. Smarandache and C. S. Kim, Neutrosophic cubic sets, *New. Math. Nat. Comput.*, 2015.
36. T. Mehmood, F. Mehmood and Q. Khan, Cubic Hesitant Fuzzy Sets and Their Applications to Multi Criteria Decision Making, *International Journal of Algebra and Statistics*, Volume 5:1 (2016), 19—51.
37. Ye, J. Multiple-attribute decision-making method using similarity measures of single-valued neutrosophic hesitant fuzzy sets based on least common multiple cardinality. *J. Intell. Fuzzy Syst.* 2018, 34, 4203–4211.
38. P. Liu and L. Shi, The generalized hybrid weighted average operator based on interval neutrosophic hesitant set and Its application to multiple attribute decision making, *Neural Computing and Applications* 26(2) (2015), 457–471.
39. B. Zhu, Z.S. Xu, J.P. Xu, Deriving a ranking from hesitant fuzzy preference relations under group decision making, *IEEE Trans. Cybern.* 44 (8) (2014).
40. E. Turkarslan, J. Ye, M. Unver and M. Olgun, "Consistency Fuzzy Sets and a Cosine Similarity Measure in Fuzzy Multiset Setting and Application to Medical Diagnosis," *Mathematical Problems in Engineering*, vol. 2021, pp. 9, 2021.
41. R. Chinram, T. Mehmood, U. Rehman, Z. Ali and A. Iampan, "Some Novel Cosine Similarity Measures Based on Complex Hesitant Fuzzy Sets and Their Applications," *J. of Maths.*, vol. 2021, pp. 20, 2021.

Received: Feb 4, 2022. Accepted: Jun 10, 2022