



# SINGLE VALUED $(2N+1)$ SIDED POLYGONAL NEUTROSOPHIC NUMBERS AND SINGLE VALUED $(2N)$ SIDED POLYGONAL NEUTROSOPHIC NUMBERS

Said Broumi<sup>1</sup>, Mullai Murugappan<sup>2</sup>, Mohamed Talea<sup>1</sup>, Assia Bakali<sup>3</sup>, Florentin Smarandache<sup>4</sup>, Prem Kumar Singh<sup>5</sup>, Arindam Dey<sup>6</sup>

<sup>1</sup> Laboratory of Information Processing, Faculty of Science Ben M'Sik, University Hassan II, B.P 7955, Sidi Othman, Casablanca, Morocco. E-mail: broumisaid78@gmail.com; taleamohamed@yahoo.fr

<sup>2</sup> Department of Mathematics, Alagappa University, Tamilnadu, India. mullaialu25@gmail.com

<sup>3</sup> Ecole Royale Navale, Boulevard Sour Jdid, B.P 16303 Casablanca, Morocco. E-mail: assiabakali@yahoo.fr

<sup>4</sup> Department of Mathematics, University of New Mexico, 705 Gurley Avenue, Gallup, NM 87301, USA. E-mail: smarand@unm.edu

<sup>5</sup> Amity Institute of Information Technology, Amity University- Sector 125, Noida-Uttar Pradesh, India. E-mail: premsingh.csjm@gmail.com

<sup>6</sup> Department of Computer Science and Engineering, Saroj Mohan Institute of Technology, Hooghly 712512, West Bengal, India. arindam84nit@gmail.com

**Abstract.** This paper introduces a single valued  $(2n)$  as well as  $(2n+1)$  sided polygonal neutrosophic numbers in continuation with other defined single valued neutrosophic numbers. The paper provides basic algebra like addition, subtraction and multiplication of a single valued  $(2n)$  as well as  $(2n+1)$  sided polygonal neutrosophic numbers with examples. In addition, the paper introduces matrix for single valued  $(2n)$  as well as  $(2n+1)$  sided polygonal neutrosophic matrix and its properties.

**Keywords:** Fuzzy numbers, Intuitionistic fuzzy numbers, Single valued trapezoidal neutrosophic numbers, Single valued triangular neutrosophic numbers, Neutrosophic matrix.

## 1 Introduction

In the real world problems, uncertainty occurs in many situations which cannot be handled precisely via crisp set theory. To approximate those uncertainties exists in the given linguistics words the fuzzy set theory is introduced by Zadeh [10]. After that, Dubois and Prade [2] defined the fuzzy number as a generalization of real number. In continuation, many authors [5-8, 11-23] introduced various types of fuzzy numbers such as triangular, trapezoidal, pentagonal, hexagonal fuzzy numbers etc. with their membership functions. Atanassov [1] introduced the concept of intuitionistic fuzzy sets that provides precise solutions to the problems in uncertain situations than fuzzy sets with membership and non-membership functions. After developing intuitionistic fuzzy sets, authors in [4, 6, 10, 19] defined various types of intuitionistic fuzzy numbers and different types of operations on intuitionistic fuzzy sets are also established by suitable examples. Smarandache [9] introduced the generalization of both fuzzy and intuitionistic fuzzy sets and named it as neutrosophic set. The Single valued neutrosophic number and its applications are described in [3]. The results of the problems using neutrosophic sets are more accurate than the results given by fuzzy and intuitionistic fuzzy sets [11-20]. Due to which it is applied in various fields for multi-decision tasks [20-32]. The applications of  $n$ -valued neutrosophic set [24-26] in data analytics research fields given a thrust to study the neutrosophic numbers. This paper focuses on introducing mathematical operation of  $2n$  and  $2n+1$  sided polygonal neutrosophic numbers and its matrices with examples.

The rest of the paper is organized as follows: The section 2 contains preliminaries. Section 3 explains single valued  $2n+1$  polygonal neutrosophic numbers whereas the Section 4 demonstrates Single valued  $2n$  sided polygonal neutrosophic numbers. Section 5 provides conclusions followed by acknowledgements and references.

## 2. Preliminaries

**Definition 1 (Fuzzy Number)[4]:** A fuzzy number is nothing but an extension of a regular number in the sense that it does not refer to one single value but rather to a connected set of possible values, where each of the possible value has its own weight between 0 and 1. This weight is called the membership function. The complex fuzzy set for a given fuzzy number  $\tilde{A}$  can be defined as  $\mu_{\tilde{A}}(x)$  is non-decreasing for  $x \leq x_0$  and non-increasing for  $\geq x_0$ . Similarly other properties can be defined.

**Definition 2** (Triangular fuzzy number [4]): A fuzzy number  $\tilde{A} = \{a, b, c\}$  is said to be a triangular fuzzy number if its membership function is given by, where  $a \leq b \leq c$

$$\mu_{\tilde{A}}(x) = \begin{cases} \frac{(x-a)}{(b-a)} & \text{for } a \leq x \leq b \\ \frac{(c-x)}{(c-b)} & \text{for } b \leq x \leq c \\ 0 & \text{otherwise} \end{cases}$$

**Definition 3** (Trapezoidal fuzzy number [4])

A Trapezoidal fuzzy number (TrFN) denoted by  $\tilde{A}_p$  is defined as  $(a, b, c, d)$ , where the membership function

$$\mu_{\tilde{A}_p}(x) = \begin{cases} 0 & \text{for } x \leq a \\ \frac{(x-a)}{(b-a)} & \text{for } a \leq x \leq b \\ 1 & \text{for } b \leq x \leq c \\ \frac{(d-x)}{(d-c)} & \text{for } c \leq x \leq d \\ 0 & \text{for } x \geq d \end{cases}$$

$$\text{Or, } \mu_{\tilde{A}_p}(x) = \max \left( \min \left( \frac{(x-a)}{(b-a)}, 1, \frac{(d-x)}{(d-c)} \right), 0 \right)$$

**Definition 4 (Generalized Trapezoidal Fuzzy Number) (GTrFNs)**

A Generalized Fuzzy Number  $(a, b, c, d, w)$ , is called a Generalized Trapezoidal Fuzzy Number “x” if its membership function is given by

$$\mu(x) = \begin{cases} 0 & \text{for } x \leq a \\ \frac{(x-a)}{(b-a)} w & \text{for } a \leq x \leq b \\ w & \text{for } b \leq x \leq c \\ \frac{(d-x)}{(d-c)} w & \text{for } c \leq x \leq d \\ 0 & \text{for } x \geq d \end{cases}$$

$$\text{Or, } \mu_{\tilde{A}_p}(x) = \max \left( \min \left( w \frac{(x-a)}{(b-a)}, w, w \frac{(d-x)}{(d-c)} \right), 0 \right)$$

**Definition 5** (Pentagonal fuzzy number [4])

A pentagonal fuzzy number (PFN) of a fuzzy set  $\tilde{A}_p = \{a, b, c, d, e\}$  and its membership function is given by,

$$\mu_{\tilde{A}_p}(x) = \begin{cases} 0 & \text{for } x < a \\ \frac{(x-a)}{(b-a)} & \text{for } a \leq x \leq b \\ \frac{(x-b)}{(c-b)} & \text{for } b \leq x \leq c \\ 1 & \text{for } x = c \\ \frac{(d-x)}{(d-c)} & \text{for } c \leq x \leq d \\ \frac{(e-x)}{(e-d)} & \text{for } d \leq x \leq e \\ 0 & \text{for } x > d \end{cases}$$

**Definition 6** (Hexagonal fuzzy number [4])

A Hexagonal fuzzy number (HFN) of a fuzzy set  $\tilde{A}_p = \{a, b, c, d, e, f\}$  and its membership function is given by,

$$\mu_{\tilde{A}_p}(x) = \begin{cases} 0 & \text{for } x < a \\ \frac{1}{2} \left( \frac{x-a}{b-a} \right) & \text{for } a \leq x \leq b \\ \frac{1}{2} + \frac{1}{2} \left( \frac{x-b}{c-b} \right) & \text{for } b \leq x \leq c \\ 1 & \text{for } c \leq x \leq d \\ 1 - \frac{1}{2} \left( \frac{x-d}{e-d} \right) & \text{for } d \leq x \leq e \\ \frac{1}{2} \left( \frac{f-x}{f-e} \right) & \text{for } e \leq x \leq f \\ 0 & \text{for } x > f \end{cases}$$

**Definition 7** (Octagonal fuzzy number [4])

A Octagonal fuzzy number (OFN) of a fuzzy set  $\tilde{A}_p = \{a_1, a_2, a_3, a_4, a_5, a_6, a_7, a_8\}$  and its membership function is given by,

$$\mu_{\tilde{A}_p} = \begin{cases} k \frac{x-a_1}{a_2-a_1}, & a_1 \leq x \leq a_2 \\ k, & a_2 \leq x \leq a_3 \\ k + (1-k) \frac{x-a_3}{a_4-a_3}, & a_3 \leq x \leq a_4 \\ 1, & a_4 \leq x \leq a_5 \\ k + (1-k) \frac{a_6-x}{a_6-a_5}, & a_5 \leq x \leq a_6 \\ k, & a_6 \leq x \leq a_7 \\ k \frac{a_8-x}{a_8-a_7}, & a_7 \leq x \leq a_8 \\ 0, & \text{Otherwise} \end{cases}$$

Where  $k = \max\{a_1, a_2, a_3, a_4, a_5, a_6, a_7, a_8\}$

**Definition 8 (A triangular intuitionistic fuzzy number)[4]**

A triangular intuitionistic fuzzy number  $\tilde{a}$  is denoted as  $\tilde{a} = ((a, b, c), (a', b', c'))$ , where  $a' \leq a \leq b \leq b' \leq c \leq c'$

with the following membership function  $\mu_{\tilde{a}}(x)$  and non-membership function  $\nu_{\tilde{a}}(x)$

$$\mu_{\tilde{a}}(x) = \begin{cases} \frac{x-a}{b-a}, & a \leq x < b \\ \frac{c-x}{c-b}, & b \leq x < c \\ 0, & \text{otherwise} \end{cases}$$

$$\nu_{\tilde{a}}(x) = \begin{cases} \frac{b-x}{b-a'}, & a' \leq x < b \\ \frac{x-b}{c'-b}, & b \leq x < c' \\ 1, & \text{otherwise} \end{cases}$$

**Definition 9 (Trapezoidal Intuitionistic fuzzy number)**

$$\mu_{\tilde{a}}(x) = \begin{cases} 0 & x \leq a \\ \frac{(x-a)}{(b-a)} & \text{for } a < x < b \\ w & \text{for } b \leq x \leq c \\ \frac{(d-x)}{(d-c)} & \text{for } c < x < d \\ 0 & \text{otherwise} \end{cases}, \nu_{\tilde{a}}(x) = \begin{cases} 1 & x \leq a \\ \frac{(b-x+u_{\tilde{a}}(x-a))}{(b-a)} & \text{for } a < x < b \\ u_{\tilde{a}} & \text{for } b \leq x \leq c \\ \frac{(x-c+u_{\tilde{a}}(d-x))}{(d-c)} & \text{for } c < x < d \\ 1 & \text{otherwise} \end{cases}$$

**Definition 10 (Single valued triangular neutrosophic number [3]):**

A triangular neutrosophic number  $\tilde{a} = \langle (a, b, c); w_{\tilde{a}}, u_{\tilde{a}}, y_{\tilde{a}} \rangle$  is a special neutrosophic set on the real number set  $\mathbb{R}$ , whose truth-membership, indeterminacy-membership and falsity-membership functions are defined as follows:

$$\mu_{\tilde{a}}(x) = \begin{cases} \frac{(x-a)}{(b-a)} w_{\tilde{a}} & \text{for } a \leq x \leq b \\ w_{\tilde{a}} & \text{for } x = b \\ \frac{(c-x)}{(c-b)} w_{\tilde{a}} & \text{for } b \leq x \leq c \\ 0 & \text{otherwise} \end{cases}, \nu_{\tilde{a}}(x) = \begin{cases} \frac{(b-x+u_{\tilde{a}}(x-a))}{(b-a)} & \text{for } a \leq x \leq b \\ u_{\tilde{a}} & \text{for } x = b \\ \frac{(x-b+u_{\tilde{a}}(c-x))}{(c-b)} & \text{for } b \leq x \leq c \\ 1 & \text{otherwise} \end{cases}$$

$$\lambda_{\tilde{a}}(x) = \begin{cases} \frac{(b-x+y_{\tilde{a}}(x-a))}{(b-a)} & \text{for } a \leq x \leq b \\ y_{\tilde{a}} & \text{for } x = b \\ \frac{(x-b+y_{\tilde{a}}(c-x))}{(c-b)} & \text{for } b \leq x \leq c \\ 1 & \text{otherwise} \end{cases}$$

A triangular neutrosophic number  $\tilde{a} = \langle (a, b, c); w_{\tilde{a}}, u_{\tilde{a}}, y_{\tilde{a}} \rangle$  may express an ill-known quantity about  $b$  which is approximately equal to  $b$ .

**Definition 11 (Single valued trapezoidal neutrosophic number [3]):**

A triangular neutrosophic number  $\tilde{a} = \langle (a, b, c, d); w_{\tilde{a}}, u_{\tilde{a}}, y_{\tilde{a}} \rangle$  is a special neutrosophic set on the real number set  $\mathbb{R}$ , whose truth-membership, indeterminacy-membership and falsity-membership function are defined as follows:

$$\mu_{\tilde{a}}(x) = \begin{cases} \frac{(x-a)}{(b-a)} w_{\tilde{a}} & \text{for } a \leq x \leq b \\ w_{\tilde{a}} & \text{for } b \leq x \leq c \\ \frac{(d-x)}{(d-c)} w_{\tilde{a}} & \text{for } c \leq x \leq d \\ 0 & \text{otherwise} \end{cases}, \nu_{\tilde{a}}(x) = \begin{cases} \frac{(b-x+u_{\tilde{a}}(x-a))}{(b-a)} & \text{for } a \leq x \leq b \\ u_{\tilde{a}} & \text{for } b \leq x \leq c \\ \frac{(x-c+u_{\tilde{a}}(d-x))}{(d-c)} & \text{for } c \leq x \leq d \\ 1 & \text{otherwise} \end{cases}$$

$$\lambda_{\tilde{a}}(x) = \begin{cases} \frac{(b-x+y_{\tilde{a}}(x-a))}{(b-a)} & \text{for } a \leq x \leq b \\ y_{\tilde{a}} & \text{for } b \leq x \leq c \\ \frac{(x-c+y_{\tilde{a}}(d-x))}{(d-c)} & \text{for } c \leq x \leq d \\ 1 & \text{otherwise} \end{cases}$$

The single valued trapezoidal neutrosophic numbers are a generalization of the intuitionistic trapezoidal fuzzy numbers, Thus, the neutrosophic number may express more uncertainty than the intuitionistic fuzzy number.

**3. Single valued 2n+1 polygonal neutrosophic numbers**

**Definition 12 (Single valued 2n+1 polygonal neutrosophic number):**

A single valued 2n+1 sided polygonal neutrosophic number  $\tilde{a} = \langle (a_1, a_2, \dots, a_n, \dots, a_{2n}, a_{2n+1}); w_{\tilde{a}}, u_{\tilde{a}}, y_{\tilde{a}} \rangle$  is a special neutrosophic set on the real number set  $\mathbb{R}$ , whose truth-membership, indeterminacy-membership and falsity-membership functions are defined as follows:

$$\begin{aligned}
 T_A(x) = & \left\{ \begin{array}{l} \frac{x - a_1}{a_2 - a_1} w_{\tilde{a}}, \quad a_1 \leq a_2 \\ \frac{x - a_2}{a_3 - a_2} w_{\tilde{a}}, \quad a_2 \leq a_3 \\ \dots \\ \dots \\ \frac{x - a_n}{a_{n+1} - a_n} w_{\tilde{a}}, \quad a_n \leq a_{n+1} \\ w_{\tilde{a}} \quad x = a_{n+1} \\ \frac{a_{n+2} - x}{a_{n+2} - a_{n+1}} w_{\tilde{a}}, \quad a_{n+1} \leq a_{n+2} \\ \frac{a_{n+3} - x}{a_{n+3} - a_{n+2}} w_{\tilde{a}}, \quad a_{n+2} \leq a_{n+3} \\ \dots \\ \dots \\ \frac{a_{2n+1} - x}{a_{2n+1} - a_{2n}} w_{\tilde{a}}, \quad a_{2n} \leq a_{2n+1} \\ 0, \text{ Otherwise} \end{array} \right. & \quad I_A(x) = \left\{ \begin{array}{l} \frac{a_2 - x + u_{\tilde{a}}(x - a_1)}{a_2 - a_1}, \quad a_1 \leq a_2 \\ \frac{a_3 - x + u_{\tilde{a}}(x - a_2)}{a_3 - a_2}, \quad a_2 \leq a_3 \\ \dots \\ \dots \\ \frac{a_{n+1} - x + u_{\tilde{a}}(x - a_n)}{a_{n+1} - a_n}, \quad a_n \leq a_{n+1} \\ u_{\tilde{a}}, \quad x = a_{n+1} \\ \frac{x - a_{n+1} + u_{\tilde{a}}(a_{n+2} - x)}{a_{n+2} - a_{n+1}}, \quad a_{n+1} \leq a_{n+2} \\ \frac{x - a_{n+2} + u_{\tilde{a}}(a_{n+3} - x)}{a_{n+3} - a_{n+2}}, \quad a_{n+2} \leq a_{n+3} \\ \dots \\ \dots \\ \frac{x - a_{2n} + u_{\tilde{a}}(a_{2n+1} - x)}{a_{2n+1} - a_{2n}}, \quad a_{2n} \leq a_{2n+1} \\ 1, \text{ Otherwise} \end{array} \right. \\
 F_A(x) = & \left\{ \begin{array}{l} \frac{a_2 - x + y_{\tilde{a}}(x - a_1)}{a_2 - a_1}, \quad a_1 \leq a_2 \\ \frac{a_3 - x + y_{\tilde{a}}(x - a_2)}{a_3 - a_2}, \quad a_2 \leq a_3 \\ \dots \\ \dots \\ \frac{a_{n+1} - x + y_{\tilde{a}}(x - a_n)}{a_{n+1} - a_n}, \quad a_n \leq a_{n+1} \\ y_{\tilde{a}}, \quad x = a_{n+1} \\ \frac{x - a_{n+1} + y_{\tilde{a}}(a_{n+2} - x)}{a_{n+2} - a_{n+1}}, \quad a_{n+1} \leq a_{n+2} \\ \frac{x - a_{n+2} + y_{\tilde{a}}(a_{n+3} - x)}{a_{n+3} - a_{n+2}}, \quad a_{n+2} \leq a_{n+3} \\ \dots \\ \dots \\ \frac{x - a_{2n} + y_{\tilde{a}}(a_{2n+1} - x)}{a_{2n+1} - a_{2n}}, \quad a_{2n} \leq a_{2n+1} \\ 1, \text{ Otherwise} \end{array} \right.
 \end{aligned}$$

**Example:1** If  $w_{\tilde{a}} = 0.2$  ,  $u_{\tilde{a}} = 0.4$   $y_{\tilde{a}} = 0.3$  and  $n= 4$  , then we have an nanogonal neutrosophic number  $\tilde{a}$  and it is taken as  $\tilde{a} = \langle (3,6,8,10,11,21,43,44,56) \rangle$ . Figure 1 demonstrates the Example 1.




Figure: 1

**Example: 2**

If  $w_{\tilde{a}} = 0.2$  ,  $u_{\tilde{a}} = 0.4$   $y_{\tilde{a}} = 0.3$  and  $n= 4$  , then we have an nanogonal neutrosophic number  $\tilde{a}$  and it is taken as  $\tilde{a} = \langle (3,6,8,10,1,2,4,7,5) \rangle$ . Figure 2 demonstrates the Example 2 and its neutrosophic membership.




Figure: 2

**Note**

The single valued triangular neutrosophic number can be generalized to a single valued  $2n+1$  polygonal neutrosophic number, where  $n=1,2,3,\dots,n$

$\tilde{a} = \langle (a_1, a_2, \dots, a_n, \dots, a_{2n}, a_{2n+1}) ; w_{\tilde{a}}, u_{\tilde{a}}, y_{\tilde{a}} \rangle$ , where  $\tilde{a}$  may express an ill –known quantity about  $a_n$  which is gradually equal to  $a_n$ .

We mean that  $a_2$  approximates  $a_n$ ,  $a_3$  approximates  $a_n$  a littel better than  $a_2, \dots, \dots, \dots, a_{n-1}$  approximates  $a_n$  a litte better than all previous  $a_1, a_2, \dots, a_n$ ,

**Remark**

If  $0 \leq w_{\tilde{a}}, u_{\tilde{a}} \cdot y_{\tilde{a}} \leq 1, 0 \leq w_{\tilde{a}} + u_{\tilde{a}} \cdot y_{\tilde{a}} \leq 1, y_{\tilde{a}} = 0$  and the single valued  $2n+1$  sided polygonal neutrosophic number reduced to the case single valued  $2n+1$  sided polygonal fuzzy number.

**3.1. Operations of single valued  $2n+1$  sided polygonal neutrosophic numbers**

Following are the three operations that can be performed on single valued  $2n+1$  polygonal neutrosophic numbers suppose  $A_{PNN} = \langle (a_1, a_2, \dots, a_n, \dots, a_{2n}, a_{2n+1}); w_{\tilde{a}}, u_{\tilde{a}}, y_{\tilde{a}} \rangle$  and  $B_{PNN} = \langle (b_1, b_2, \dots, b_n, \dots, b_{2n}, b_{2n+1}); w_{\tilde{b}}, u_{\tilde{b}}, y_{\tilde{b}} \rangle$  are two single valued  $2n+1$  polygonal neutrosophic numbers then

(i) **Addition:**

$$A_{PNN} + B_{PNN} = \langle (a_1 + b_1, a_2 + b_2, \dots, a_n + b_n, \dots, a_{2n} + b_{2n}, a_{2n+1} + b_{2n+1}); w_{\tilde{a}} + w_{\tilde{b}} - w_{\tilde{a}} \cdot w_{\tilde{b}}, u_{\tilde{a}} \cdot u_{\tilde{b}}, y_{\tilde{a}} \cdot y_{\tilde{b}} \rangle$$

(ii) **Subtraction:**

$$A_{PNN} - B_{PNN} = \langle (a_1 - b_1, a_2 - b_2, \dots, a_n - b_n, \dots, a_{2n} - b_{2n}, a_{2n+1} - b_{2n+1}); w_{\tilde{a}} + w_{\tilde{b}} - w_{\tilde{a}} \cdot w_{\tilde{b}}, u_{\tilde{a}} \cdot u_{\tilde{b}}, y_{\tilde{a}} \cdot y_{\tilde{b}} \rangle$$

**Multiplication:**

$$A_{PNN} * B_{PNN} = \langle (a_1 \cdot b_1, a_2 \cdot b_2, \dots, a_n \cdot b_n, \dots, a_{2n} \cdot b_{2n}, a_{2n+1} \cdot b_{2n+1}); w_{\tilde{a}} \cdot w_{\tilde{b}}, u_{\tilde{a}} + u_{\tilde{b}} - u_{\tilde{a}} \cdot u_{\tilde{b}}, y_{\tilde{a}} + y_{\tilde{b}} - y_{\tilde{a}} \cdot y_{\tilde{b}} \rangle$$

**Remark**

If  $w_{\tilde{a}} = 1, u_{\tilde{a}} = 0, y_{\tilde{a}} = 0$  then single valued  $2n+1$  sided polygonal neutrosophic number  $A_{PNN} = \langle (a_1, a_2, \dots, a_n, \dots, a_{2n}, a_{2n+1}); w_{\tilde{a}}, u_{\tilde{a}}, y_{\tilde{a}} \rangle$  reduced to the case of single valued  $2n+1$  sided polygonal fuzzy number  $A_{PFN} = \langle (a_1, a_2, \dots, a_n, \dots, a_{2n}, a_{2n+1}) \rangle, n=1,2,3,\dots,n$ .

**Remark**

If  $0 \leq w_{\tilde{a}}, u_{\tilde{a}} \cdot y_{\tilde{a}} \leq 1, 0 \leq w_{\tilde{a}} + u_{\tilde{a}} \cdot y_{\tilde{a}} \leq 3,$  and  $n=1,$  the single valued  $2n+1$  -sided polygonal neutrosophic number reduced to the case of the single valued triangular neutrosophic number  $A_{PNN} = \langle (a_1, a_2, a_3); w_{\tilde{a}}, u_{\tilde{a}}, y_{\tilde{a}} \rangle [3]$ .

**Example 3:** Let  $w_{\tilde{a}} = 1, u_{\tilde{a}} = 0, y_{\tilde{a}} = 0$  and  $n = 1$

If  $w_{\tilde{a}} = 1, u_{\tilde{a}} = 0, y_{\tilde{a}} = 0$  and  $n = 2,$  then we have an Pentagonal fuzzy number [5]:

Let  $A = (1, 2, 3, 4, 5)$  and  $B = (2, 3, 4, 5, 6)$  be two Pentagonal fuzzy numbers, then

- i.  $A + B = (3, 5, 7, 9, 11)$
- ii.  $A - B = (-1, -1, -1, -1, -1)$
- iii.  $2A = (2, 4, 6, 8, 10)$
- iv.  $A \cdot B = (2, 6, 12, 20, 30)$




Figure: 3

Figure 3 demonstrates operation given in Example 3. The single valued  $2n+1$  polygonal neutrosophic number are generalization of the Pentagonal fuzzy number numbers [5], and single valued triangular neutrosophic number [3]

4. Single valued  $2n$ -sided polygonal neutrosophic numbers

**Definition 13:** The single valued trapezoidal neutrosophic number can be extended to a single valued  $2n$  sided polygonal neutrosophic number  $\tilde{a} = \langle (a_1, a_2, \dots, a_n, a_{n+1}, \dots, a_{2n-1}, a_{2n}); w_{\tilde{a}}, u_{\tilde{a}}, y_{\tilde{a}} \rangle$  where  $n=1,2,3,\dots,n$ , whose truth-membership, indeterminacy-membership and falsity-membership functions are defined as follows:

$$T_A(x) = \begin{cases} k \frac{x-a_1}{a_2-a_1} w_{\tilde{a}}, & a_1 \leq x \leq a_2 \\ k + (1-k) \frac{x-a_2}{a_3-a_2} w_{\tilde{a}}, & a_2 \leq x \leq a_3 \\ \dots \\ \dots \\ k + (1-mk) \frac{x-a_{n-1}}{a_n-a_{n-1}} w_{\tilde{a}}, & a_{n-1} \leq x \leq a_n \\ w_{\tilde{a}}, & a_n \leq x \leq a_{n+1} \\ k + (1-mk) \frac{a_{n+2}-x}{a_{n+2}-a_{n+1}} w_{\tilde{a}}, & a_{n+1} \leq x \leq a_{n+2} \\ \dots \\ \dots \\ k + (1-k) \frac{a_{2n-1}-x}{a_{2n-1}-a_{2n-2}} w_{\tilde{a}}, & a_{2n-2} \leq x \leq a_{2n-1} \\ k \frac{a_{2n}-x}{a_{2n}-a_{2n-1}} w_{\tilde{a}}, & a_{2n-1} \leq x \leq a_{2n} \\ 0, & \text{Otherwise} \end{cases}$$



$$I_A(x) = \left\{ \begin{array}{l}
 k + (1 - mk) \frac{a_2 - x}{a_2 - a_1} u_{\tilde{a}}, \quad a_1 \leq x \leq a_2 \\
 k + (1 - (m - 1)k) \frac{a_3 - x}{a_3 - a_2} u_{\tilde{a}}, \quad a_2 \leq x \leq a_3 \\
 \dots \\
 \dots \\
 \dots \\
 k + (1 - k) \frac{a_{n-1} - x}{a_{n-1} - a_{n-2}} u_{\tilde{a}}, \quad a_{n-2} \leq x \leq a_{n-1} \\
 k \frac{a_n - x}{a_n - a_{n-1}} u_{\tilde{a}}, \quad a_{n-1} \leq x \leq a_n \\
 0, \quad a_n \leq x \leq a_{n+1} \\
 k \frac{x - a_{n+1}}{a_{n+2} - a_{n+1}} u_{\tilde{a}}, \quad a_{n+1} \leq x \leq a_{n+2} \\
 k + (1 - k) \frac{x - a_{n+2}}{a_{n+3} - a_{n+2}} u_{\tilde{a}}, \quad a_{n+2} \leq x \leq a_{n+3} \\
 \dots \\
 \dots \\
 \dots \\
 k + (1 - (m - 1)k) \frac{x - a_{2n-2}}{a_{2n-1} - a_{2n-2}} u_{\tilde{a}}, \quad a_{2n-2} \leq x \leq a_{2n-1} \\
 k + (1 - mk) \frac{x - a_{2n-1}}{a_{2n} - a_{2n-1}} u_{\tilde{a}}, \quad a_{2n-1} \leq x \leq a_{2n} \\
 1, \text{ Otherwise}
 \end{array} \right.$$

$$F_A(x) = \left\{ \begin{array}{l} k + (1 - mk) \frac{a_2 - x}{a_2 - a_1} y_{\tilde{a}}, \quad a_1 \leq x \leq a_2 \\ k + (1 - (m - 1)k) \frac{a_3 - x}{a_3 - a_2} y_{\tilde{a}}, \quad a_2 \leq x \leq a_3 \\ \dots \\ \dots \\ \dots \\ k + (1 - k) \frac{a_{n-1} - x}{a_{n-1} - a_{n-2}} y_{\tilde{a}}, \quad a_{n-2} \leq x \leq a_{n-1} \\ k \frac{a_n - x}{a_n - a_{n-1}} y_{\tilde{a}}, \quad a_{n-1} \leq x \leq a_n \\ 0, \quad a_n \leq x \leq a_{n+1} \\ k \frac{x - a_{n+1}}{a_{n+2} - a_{n+1}} y_{\tilde{a}}, \quad a_{n+1} \leq x \leq a_{n+2} \\ k + (1 - k) \frac{x - a_{n+2}}{a_{n+3} - a_{n+2}} y_{\tilde{a}}, \quad a_{n+2} \leq x \leq a_{n+3} \\ \dots \\ \dots \\ \dots \\ k + (1 - (m - 1)k) \frac{x - a_{2n-2}}{a_{2n-1} - a_{2n-2}} y_{\tilde{a}}, \quad a_{2n-2} \leq x \leq a_{2n-1} \\ k + (1 - mk) \frac{x - a_{2n-1}}{a_{2n} - a_{2n-1}} y_{\tilde{a}}, \quad a_{2n-1} \leq x \leq a_{2n} \\ 1, \text{ Otherwise} \end{array} \right.$$

where  $\tilde{a}$  may represent an ill-known quantity of range, which is gradually approximately equal to the interval  $[a_n, a_{n+1}]$ .

We mean that  $(a_2, a_{2n-1})$  approximates  $[a_n, a_{n+1}]$ ,  $(a, a_{2n-2})$  approximates  $[a_n, a_{n+1}]$  a little better than  $(a_2, a_{2n-1})$ , .....  $(a_n, a_{n+1})$  approximates  $[a_n, a_{n+1}]$  a little better than all previous intervals.

**Remark**

If  $0 \leq w_{\tilde{a}}, u_{\tilde{a}}, y_{\tilde{a}} \leq 1, 0 \leq w_{\tilde{a}} + u_{\tilde{a}} + y_{\tilde{a}} \leq 1, y_{\tilde{a}} = 0$  and the single valued  $2n$ -sided polygonal neutrosophic number reduced to the case of single valued  $2n$ -sided polygonal fuzzy number.

**4.1 Single valued 2n-sided polygonal neutrosophic number**

Following are the three operations that can be performed on single valued  $2n$ -sided polygonal neutrosophic numbers suppose  $A_{PNN} = \langle (a_1, a_2, \dots, a_n, a_{n+1}, \dots, a_{2n-1}, a_{2n}); w_{\tilde{a}}, u_{\tilde{a}}, y_{\tilde{a}} \rangle$  and  $B_{PNN} = \langle (b_1, b_2, \dots, b_n, b_{n+1}, \dots, b_{2n-1}, b_{2n}); w_{\tilde{b}}, u_{\tilde{b}}, y_{\tilde{b}} \rangle$  are two  $2n$ -sided polygonal neutrosophic number.

- (i) **Addition:**  $A_{PNN} + B_{PNN} = \langle (a_1 + b_1, a_2 + b_2, \dots, a_n + b_n, a_{n+1} + b_{n+1}, \dots, a_{2n-1} + b_{2n-1}, a_{2n} + b_{2n}); w_{\tilde{a}} + w_{\tilde{b}}, u_{\tilde{a}} \cdot u_{\tilde{b}}, y_{\tilde{a}} \cdot y_{\tilde{b}} \rangle$
- (ii) **Subtraction:**  $A_{PNN} - B_{PNN} = \langle (a_1 - b_1, a_2 - b_2, \dots, a_n - b_n, a_{n+1} - b_{n+1}, \dots, a_{2n-1} - b_{2n-1}, a_{2n} - b_{2n}); w_{\tilde{a}} + w_{\tilde{b}}, u_{\tilde{a}} \cdot u_{\tilde{b}}, y_{\tilde{a}} \cdot y_{\tilde{b}} \rangle$
- (iii) **Multiplication:**  $A_{PNN} * B_{PNN} = \langle (a_1 \cdot b_1, a_2 \cdot b_2, \dots, a_n \cdot b_n, a_{n+1} \cdot b_{n+1}, \dots, a_{2n-1} \cdot b_{2n-1}, a_{2n} \cdot b_{2n}); w_{\tilde{a}} \cdot w_{\tilde{b}}, u_{\tilde{a}} + u_{\tilde{b}}, y_{\tilde{a}} + y_{\tilde{b}} - y_{\tilde{a}} \cdot y_{\tilde{b}} \rangle$

**Remark**

If  $w_{\tilde{a}} = 1, u_{\tilde{a}} = 0, y_{\tilde{a}} = 0$  then single valued  $2n$ -sided polygonal neutrosophic number  $A_{PNN} = \langle (a_1, a_2, \dots, a_n, a_{n+1}, \dots, a_{2n-1}, a_{2n}); w_{\tilde{a}}, u_{\tilde{a}}, y_{\tilde{a}} \rangle$  reduced to the case of single valued  $2n$ -sided polygonal fuzzy number  $A_{PFN} = \langle (a_1, a_2, \dots, a_n, a_{n+1}, \dots, a_{2n-1}, a_{2n}) \rangle$  for  $n = 1, 2, 3, \dots, n$ .


**Remark**

If  $0 \leq w_{\bar{a}}, u_{\bar{a}}, y_{\bar{a}} \leq 1$ ,  $0 \leq w_{\bar{a}} + u_{\bar{a}} + y_{\bar{a}} \leq 3$ , and  $n=2$ , the single valued  $2n$ -sided polygonal neutrosophic number reduced to the case of single valued trapezoidal neutrosophic number  $A_{PNN} = \langle (a_1, a_2, a_3, a_4); w_{\bar{a}}, u_{\bar{a}}, y_{\bar{a}} \rangle [x]$ .

**Example 4:** if  $w_{\bar{a}} = 1$ ,  $u_{\bar{a}} = 0$ ,  $y_{\bar{a}} = 0$  and  $n=3$  then we have an Hexagonal fuzzy number [7-8]:

Let  $A = (1, 2, 3, 5, 6)$  and  $B = (2, 4, 6, 8, 10, 12)$  be two Hexagonal fuzzy numbers then

$A + B = (3, 6, 9, 13, 16, 19)$



**Figure: 4**

Figure 4 demonstrates operation given in Example 4.

The single valued  $2n$ -sided polygonal neutrosophic number are generalization of the hexagonal fuzzy numbers [8], intuitionistic trapezoidal fuzzy numbers [x] and single valued trapezoidal neutrosophic number [3] with its application [12-23] for multi-decision process [24-26].

**5. Conclusion:**

This paper introduces single valued ( $2n$  and  $2n+1$ ) sided polygonal neutrosophic numbers its addition, subtraction, multiplication as well as polygonal neutrosophic matrix with an illustrative example. In near future our focus will be on applications of single-valued  $2n$  sided polygonal neutrosophic numbers and its other mathematical algebra.

**Acknowledgement:**

Authors thank the reviewer for their useful comments and suggestions.

**References**

- [1]. Atanassov K.T.: Intuitionistic Fuzzy Sets, Fuzzy sets and systems, vol.20 (1), 1986, pp: 87-96.
- [2]. Dubois.D and Prade.H: Operations on fuzzy numbers, International Journal of Systems Science, vol.9, no.6., 1978, pp.613-626.
- [3]. Irfan Deli, Yusuf Subas: Single valued neutrosophic numbers and their applications to multicriteria decision making problem.
- [4]. G.Menaka: Ranking of Octagonal Intuitionistic Fuzzy Numbers, IOSR journal of Mathematics, e-ISSN: 2278-5728, P.ISSN:2319-765X. Vol.13.Issue 3 Ver.II(May-June 2017), pp. P.63-71.
- [5]. T. Pathinathan, K. Ponnivalavan: Pentagonal fuzzy number, International Journal of computing Algorithm, Vol 3, 2014, pp. 103-1005.
- [6]. K. Ponnivalavan, T. Pathinathan: Intuitionistic Pentagonal fuzzy number, ARPN Journal of Engineering and Applied Sciences, vol. 10, no 12, 2015, .
- [7]. P. Rajarajeswari, A. SahayaSudha, R. Karhika: A New operation on Hexagonal Fuzzy Number, International Journal of fuzzy Logic Systems(IJFLS), vol 3, no 3, 2013, pp. 15-23.
- [8]. P. Rajarajeswari, A. Sahaya Sudha : Ranking of Hexagonal Fuzzy Numbers for Solving Multi-Objective Fuzzy Linear Programming Problem, International Journal of Computer Applications, vol 84, no 8, 2013, pp. 0975-8887.
- [9]. F. Smarandache: Neutrosophic set, a generalization of the intuitionistic fuzzy sets. Int J Pure Appl Math vol. 24, 2005, pp. 287-297
- [10]. L. A. Zadeh: Fuzzy Sets. Information and Control vol. 8, 1965, pp. 338-353.

- [11]. I. Deli, Y. Şubaş: A ranking method of single valued neutrosophic numbers and its applications to multiattribute decision making problems, *International Journal of Machine Learning and Cybernetics*, vol. 8, no. 2, 2017, pp. 1309–1322. DOI: 10.1007/s13042-016-0505-3.
- [12] I. Deli , Y. Şubaş: Some weighted geometric operators with SVTrN-numbers and their application to multi-criteria decision making problems, *Journal of Intelligent & Fuzzy Systems*, vol. 32, no. 1, 2017, pp. 291-301
- [13] I. Deli: Operators on Single Valued Trapezoidal Neutrosophic Numbers and SVTN-Group Decision Making, *neutrosophic set and system*, vol. 22, 2018, pp. 131-151.
- [14] İ. Deli, S. Broumi: Neutrosophic Soft Matrices and NSM-decision Making, *Journal of Intelligent and Fuzzy Systems*, vol. 28, no. 5, 2015, pp. 2233–2241.
- [15] M. Ali, I. Deli, F. Smarandache: The theory of neutrosophic cubic sets and their applications in pattern recognition, *Journal of Intelligent and Fuzzy Systems*, vol. 30, no. 4, 2016, pp. 1957–1963.
- [16] I. Deli: Interval-valued neutrosophic soft sets and its decision making, *International Journal of Machine Learning and Cybernetics*, vol. 8, no. 2, 2017, pp. 665–676. DOI: 10.1007/s13042-015-0461-3.
- [17]P. Biswas, S. Pramanik, B.C. Giri: Multi-attribute group decision making based on expected value of neutrosophic trapezoidal numbers. In F.Smarandache, & S. Pramanik (Eds., vol.2), *New trends in neutrosophic theory and applications*, 2018, pp. 103-124. Brussels: Pons Editions.
- [18] S. Pramanik, R. Mallick : VIKOR based MAGDM Strategy with Trapezoidal Neutrosophic Numbers, *Neutrosophic Sets and Systems*, vol. 22, 2018, pp. 118-130. DOI: 10.5281/zenodo.2160840
- [19] P. Biswas, S. Pramanik, B. C. Giri: TOPSIS Strategy for Multi-Attribute Decision Making with Trapezoidal Neutrosophic Numbers, *Neutrosophic Sets and Systems*, vol. 19, 2018, pp. 29-39.
- [20] P. Biswas, S. Pramanik, B. C. Giri: Distance Measure Based MADM Strategy with Interval Trapezoidal Neutrosophic Numbers, *Neutrosophic Sets and Systems*, vol. 19, 2018, pp. 40-46. <http://doi.org/10.5281/zenodo.1235165>
- [21] P. Biswas, S. Pramanik, B. C. Giri: Value and ambiguity index based ranking method of single-valued trapezoidal neutrosophic numbers and its application to multi-attribute decision making, *Neutrosophic Sets and Systems*, vol. 12, 2016, pp. 127-138. <doi.org/10.5281/zenodo.571154>
- [22] P. Biswas, S. Pramanik, B. C. Giri, Cosine Similarity Measure Based Multi-Attribute Decision-making with Trapezoidal Fuzzy Neutrosophic Numbers, *Neutrosophic Sets and Systems*, vol. 8, 2015, pp. 46-56. <doi.org/10.5281/zenodo.571274>.
- [23] P. Biswas, S. Pramanik, B. C. Giri, Aggregation of triangular fuzzy neutrosophic set information and its application to multi-attribute decision making, *Neutrosophic Sets and Systems*, Vol. 12, 2016, pp. 20-40. <doi.org/10.5281/zenodo.571125>
- [24] Prem Kumar Singh, Three-way n-valued neutrosophic concept lattice at different granulation, *International Journal of Machine Learning and Cybernetics*, Vol 9, Issue 11, 2018, pp. 1839-1855, doi: 10.1007/s13042-018-0860-3
- [25] Prem Kumar Singh, m-polar fuzzy graph representation of concept lattice, *Engineering Applications of Artificial Intelligence*, vol. 67, 2018, pp. 52-62, <https://doi.org/10.1016/j.engappai.2017.09.011>
- [26] Prem Kumar Singh, Interval-valued neutrosophic graph representation of concept lattice and its  $(\alpha, \beta, \gamma)$ -decomposition, *Arabian Journal for Science and Engineering*, vol. 43, Issue 2, 2018, pp. 723-740, DOI: 10.1007/s13369-017-2718-5
- [27] M.Abdel-Basset, M.Gunasekaran, M.Mohamed & N.Chilamkurti. Three-way decisions based on neutrosophic sets and AHP-QFD framework for supplier selection problem. *Future Generation Computer Systems*, 2018
- [28] V.Chang, , M.Abdel-Basset, & M.Ramachandran. Towards a Reuse Strategic Decision Pattern Framework—from Theories to Practices. *Information Systems Frontiers*, 2018, pp.1-18.
- [29] M.Abdel-Basset, G.Manogaran, A.Gamal & F. Smarandache.A hybrid approach of neutrosophic sets and DEMATEL method for developing supplier selection criteria. *Design Automation for Embedded Systems*, 2018, pp.1-22.
- [30] Nabeeh, N. A., Smarandache, F., Abdel-Basset, M., El-Ghareeb, H. A., & Aboelfetouh, A. (2019). An Integrated Neutrosophic-TOPSIS Approach and Its Application to Personnel Selection: A New Trend in Brain Processing and Analysis. *IEEE Access*, 7, 29734-29744.
- [31] Abdel-Basset, M., Saleh, M., Gamal, A., & Smarandache, F. (2019). An approach of TOPSIS technique for developing supplier selection with group decision making under type-2 neutrosophic number. *Applied Soft Computing*, 77, 438-452.
- [32] Abdel-Baset, M., Chang, V., Gamal, A., & Smarandache, F. (2019). An integrated neutrosophic ANP and VIKOR method for achieving sustainable supplier selection: A case study in importing field. *Computers in Industry*, 106, 94-110.

Received: January 07, 2019, Accepted: March 01, 2019