

## MOMENTS OF THE SMARANDACHE FUNCTION

**Steven R. Finch**  
 MathSoft Inc.  
 101 Main Street  
 Cambridge, MA, USA 02142  
*sfinch@mathsoft.com*

Given a positive integer  $n$ , let  $P(n)$  denote the largest prime factor of  $n$  and  $S(n)$  denote the smallest integer  $m$  such that  $n$  divides  $m!$

This paper extends earlier work [1] on the average value of the Smarandache function  $S(n)$  and is based on a recent asymptotic result [2]:

$$|\{n \leq N : P(n) < S(n)\}| = o\left(\frac{N}{\ln(N)^j}\right) \quad \text{for any positive integer } j$$

due to Ford. We first prove:

**Theorem 1.** 
$$E(S(N)^k) = \frac{1}{N} \cdot \sum_{n=1}^N S(n)^k = \frac{\zeta(k+1)}{k+1} \cdot \frac{N^k}{\ln(k)} + O\left(\frac{N^k}{\ln(N)^2}\right)$$

where  $\zeta(x)$  is the Riemann zeta function. In particular,

$$\lim_{N \rightarrow \infty} \frac{\ln(N)}{N} \cdot E(S(N)) = \frac{\pi^2}{12} = 0.82246703\dots$$

$$\lim_{N \rightarrow \infty} \frac{\ln(N)}{N^2} \cdot \text{Var}(S(N)) = \frac{\zeta(3)}{3} = 0.40068563\dots$$

**Sketch of Proof.** On one hand,

$$L(k) = \lim_{n \rightarrow \infty} \frac{\ln(n)}{n^k} \cdot E(P(n)^k) \leq \lim_{n \rightarrow \infty} \frac{\ln(n)}{n^k} \cdot E(S(n)^k) = \lim_{N \rightarrow \infty} \frac{\ln(N)}{N^{k+1}} \cdot \sum_{n=1}^N S(n)^k$$

The above summation, on the other hand, breaks into two parts:

$$\lim_{N \rightarrow \infty} \frac{\ln(N)}{N^{k+1}} \cdot \left( \sum_{P(n)=S(n)} P(n)^k + \sum_{P(n)<S(n)} S(n)^k \right)$$

The second part vanishes:

$$\lim_{N \rightarrow \infty} \frac{\ln(N)}{N} \cdot \left( \sum_{P(n) < S(n)} \left( \frac{S(n)}{N} \right)^k \right) \leq \lim_{N \rightarrow \infty} \frac{\ln(N)}{N} \cdot \left( \sum_{P(n) < S(n)} 1 \right) = \lim_{N \rightarrow \infty} \frac{\ln(N)}{N} \cdot o\left(\frac{N}{\ln(N)}\right) = 0$$

while the first part is bounded from above:

$$\lim_{N \rightarrow \infty} \frac{\ln(N)}{N^{k+1}} \cdot \left( \sum_{P(n)=S(n)} P(n)^k \right) \leq \lim_{N \rightarrow \infty} \frac{\ln(N)}{N^{k+1}} \cdot \sum_{n=1}^N P(n)^k = \lim_{n \rightarrow \infty} \frac{\ln(n)}{n^k} \cdot E(P(n)^k) = L(k)$$

A formula for  $L(k)$  was found by Knuth and Trabb Pardo [3] and the remaining second-order details follow similarly.

Observe that the ratio  $\sqrt{\text{Var}(S(N))} / E(S(N)) \rightarrow \infty$  as  $N \rightarrow \infty$ , which indicates that the traditional sample moments are unsuitable for estimating the probability distribution of  $S(N)$ . An alternative estimate involves the relative number of digits in the output of  $S$  per digit in the input. A proof of the following is similar to [1]; the integral formulas were discovered by Shepp and Lloyd [4].

**Theorem 2.**

$$\lim_{N \rightarrow \infty} E \left( \left\{ \frac{\ln(S(N))}{\ln(N)} \right\}^k \right) = \int_0^{\infty} \frac{x^{k-1}}{k!} \cdot \exp \left( -x - \int_x^{\infty} \frac{e^{-y}}{y} dy \right) dx = \begin{cases} 0.62432998 & \text{if } k = 1 \\ 0.42669576 & \text{if } k = 2 \\ 0.31363067 & \text{if } k = 3 \\ 0.24387660 & \text{if } k = 4 \\ 0.19792289 & \text{if } k = 5 \end{cases}$$

The mean output of  $S$  hence has asymptotically 62.43% of the number of digits of the input, with a standard deviation of 19.21%. A web-based essay on the Golomb-Dickman constant 0.62432998... appears in [5] and has further extensions and references.

## References

1. S. R. Finch, The average value of the Smarandache function, *Smarandache Notions Journal* 9 (1998) 95-96.
2. K. Ford, The normal behavior of Smarandache function, *Smarandache Notions Journal* 9 (1998) 81-86.
3. D. E. Knuth and L. Trabb Pardo, Analysis of a simple factorization algorithm, *Theoret. Comp. Sci.* 3 (1976) 321-348.
4. L. A. Shepp and S. P. Lloyd, Ordered cycle lengths in a random permutation, *Trans. Amer. Math. Soc.* 121 (1966) 350-557.
5. S. R. Finch, *Favorite Mathematical Constants*, website URL <http://www.mathsoft.com/asolve/constant/constant.html>, MathSoft Inc.