

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/368024084>

# A Study of Derivative and Integration a Neutrosophic Functions

Article · January 2023

DOI: 10.54216/JNFS.050203

---

CITATIONS

0

---

READS

20

3 authors, including:



**A. A. Salama**

Port Said University

285 PUBLICATIONS 4,287 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



New Trends in Neutrosophic Theory and Applications [View project](#)



Call for book-chapters for the collective book "Theory and Applications of NeutroAlgebras as Generalizations of Classical Algebras" [View project](#)



## A Study of Derivative and Integration a Neutrosophic Functions

Ahmed Salamah<sup>1</sup>, Malath F Alaswad<sup>2</sup>, Rasha Dallah<sup>3</sup>

<sup>1</sup> Department of Mathematics, port said, Egypt,

<sup>2,3</sup> Department of Mathematics, Albaath University, Homs, Syria

Emails: [drsalama44@gmail.com](mailto:drsalama44@gmail.com); [Malaz.Aswad@yahoo.com](mailto:Malaz.Aswad@yahoo.com); [rasha.dallah20@gmail.com](mailto:rasha.dallah20@gmail.com)

### Abstract

The objective of this paper is to study and define the neutrosophic real functions with one neutrosophic variable depending on the geometric isometry (AH-Isometry), with a lot of concepts from real analysis including continuity, differentiability, derivativity, integrability. We have presented the formal forms of different popular functions in neutrosophic environment like logarithmic function, exponential function, trigonometric functions. Rising neutrosophic derivative, indefinite integral, and definite integral well defined including rising to neutrosophic functions.

**Keywords:** Neutrosophic real analysis; AH-isometry; Integration; Derivative neutrosophic function; Indefinite integral neutrosophic function; Definite integral neutrosophic function.

### 1.Introduction

Neutrosophy is a new branch of philosophy concerns with the indeterminacy in all areas of life and science. It has become a useful tool in generalizing many classical systems such as equations [1,9], number theory [2,3], topology [4,5], linear spaces [6,10], modules [4,5], and ring of matrices [7,8].

In the literature, we find many studies about neutrosophic calculus, where some definitions and properties were presented about neutrosophic real functions and numbers [10].

The neutrosophic real functions with one variable were defined only in a special case [11], as follows:

$f(x) = g(x) + h(x)I$  where  $I$  takes an interval value defining what is called by neutrosophic thick functions . For example  $f(x) = 2x + 5xI, I \in [0, 0.01]$  is a neutrosophic real thick function.

The problem with this definition, that it does not consider the general case  $f: R(I) \rightarrow R(I); f = f(X)$

And  $X = x + yI \in R(I)$ .

Recently, Abobala et.al, have presented the concept of two-dimensional AH-isometry to study the correspondence between neutrosophic plane  $R(I) \times R(I)$  and the classical module  $R^2 \times R^2$ . Also, the one-dimensional AH-isometry between  $R(I)$  and  $R \times R$ . This isometry was useful in defining inner products and norms [10], ordering [9], and neutrosophic geometrical shapes [10].

In this work, we use the one-dimensional AH-isometry to turn the general case of neutrosophic real functions with one variable into two classical real functions so we will go from  $R(I)$  space into  $R \times R$  space, we study the properties of our functions then we go back to  $R(I)$  space using AH-isometry.

**2 Definitions.**

**Definition 2.1. Neutrosophic Real Number:** [12]

Suppose that  $w$  is a neutrosophic number, then it takes the following standard form:  $w = a + bI$  where  $a, b$  are real coefficients, and  $I$  represents the indeterminacy, where  $0.I = 0$  and  $I^n = I$  for all positive integers  $n$ .

For example:

$$w = 1 + 2I, w = 3 = 3 + 0I.$$

**Definition 2.2. Division of neutrosophic real numbers:** [13]

Suppose that  $w_1, w_2$  are two neutrosophic number, where

$$w_1 = a_1 + b_1I, w_2 = a_2 + b_2I$$

Then:

$$\frac{w_1}{w_2} = \frac{a_1 + b_1I}{a_2 + b_2I} = \frac{a_1}{a_2} + \frac{a_2b_1 - a_1b_2}{a_2(a_2 + b_2)}I$$

**3 Neutrosophic Functions on  $R(I)$ .**

**Definition 3.1** [14]

Let  $R(I) = \{a + bI ; a, b \in R\}$  where  $I^2 = I$  be the neutrosophic field of reals. The one-dimensional isometry (AH-Isometry) is defined as follows: [49]

$$T: R(I) \rightarrow R \times R ; T(a + bI) = (a, a + b)$$

**Remark 3.1.** [14]

$T$  is an algebraic isomorphism between two rings, it has the following properties:

- 1)  $T$  is bijective.
- 2)  $T$  preserves addition and multiplication, i.e.:

$$T[(a + bI) + (c + dI)] = T(a + bI) + T(c + dI)$$

And:

$$T[(a + bI) \cdot (c + dI)] = T(a + bI) \cdot T(c + dI)$$

- 3) Since  $T$  is bijective, then it is invertible by:

$$T^{-1}: R \times R \rightarrow R(I) ; T^{-1}(a, b) = a + (b - a)I$$

- 4)  $T$  preserves distances, i.e.:

The distance on  $R(I)$  can be defined as follows:

$$\text{Let } A = a + bI, B = c + dI \text{ be two neutrosophic real numbers, then } L = \|\overline{AB}\| = d[(a + bI, c + dI)] = |a + bI - (c + dI)| = |(a - c) + I(b - d)| = |a - c| + I|a + b - c - d| = |a - c|.$$

On the other hand, we have:

$$T(\|\overline{AB}\|) = (|a - c|, |(a + b) - (c + d)|) = (d(a, c), d(a + b, c + d)) = d[(a, a + b), (c, c + d)] = d(T(a + bI), T(c + dI))$$

$$= \|\overline{T(AB)}\|.$$

This implies that the distance is preserved up to isometry. i.e.  $\|T(AB)\| = T(\|AB\|)$

**Definition 3.2.** [15]

Let  $f: R(I) \rightarrow R(I)$ ;  $f = f(X)$  and  $X = x + yI \in R(I)$  the  $f$  is called a neutrosophic real function with one neutrosophic variable.

a neutrosophic real function  $f(X)$  written as follows:

$$f(X) = f(x + yI) = f(x) + I[f(x + y) - f(x)]$$

**Theorem 3.1.** any neutrosophic real function into two classical real functions, i.e., to the classical Euclidean plane  $R \times R$ .

**Proof.**

Let  $f(X) = f(x + yI) = f(x) + I[f(x + y) - f(x)]$  a neutrosophic real function.

Now, Using the one-dimensional AH-isometry, we have.

$T(f(X)) = T(f(x) + I[f(x + y) - f(x)])$ , then.

$(f_1, f_2) = (f(x), f(x + y))$ , then, we have.

$$\begin{cases} f_1 = f(x) \\ f_2 = f(x + y) \end{cases}$$

the functions  $f(x), f(x + y)$  are a real functions.

**Definition 3.3. Neutrosophic exponential and logarithmic functions.** [14]

Let  $R(I)$  be the neutrosophic field of reals, we have:

1.  $f(X) = e^X = e^{x+yI} = e^x + I(e^{x+y} - e^x)$
2.  $f(X) = \ln(X) = \ln(x + yI) = \ln x + I(\ln(x + y) - \ln(x))$ , where  $x + yI > 0$ .

**Definition 3.4.** [14]

Let  $R(I)$  be the neutrosophic field of reals, we have:

$$f(X) = \sqrt{x + yI} = \sqrt{x} + I(\sqrt{x + y} - \sqrt{x})$$

**Definition 3.5.**

Let  $R(I)$  be the neutrosophic field of reals, we have:

$$\begin{aligned} f(X) &= A_n X^n + A_{n-1} X^{n-1} + \dots + A_1 X + A_0 \\ &= a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0 \\ &\quad + I[(b_n (x + y)^n + b_{n-1} (x + y)^{n-1} + \dots + b_1 (x + y) + b_0 \\ &\quad - (a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0)] \end{aligned}$$

**Definition 3.6. Neutrosophic Trigonometric Functions:**

Let  $R(I)$  be the neutrosophic field of reals, we have:

1.  $\sin(x + yI) = \sin x + I[\sin(x + y) - \sin x]$
2.  $\cos(x + yI) = \cos x + I[\cos(x + y) - \cos x]$
3.  $\tan(x + yI) = \tan x + I[\tan(x + y) - \tan x]$

**4. derivative a Neutrosophic Functions on  $R(I)$ .**

**Definition 4.1.**

Let  $f(X) = f(x + yI) = f(x) + I[f(x + y) - f(x)]$  a neutrosophic function on  $R(I)$ , the we define a derivative of a neutrosophic function  $f(X)$  as follows:

$$f'(X) = f'(x + yI) = f'(x) + I[f'(x + y) - f'(x)]$$

**Examples 4.1.**

1.  $f(X) = e^X = e^x + I(e^{x+y} - e^x)$ . We have.

$$f'(X) = (e^x)' + I[(e^{x+y})' - (e^x)'] = e^x + I(e^{x+y} - e^x) = e^x.$$

2.  $f(X) = \ln(X) = \ln x + I[\ln(x+y) - \ln(x)]$ . We have.

$$\begin{aligned} f'(X) &= (\ln x)' + I[(\ln(x+y))' - (\ln x)'] = \frac{1}{x} + I\left[\frac{1}{(x+y)} - \frac{1}{x}\right] = \frac{1}{x} + I\left[\frac{x-x-y}{x(x+y)}\right] = \frac{1}{x} + I\left[\frac{-y}{x(x+y)}\right] \\ &= \frac{1+0I}{x+yI} = \frac{1}{X} \end{aligned}$$

3.  $f(X) = Xe^X = xe^x + I((x+y)e^{x+y} - xe^x)$ . We have.

$$f'(X) = (xe^x)' + I[((x+y)e^{x+y})' - (xe^x)'] = e^x + xe^x + I(e^{x+y} + (x+y)e^{x+y} - (e^x + xe^x)) = e^x + I(e^{x+y} - e^x) + [xe^x + I((x+y)e^{x+y} - xe^x)] = e^x + Xe^X.$$

4.  $f(X) = \sin X = \sin(x+yI) = \sin x + I[\sin(x+y) - \sin x]$ .

We have.

$$f'(X) = (\sin x)' + I[(\sin(x+y))' - (\sin x)'] = \cos x + I[\cos(x+y) - \cos x] = \cos X.$$

5.  $f(X) = \cos X = \cos(x+yI) = \cos x + I[\cos(x+y) - \cos x]$ . We have.

$$f'(X) = (\cos x)' + I[(\cos(x+y))' - (\cos x)'] = -\sin x + I[-\sin(x+y) + \sin x] = -(\sin x + I[\sin(x+y) - \sin x]) = -\sin X.$$

6.  $f(X) = e^X \ln(X) + X^2 = e^x \ln(x) + x^2 + I[e^{(x+y)} \ln(x+y) + (x+y)^2 - (e^x \ln(x) + x^2)]$ . We have.

$$f'(X) = (e^x \ln(x) + x^2)' + I[(e^{(x+y)} \ln(x+y) + (x+y)^2)' - (e^x \ln(x) + x^2)'] = \left(e^x \ln(x) + \frac{e^x}{x} + 2x\right)' + I\left[e^{(x+y)} \ln((x+y)) + \frac{e^{(x+y)}}{x+y} + 2(x+y) - \left(e^x \ln(x) + \frac{e^x}{x} + 2x\right)\right] = e^x \ln(X) + \frac{e^X}{X} + 2X.$$

7.  $f(X) = \sin^2 X \cdot \cos X = \sin^2 x \cdot \cos x + I[(\sin^2(x+y) \cdot \cos(x+y) - \sin^2 x \cdot \cos x)]$ . We have.

$$f'(X) = (\sin^2 x \cdot \cos x)' + I[(\sin^2(x+y) \cdot \cos(x+y))' - (\sin^2 x \cdot \cos x)'] = 2\sin x \cdot \cos^2 x - \sin^3 x + I[2\sin(x+y) \cdot \cos^2(x+y) - \sin^3(x+y) - (2\sin x \cdot \cos^2 x - \sin^3 x)] = 2\sin X \cdot \cos^2 X - \sin^3 X.$$

**5. integral a Neutrosophic Functions on  $R(I)$ .****Definition 5.1.**

Let  $f(X) = f(x+yI) = f(x) + I[f(x+y) - f(x)]$  a neutrosophic function on  $R(I)$ , the we define a integration of a neutrosophic function  $f(X)$  as follows:

$$\int f(X) dX = \int f(x) dx + I\left[\int f(x+y) d(x+y) - \int f(x) dx\right] + (a + bI)$$

Where  $(a + bI)$  is a neutrosophic constant number, and  $\int f(X) dX = F(X) = F(x) + I[F(x+y) - F(x)]$ .

**Examples 5.1.**

1.  $f(X) = e^X = e^x + I(e^{x+y} - e^x)$ . We have.

$$\int e^X dX = \int e^x dx + I[\int e^{x+y} d(x+y) - \int e^x dx] = e^x + I(e^{x+y} - e^x) + (a + bI) = e^X + (a + bI).$$

2.  $f(X) = X^2 e^X = x^2 e^x + I((x+y)^2 e^{x+y} - x^2 e^x)$ . We have.

$$\int X^2 e^X dX = \int x^2 e^x dx + I[\int (x+y)^2 e^{x+y} d(x+y) - \int x^2 e^x dx] = (x^2 - 2x + 2)e^x + I[(x+y)^2 - 2(x+y) + 2]e^{x+y} - (x^2 - 2x + 2)e^x + (a + bI) = (X^2 - 2X + 2)e^X + (a + bI).$$

3.  $f(X) = Xe^{X^2} = xe^{x^2} + I((x+y)e^{(x+y)^2} - xe^{x^2})$ . We have.

$$\begin{aligned} \int Xe^{X^2} dX &= \int xe^{x^2} dx + I[\int (x+y)e^{(x+y)^2} d(x+y) - \int xe^{x^2} dx] = \frac{1}{2}e^{x^2} + I\left(\frac{1}{2}e^{(x+y)^2} - \frac{1}{2}e^{x^2}\right) + (a + bI) \\ &= \frac{1}{2}[e^{x^2} + I(e^{(x+y)^2} - e^{x^2})] + (a + bI) = \frac{1}{2}e^{X^2} + (a + bI). \end{aligned}$$

4.  $f(X) = \sin^3 X = \sin^3 x + I[\sin^3(x+y) - \sin^3 x]$  We have.

$$\begin{aligned} \int Xe^{X^2} dX &= \int \sin^3 x dx + I[\int \sin^3(x+y) d(x+y) - \int \sin^3 x dx] = \frac{\cos^3 x}{3} - \cos x + I\left[\frac{\cos^3(x+y)}{3} - \cos(x+y) - \left(\frac{\cos^3 x}{3} - \cos x\right)\right] + (a + bI) \\ &= \frac{\cos^3 X}{3} - \cos X + (a + bI). \end{aligned}$$

5.  $f(X) = X^3 \ln(X) = x^3 \ln x + I[(x+y)^3 \ln(x+y) - x^3 \ln(x)]$ . We have.

$$\int X^3 \ln(X) dX = \int x^3 \ln x dx + I[\int (x + y)^3 \ln(x + y) d(x + y) - \int x^3 \ln x dx] = \frac{x^4}{4} \ln x - \frac{1}{16} x^4 + I\left[\frac{(x+y)^4}{4} \ln(x + y) - \frac{1}{16} (x + y)^4 - \left(\frac{x^4}{4} \ln x - \frac{1}{16} x^4\right)\right] + (a + bI) = \frac{x^4}{4} \ln X - \frac{1}{16} X^4 + (a + bI).$$

6.  $f(X) = \frac{X^2}{1-X^2} = \frac{x^2}{1-x^2} + I\left(\frac{(x+y)^2}{1-(x+y)^2} - \frac{x^2}{1-x^2}\right)$ . We have.

$$\int \frac{x^2}{1-x^2} dX = \int \frac{x^2}{1-x^2} dx + I\left[\int \frac{(x+y)^2}{1-(x+y)^2} d(x + y) - \int \frac{x^2}{1-x^2} dx\right] = -x + \frac{1}{2} \ln \left|\frac{1+x}{1-x}\right| + I\left[-(x + y) + \frac{1}{2} \ln \left|\frac{1+(x+y)}{1-(x+y)}\right| - \left(-x + \frac{1}{2} \ln \left|\frac{1+x}{1-x}\right|\right)\right] + (a + bI) = -X + \frac{1}{2} \ln \left|\frac{1+X}{1-X}\right| + (a + bI).$$

7.  $f(X) = \frac{\sqrt{1+X^2} + \sqrt{1-X^2}}{\sqrt{1-X^4}} = \frac{\sqrt{1+x^2} + \sqrt{1-x^2}}{\sqrt{1-x^4}} + I\left(\frac{\sqrt{1+(x+y)^2} + \sqrt{1-(x+y)^2}}{\sqrt{1-(x+y)^4}} - \frac{\sqrt{1+x^2} + \sqrt{1-x^2}}{\sqrt{1-x^4}}\right)$ . We have.

$$\int \frac{\sqrt{1+X^2} + \sqrt{1-X^2}}{\sqrt{1-X^4}} dX = \int \frac{\sqrt{1+x^2} + \sqrt{1-x^2}}{\sqrt{1-x^4}} dx + I\left[\int \frac{\sqrt{1+(x+y)^2} + \sqrt{1-(x+y)^2}}{\sqrt{1-(x+y)^4}} d(x + y) - \int \frac{\sqrt{1+x^2} + \sqrt{1-x^2}}{\sqrt{1-x^4}} dx\right] = \int \left(\frac{1}{\sqrt{1-x^2}} + \frac{1}{\sqrt{1+x^2}}\right) dx + I\left[\int \left(\frac{1}{\sqrt{1-(x+y)^2}} + \frac{1}{\sqrt{1+(x+y)^2}}\right) d(x + y) - \int \left(\frac{1}{\sqrt{1-x^2}} + \frac{1}{\sqrt{1+x^2}}\right) dx\right] = \text{arc sin} x + \ln|x + \sqrt{1 + x^2}| + I[\text{arc sin}(x + y) + \ln|(x + y) + \sqrt{1 + (x + y)^2}| - (\text{arc sin} x + \ln|x + \sqrt{1 + x^2}|)] + (a + bI) = \text{arc sin} X + \ln|X + \sqrt{1 + X^2}| + (a + bI).$$

**6. The definite integration a Neutrosophic Functions on R(I).**

**Definition 6.1.**

Let  $a + bI, c + dI \in R(I)$ , the we say  $a + bI \leq c + dI$  if  $a \leq c$  and  $a + b \leq c + d$ .

**Definition 6.2.**

Let  $f(X) = f(x + yI) = f(x) + I[f(x + y) - f(x)]$  a neutrosophic function on  $R(I)$ , we define the definite integration of a neutrosophic function  $f(X)$  as follows:

$$\int_{a+bI}^{c+dI} f(X) dX = \int_a^c f(x) dx + I\left[\int_{a+b}^{c+d} f(x + y) d(x + y) - \int_a^c f(x) dx\right]$$

**Examples 5.1.**

1.  $J = \int_{0+0I}^{1+1I} e^X dX = \int_0^1 e^x dx + I\left[\int_0^2 e^{(x+y)} d(x + y) - \int_0^1 e^x dx\right] = [e^x]_0^1 + I\left[[e^{(x+y)}]_0^2 - [e^x]_0^1\right]$   
 $J = (e - e^0) + I[(e^2 - e^0) - (e - e^0)] = (e - 1) + I[(e^2 - 1) - (e - 1)] = e - 1 + I[e^2 - e]$

2.  $J = \int_{\frac{\pi}{6} + \frac{\pi}{6}I}^{\frac{\pi}{2} + \frac{\pi}{2}I} \sin X dX = \int_{\frac{\pi}{6}}^{\frac{\pi}{2}} \sin x dx + I\left[\int_{\frac{\pi}{2}}^{\frac{\pi}{2}} \sin(x + y) d(x + y) - \int_{\frac{\pi}{6}}^{\frac{\pi}{6}} \sin x dx\right] = [-\cos x]_{\frac{\pi}{6}}^{\frac{\pi}{2}} + I\left[-\cos(x + y)]_{\frac{\pi}{2}}^{\frac{\pi}{2}} + [\cos x]_{\frac{\pi}{6}}^{\frac{\pi}{6}}\right]$   
 $J = \left(-0 + \frac{\sqrt{3}}{2}\right) + I\left[(1 - 0) - \left(-0 + \frac{\sqrt{3}}{2}\right)\right] = \frac{\sqrt{3}}{2} + I\left[1 - \frac{\sqrt{3}}{2}\right]$

**3. Conclusion**

In this paper, we have defined the concept of a neutrosophic functions. Also, we have discussed some of their elementary properties such as derivative, indefinite integral, definite integral.

**References**

[1] Abobala, M., "AH-Subspaces in Neutrosophic Vector Spaces", International Journal of Neutrosophic Science, Vol. 6 , pp. 80-86. 2020.

- [2] Abobala, M., "A Study of AH-Substructures in  $n$ -Refined Neutrosophic Vector Spaces", *International Journal of Neutrosophic Science*, Vol. 9, pp.74-85. 2020.
- [3] Sankari, H., and Abobala, M., "Neutrosophic Linear Diophantine Equations With two Variables", *Neutrosophic Sets and Systems*, Vol. 38, pp. 22-30, 2020.
- [4] Sankari, H., and Abobala, M."  $n$ -Refined Neutrosophic Modules", *Neutrosophic Sets and Systems*, Vol. 36, pp. 1-11. 2020.
- [5] Alhamido, R., and Abobala, M., "AH-Substructures in Neutrosophic Modules", *International Journal of Neutrosophic Science*, Vol. 7, pp. 79-86 . 2020.
- [6] Smarandache, F., " A Unifying Field in Logics: Neutrosophic Logic, Neutrosophy, Neutrosophic Set, Neutrosophic Probability", American Research Press. Rehoboth, 2003.
- [7] Suresh, R., and S. Palaniammal,. "Neutrosophic Weakly Generalized open and Closed Sets", *Neutrosophic Sets and Systems*, Vol. 33, pp. 67-77,. 2020.
- [8] Olgun, N., and Hatip, A., "The Effect Of The Neutrosophic Logic On The Decision Making, in Quadruple Neutrosophic Theory And Applications", Belgium, EU, Pons Editions Brussels,pp. 238-253. 2020.
- [9] Hatip, A., Alhamido, R., and Abobala, M., "A Contribution to Neutrosophic Groups", *International Journal of Neutrosophic Science*", Vol. 0, pp. 67-76 . 2019.
- [10] Abobala, M., "  $n$ -Refined Neutrosophic Groups I", *International Journal of Neutrosophic Science*, Vol. 0, pp. 27-34. 2020.
- [11] Abobala, M., "Classical Homomorphisms Between  $n$ -refined Neutrosophic Rings", *International Journal of Neutrosophic Science*", Vol. 7, pp. 74-78. 2020.
- [12] Smarandache, F., and Abobala, M.,  $n$ -Refined neutrosophic Rings, *International Journal of Neutrosophic Science*, Vol. 5 , pp. 83-90, 2020.
- [13] Abobala, M., On Some Special Substructures of Neutrosophic Rings and Their Properties, *International Journal of Neutrosophic Science*", Vol. 4 , pp. 72-81, 2020.
- [14] Abobala, M., "On Some Special Substructures of Refined Neutrosophic Rings", *International Journal of Neutrosophic Science*, Vol. 5, pp. 59-66. 2020.
- [15] Sankari, H., and Abobala, M.," AH-Homomorphisms In neutrosophic Rings and Refined Neutrosophic Rings", *Neutrosophic Sets and Systems*, Vol. 38, pp. 101-112, 2020.