

CONJECTURES ON PARTITIONS OF INTEGERS AS SUMMATIONS OF PRIMES

Florentin Smarandache, Ph D
Associate Professor
Chair of Department of Math & Sciences
University of New Mexico
200 College Road
Gallup, NM 87301, USA
E-mail: smarand@unm.edu

Abstract.

In this short note many conjectures on partitions of integers as summations of prime numbers are presented, which are extension of Goldbach conjecture.

A) Any odd integer n can be expressed as a combination of three primes as follows:

- 1) As a sum of two primes minus another prime: $n = p + q - r$, where p, q, r are all prime numbers.

Do not include the trivial solution: $p = p + q - q$ when p, q are prime.

For example:

$$1 = 3 + 5 - 7 = 5 + 7 - 11 = 7 + 11 - 17 = 11 + 13 - 23 = \dots;$$

$$3 = 5 + 5 - 7 = 7 + 19 - 23 = 17 + 23 - 37 = \dots;$$

$$5 = 3 + 13 - 11 = \dots;$$

$$7 = 11 + 13 - 17 = \dots$$

$$9 = 5 + 7 - 3 = \dots;$$

$$11 = 7 + 17 - 13 = \dots;$$

- a) Is this a conjecture equivalent to Goldbach's Conjecture (any odd integer ≥ 9 is the sum of three primes)?
b) Is the conjecture true when all three prime numbers are different?
c) In how many ways can each odd integer be expressed as above?

- 2) As a prime minus another prime and minus again another prime:
 $n = p - q - r$, where p, q, r are all prime numbers.

For example:

$$1 = 13 - 5 - 7 = 17 - 5 - 11 = 19 - 5 - 13 = \dots;$$

$$3 = 13 - 3 - 7 = 23 - 7 - 13 = \dots;$$

$$5 = 13 - 3 - 5 = \dots;$$

$$7 = 17 - 3 - 7 = \dots;$$

$$9 = 17 - 3 - 5 = \dots;$$

$$11 = 19 - 3 - 5 = \dots$$

- a) In this conjecture equivalent to Goldbach's Conjecture?
- b) Is the conjecture true when all three prime numbers are different?
- c) In how many ways can each odd integer be expressed as above?

B) Any odd integer n can be expressed as a combination of five primes as follows:

3) $n = p + q + r + t - u$, where p, q, r, t, u are all prime numbers, and $t \neq u$.

For example:

$$1 = 3 + 3 + 3 + 5 - 13 = 3 + 5 + 5 + 17 - 29 = \dots;$$

$$3 = 3 + 5 + 11 + 13 - 29 = \dots;$$

$$5 = 3 + 7 + 11 + 13 - 29 = \dots;$$

$$7 = 5 + 7 + 11 + 13 - 29 = \dots;$$

$$9 = 5 + 7 + 11 + 13 - 29 = \dots$$

$$11 = 5 + 7 + 11 + 17 - 29 = \dots$$

- a) Is the conjecture true when all five prime numbers are different?
- b) In how many ways can each odd integer be expressed as above?

4) $n = p + q + r - t - u$, where p, q, r, t, u are all prime numbers, and $t, u \neq p, q, r$.

For example:

$$1 = 3 + 7 + 17 - 13 - 13 = 3 + 7 + 23 - 13 - 19 = \dots;$$

$$3 = 5 + 7 + 17 - 13 - 13 = \dots;$$

$$5 = 7 + 7 + 17 - 13 - 13 = \dots;$$

$$7 = 5 + 11 + 17 - 13 - 13 = \dots;$$

$$9 = 7 + 11 + 17 - 13 - 13 = \dots;$$

$$11 = 7 + 11 + 19 - 13 - 13 = \dots$$

- a) Is the conjecture true when all five prime numbers are different?
- b) In how many ways can each odd integer be expressed as above?

5) $n = p + q - r - t - u$, where p, q, r, t, u are all prime numbers, and $r, t, u \neq p, q$

For example:

$$1 = 11 + 13 - 3 - 3 - 17 = \dots;$$

$$3 = 13 + 13 - 3 - 3 - 17 = \dots;$$

$$5 = 5 + 29 - 5 - 5 - 17 = \dots;$$

$$7 = 3 + 31 - 5 - 5 - 17 = \dots;$$

$$9 = 3 + 37 - 7 - 7 - 17 = \dots;$$

$$11 = 5 + 37 - 7 - 7 - 17 = \dots$$

- a) Is the conjecture true when all five prime numbers are different?
- b) In how many ways can each odd integer be expressed as above?

6) $n = p - q - r - t - u$, where p, q, r, t, u are all prime numbers, and $q, r, t, u \neq p$.

For example:

$$1 = 13 - 3 - 3 - 3 - 3 = \dots;$$

$$3 = 17 - 3 - 3 - 3 - 5 = \dots;$$

$$5 = 19 - 3 - 3 - 3 - 5 = \dots;$$

$$7 = 23 - 3 - 3 - 5 - 5 = \dots;$$

$$9 = 29 - 3 - 5 - 5 - 7 = \dots;$$

$$11 = 31 - 3 - 5 - 5 - 7 = \dots .$$

- a) Is the conjecture true when all five prime numbers are different?
- b) In how many ways can each odd integer be expressed as above?

GENERAL CONJECTURE:

Let $k \geq 3$, and $1 < s < k$ be integers. Then:

i) If k is odd, any odd integer can be expressed as a sum of $k - s$ primes (first set) minus a sum of s primes (second set) [such that the primes of the first set is different from the primes of the second set].

- a) Is the conjecture true when all k prime numbers are different?
- b) In how many ways can each odd integer be expressed as above?

ii) If k is even, any even integer can be expressed as a sum of $k - s$ primes (first set) minus a sum of s primes (second set) [such that the primes of the first set is different from the primes of the second set].

- a) Is the conjecture true when all k prime numbers are different?
- b) In how many ways can each even integer be expressed as above?

REFERENCE

- [1] Smarandache, Florentin, "Collected Papers", Vol. II, Moldova State University Press at Kishinev, article "Prime Conjecture", p. 190, 1997.

[Published in "Math Power", Pima Community College, Tucson, AZ, USA, Vol. 5, No. 9, pp. 2-4, September 1999;
and in "Octagon", Braşov, Romania, Vol. 8, No. 1, pp. 189-191, 2000.]