

TODIM Method for Group Decision Making under Bipolar Neutrosophic Set Environment

Surapati Pramanik^{1*}, Shyamal Dalapati², Shariful Alam³, Tapan Kumar Roy⁴

¹ Department of Mathematics, Nandalal Ghosh B.T. College, Panpur, P.O.-Narayanpur, District –North 24 Parganas, Pin code-743126, West Bengal, India. ^{1*}E-mail: sura_pati@yahoo.co.in

² Department of Mathematics, Indian Institute of Engineering Science and Technology, Shibpur, P.O.-Botanic Garden, Howrah-711103, West Bengal,

India. E-mail: dalapatishyamal30@gmail.com

³ Department of Mathematics, Indian Institute of Engineering Science and Technology, Shibpur, P.O.-Botanic Garden, Howrah-711103, West Bengal,

India. E-mail: salam50in@yahoo.co.in

⁴Department of Mathematics, Indian Institute of Engineering Science and Technology, Shibpur, P.O.-Botanic Garden, Howrah-711103, West Bengal,

India. E-mail: roy_t_k@yahoo.co.in

ABSTRACT

Classical TODIM (an acronym in Portuguese for Interactive Multi criteria Decision Making) method works on crisp numbers to solve multi-attribute group decision making problems. In this paper, we define TODIM method in bipolar neutrosophic set environment to handle multi-attribute group decision making problems, which means we combine the TODIM with bipolar neutrosophic number to deal with multi-attribute group decision making problems. We have proposed a new method for solving multi-attribute group decision making problems. Finally, we solve multi-attribute group decision making problem using our newly proposed TODIM method to show the applicability and effectiveness of the proposed method.

Keywords: Bipolar neutrosophic sets, TODIM method, Multi attribute group decision making.

1. INTRODUCTION

There exist many decision making methods (Triantaphyllou, 2000; Hwang & Yoon, 1981; Shanian & Savadogo, 2009; Chan & Tong, 2007; Rao & Davim, 2008; Gomes & Lima, 1992) in the literature to deal with multi attribute group decision making (MAGDM) problems which are frequently meet in many fields such as politics, economy, military, etc. In classical methods for MAGDM attribute values are assumed as crisp numbers. In realistic decision making problem uncertainty involves due to the complexity of the problem. So crisp numbers are not sufficient to characterize attribute values. To handle this type of difficulties, Zadeh (1965) introduced the concept of fuzzy set by defining membership function. Atanassov (1986) incorporated non-membership function as independent component and defined intuitionistic fuzzy set to deal with uncertainty. Intuitionistic fuzzy set has been rapidly applied to many MADM fields (Gumus et al., 2016; Mondal & Pramanik, 2014c; Mondal & Pramanik, 2015a; Dey et al., 2015; Pramanik & Mukhopadhyaya, 2011; Xu, 2007; Xu & Yager, 2008; Atanassov et al., 2005; Wei, 2010).

Smarandache (1998) introduced the notion of neutrosophic set by incorporating indeterminacy as independent component to intuitionistic fuzzy set. For dealing with the imperfection knowledge received from real world decision making problems, Wang et al. (2010) defined single valued neutrosophic set (SVNS), which is an instance of neutrosophic set.

Neutrosophic sets and SVNSs are essential topics for research in different route of research such as conflict resolution (Pramanik & Roy, 2014), clustering analysis (Ye, 2014a, 2014b), decision making (Biswas et al., 2014a, 2014b, 2015a, 2015b, 2016a, 2016b; Deli & Subas, 2016; Ji, Wang et al., 2016; Kharal, 2014; Pramanik, Banerjee et al., 2016; Pramanik, Dalapati et al., 2016; Ye, 2013a, 2013b, 2014c, 2014d, 2015a, 2015b, 2017), educational problem (Mondal & Pramanik 2014b, 2015b), medical diagnosis (Ye, 2015c), optimization (Pramanik, 2016a, 2016b; Roy & Das, 2015), social problem (Mondal & Pramanik, 2014a; Pramanik & Chakrabarti, 2013), and so on.

Deli et al. (2015) defined bipolar neutrosophic sets and applied it to MCDM problems. Pramanik and Mondal (2016) defined bipolar rough neutrosophic set. Dey et al. (2016) defined TOPSIS for solving MADM problems under bipolar neutrosophic set environment.

Firstly, Gomes and Lima (1992) introduced TODIM method on the basis of the prospect theory (Kahneman & Tversky, 1979).

Krohling & De Souza (2012) developed a generalized version of TODIM called fuzzy TODIM to deal with fuzzy information. Researchers presented fuzzy TODIM methods in varied fuzzy MADM or MAGDM problems (Liu & Teng, 2014; Tosun & Akyu, 2015; Gomes et al., 2013). Fan et al. (2013) extended TODIM method to deal with the hybrid MADM problems where attribute values are crisp numbers, interval numbers and fuzzy information.

Krohling et al. (2013) studied intuitionistic fuzzy TODIM for MCDM problems. Lourenzutti & Krohling (2013) proposed intuitionistic fuzzy random TODIM method which deals intuitionistic fuzzy information and an underlying random vector that affects the performance of the alternatives. Krohling, R. A., & Pacheco proposed interval-valued intuitionistic fuzzy TODIM to tackle MCDM problems involving uncertainty characterized by interval-valued intuitionistic fuzzy numbers.

Wang (2015) extended TODIM method for MCDM in multi-valued neutrosophic set environment. Ji, Zhang et al. (2016) define projection based TODIM method under multi-valued neutrosophic environment and applied it to personal selection. Zhang et al. (2016) proposed TODIM method for group decision making in neutrosophic environment using neutrosophic numbers (Smarandache, 1998) in the form $a + bI$, where 'a' denotes real part and 'bI' denotes indeterminate part. Bipolar neutrosophic numbers are more suitable to deal with the uncertain information and the TODIM is a good decision making method based on prospect theory. Our objective is to propose an extended TODIM method to deal with multi-criteria group decision making problems in which the evaluation information is expressed by bipolar neutrosophic numbers.

Literature review suggests that TODIM method in bipolar neutrosophic set is yet to appear. To fill the gap, we develop a novel TODIM method for MAGDM in bipolar neutrosophic environment. A numerical example of MAGDM problem in bipolar neutrosophic set environment is solved to show the effectiveness of the proposed method.

Rest of the paper is presented as follows: Section 2 recalls some basic definitions of neutrosophic sets, single valued neutrosophic sets, bipolar neutrosophic set. Section 3 develops a novel MAGDM method based on TODIM method in bipolar neutrosophic set environment. Section 4 solves an illustrative example of MAGDM based on proposed TODIM method in bipolar neutrosophic environment. Finally, section 5 presents concluding remarks and future scope of research.

2. PRELIMINARIES

In this section we recall some basic definitions related to neutrosophic sets, bipolar neutrosophic sets and TODIM method.

Definition 2.1: Neutrosophic Set (Smarandache, 1998)

Let U be a space of points (objects), with a generic element in U denoted by u . A neutrosophic sets A in U is characterized by a truth-membership function $\mu_A(u)$, an indeterminacy-membership function $\nu_A(u)$ and a falsity-membership function $\delta_A(u)$,

where, $\mu_A(u), \nu_A(u), \delta_A(u) : U \rightarrow]^{-}0, 1^{+}[$.

Neutrosophic set A can be written as:

$A = \{ \langle u, (\mu_A(u), \nu_A(u), \delta_A(u)) \rangle : u \in U, \mu_A(u), \nu_A(u), \delta_A(u) \in]^{-}0, 1^{+}[\}$. There is no restriction on the sum of $\mu_A(u), \nu_A(u), \delta_A(u)$ so $^{-}0 \leq \mu_A(u) + \nu_A(u) + \delta_A(u) \leq 3^{+}$.

Definition 2.2: Single Valued Neutrosophic Set (Wang et al., 2010)

Let U be a space of points (objects) with a generic element in U denoted by u . A single valued neutrosophic set H in U is characterized by a truth-membership function $\mu_H(u)$, an indeterminacy-membership function $\nu_H(u)$ and a falsity-membership function $\delta_H(u)$, where, $\mu_H(u), \nu_H(u), \delta_H(u) : U \rightarrow [0, 1]$. A single valued neutrosophic set H can be expressed by $H = \{ \langle u, (\mu_H(u), \nu_H(u), \delta_H(u)) \rangle, u \in U \}$. Therefore for each $u \in U$, $\mu_H(u), \nu_H(u), \delta_H(u) \in [0, 1]$ the sum of three functions lies between 0 and 1, i.e. $0 \leq \mu_H(u) + \nu_H(u) + \delta_H(u) \leq 3$.

Definition 2.3: Bipolar Neutrosophic Set (Deli et al., 2015)

Let U be a space of points (objects) with a generic element in U denoted by u . A bipolar neutrosophic set B in U is defined as an object of the form

$B = \{ \langle u, \mu^+(u), \nu^+(u), \delta^+(u), \mu^-(u), \nu^-(u), \delta^-(u) \rangle : u \in U \}$, where, $\mu^+(u), \nu^+(u), \delta^+(u) : U \rightarrow [0, 1]$ and $\mu^-(u), \nu^-(u), \delta^-(u) : U \rightarrow [-1, 0]$. We denote $B = \{ \langle u, \mu^+(u), \nu^+(u), \delta^+(u), \mu^-(u), \nu^-(u), \delta^-(u) \rangle : u \in U \}$ simply $b = \langle \mu^+, \nu^+, \delta^+, \mu^-, \nu^-, \delta^- \rangle$ as a bipolar neutrosophic number (BNN).

Definition 2.4: Containment of Two Bipolar Neutrosophic Sets (Deli et al., 2015)

Let $B_1 = \{ \langle u, \mu_1^+(u), \nu_1^+(u), \delta_1^+(u), \mu_1^-(u), \nu_1^-(u), \delta_1^-(u) \rangle : u \in U \}$ and

$B_2 = \{ \langle u, \mu_2^+(u), \nu_2^+(u), \delta_2^+(u), \mu_2^-(u), \nu_2^-(u), \delta_2^-(u) \rangle : u \in U \}$ be any two bipolar neutrosophic sets in U . Then $B_1 \subseteq B_2$ iff $\mu_1^+(u) \leq \mu_2^+(u)$, $\nu_1^+(u) \geq \nu_2^+(u)$, $\delta_1^+(u) \geq \delta_2^+(u)$ and $\mu_1^-(u) \geq \mu_2^-(u)$, $\nu_1^-(u) \leq \nu_2^-(u)$, $\delta_1^-(u) \leq \delta_2^-(u)$ for all $u \in U$.

Definition 2.5: Equality of Two Bipolar Neutrosophic Sets (Deli et al., 2015)

Let $B_1 = \{ \langle u, \mu_1^+(u), \nu_1^+(u), \delta_1^+(u), \mu_1^-(u), \nu_1^-(u), \delta_1^-(u) \rangle : u \in U \}$ and $B_2 = \{ \langle u, \mu_2^+(u), \nu_2^+(u), \delta_2^+(u), \mu_2^-(u), \nu_2^-(u), \delta_2^-(u) \rangle : u \in U \}$ be any two bipolar neutrosophic sets in U . Then, $B_1 = B_2$ iff $\mu_1^+(u) = \mu_2^+(u)$, $\nu_1^+(u) = \nu_2^+(u)$, $\delta_1^+(u) = \delta_2^+(u)$ and $\mu_1^-(u) = \mu_2^-(u)$, $\nu_1^-(u) = \nu_2^-(u)$, $\delta_1^-(u) = \delta_2^-(u)$ for all $u \in U$.

Definition 2.6: Union of Two Bipolar Neutrosophic Sets (Deli et al., 2015)

Let $B_1 = \{ \langle u, \mu_1^+(u), \nu_1^+(u), \delta_1^+(u), \mu_1^-(u), \nu_1^-(u), \delta_1^-(u) \rangle : u \in U \}$ and $B_2 = \{ \langle u, \mu_2^+(u), \nu_2^+(u), \delta_2^+(u), \mu_2^-(u), \nu_2^-(u), \delta_2^-(u) \rangle : u \in U \}$ be any two bipolar neutrosophic sets in U . Then, their union is defined as

$$B_3(u) = B_1(u) \cup B_2(u) = \{ \langle u, \max(\mu_1^+(u), \mu_2^+(u)), \max(\nu_1^+(u), \nu_2^+(u)), \min(\delta_1^+(u), \delta_2^+(u)), \min(\mu_1^-(u), \mu_2^-(u)), \min(\nu_1^-(u), \nu_2^-(u)), \max(\delta_1^-(u), \delta_2^-(u)) \rangle : u \in U \}, \text{ for all } u \in U.$$

Definition 2.7: Intersection of Two Bipolar Neutrosophic Sets (Deli et al., 2015)

Let $B_1 = \{ \langle u, \mu_1^+(u), \nu_1^+(u), \delta_1^+(u), \mu_1^-(u), \nu_1^-(u), \delta_1^-(u) \rangle : u \in U \}$ and $B_2 = \{ \langle u, \mu_2^+(u), \nu_2^+(u), \delta_2^+(u), \mu_2^-(u), \nu_2^-(u), \delta_2^-(u) \rangle : u \in U \}$ be any two bipolar neutrosophic sets in U . Then, their intersection is defined as

$$B_4(u) = B_1(u) \cap B_2(u) = \{ \langle u, \min(\mu_1^+(u), \mu_2^+(u)), \min(\nu_1^+(u), \nu_2^+(u)), \max(\delta_1^+(u), \delta_2^+(u)), \max(\mu_1^-(u), \mu_2^-(u)), \max(\nu_1^-(u), \nu_2^-(u)), \min(\delta_1^-(u), \delta_2^-(u)) \rangle : u \in U \} \text{ for all } u \in U.$$

Definition 2.8: Compliment of a Bipolar Neutrosophic Set (Deli et al., 2015)

Let $B_1 = \{ \langle u, \mu_1^+(u), \nu_1^+(u), \delta_1^+(u), \mu_1^-(u), \nu_1^-(u), \delta_1^-(u) \rangle : u \in U \}$ be a bipolar neutrosophic set in U . Then the compliment of B_1 is denoted by B_1^c and is defined by

$$B_1^c = \{ \langle u, 1 - \mu_1^+(u), 1 - \nu_1^+(u), 1 - \delta_1^+(u), \{-1\} - \mu_1^-(u), \{-1\} - \nu_1^-(u), \{-1\} - \delta_1^-(u) \rangle : u \in U \} \text{ for all } u \in U.$$

Definition 2.9: Score function of a BNN (Deli et al., 2015)

The score function of a bipolar neutrosophic number $b = \langle \mu^+, \nu^+, \delta^+, \mu^-, \nu^-, \delta^- \rangle$ is denoted by $Sc(b)$ and is defined by

$$Sc(b) = \frac{(\mu^+ + 1 - \nu^+ + 1 - \delta^+ + 1 + \mu^- - \nu^- - \delta^-)}{6}. \tag{1}$$

Definition 2.10: Accuracy function of a BNN (Deli et al., 2015)

The accuracy function of a bipolar neutrosophic number $b = \langle \mu^+, \nu^+, \delta^+, \mu^-, \nu^-, \delta^- \rangle$ is denoted by $Ac(b)$ and is defined by

$$Ac(b) = \mu^+ - \delta^+ + \mu^- - \delta^-. \tag{2}$$

Definition 2.11: Certainty function of a BNN (Deli et al., 2015)

The certainty function of a bipolar neutrosophic number $b = \langle \mu^+, \nu^+, \delta^+, \mu^-, \nu^-, \delta^- \rangle$ is denoted by $C(b)$ and is defined by $C(b) = \mu^+ - \delta^-$

$$\tag{3}$$

Definition 2.12: Comparison procedure of two BNNs (Deli et al., 2015)

Let $b_1 = \langle \mu_1^+, \nu_1^+, \delta_1^+, \mu_1^-, \nu_1^-, \delta_1^- \rangle$ and $b_2 = \langle \mu_2^+, \nu_2^+, \delta_2^+, \mu_2^-, \nu_2^-, \delta_2^- \rangle$ be any two bipolar neutrosophic numbers in U. The comparison procedure is stated as follows:

1. If $Sc(b_1) > Sc(b_2)$, then b_1 is greater than b_2 , denoted by $b_1 > b_2$.
2. If $Sc(b_1) = Sc(b_2)$ and $Ac(b_1) > Ac(b_2)$, then b_1 is greater than b_2 , denoted by $b_1 > b_2$.
3. If $Sc(b_1) = Sc(b_2)$, $Ac(b_1) = Ac(b_2)$ and $C(b_1) > C(b_2)$, then b_1 is greater than b_2 , denoted by $b_1 > b_2$.
4. If $Sc(b) = Sc(b)$, $Ac(b) = Ac(b)$ and $C(b_1) = C(b_2)$, then b_1 is equal to b_2 , denoted by $b_1 = b_2$.

Definition 2.13: Distance measure between two BNNs

Let $b_1 = \langle \mu_1^+, \nu_1^+, \delta_1^+, \mu_1^-, \nu_1^-, \delta_1^- \rangle$ and $b_2 = \langle \mu_2^+, \nu_2^+, \delta_2^+, \mu_2^-, \nu_2^-, \delta_2^- \rangle$ be any two bipolar neutrosophic numbers in U. Distance measure between b_1 and b_2 is denoted by $d_H(b_1, b_2)$ and defined as

$$d_H(b_1, b_2) = \frac{1}{6} [|\mu_1^+ - \mu_2^+| + |\nu_1^+ - \nu_2^+| + |\delta_1^+ - \delta_2^+| + |\mu_1^- - \mu_2^-| + |\nu_1^- - \nu_2^-| + |\delta_1^- - \delta_2^-|] \quad (4)$$

Definition 2.14: Procedure of normalization

Assume that b_{ij} be a BNN to assess i-th alternative with regarding to j-th criterion. A criterion may be benefit type or cost type. To normalize the BNN b_{ij} , we use the following formula.

$$b_{ij}^* = \langle \{1\} - \mu_{ij}^+, \{1\} - \nu_{ij}^+, \{1\} - \delta_{ij}^+, \{-1\} - \mu_{ij}^-, \{-1\} - \nu_{ij}^-, \{1\} - \delta_{ij}^- \rangle \quad (5)$$

3. TODIM METHOD FOR SOLVING MAGDM PROBLEM UNDER BIPOLAR NEUTROSOPHIC ENVIRONMENT

In this section, we propose a MAGDM method under bipolar neutrosophic environment. Assume that $P = \{p_1, p_2, p_3, \dots, p_r\}$ be a set of r alternatives and $C = \{c_1, c_2, c_3, \dots, c_s\}$ be a set of s criteria. Assume that $W = \{w_1, w_2, w_3, \dots, w_s\}$ be the weight vector of the criteria, where $w_k > 0$ and $\sum_{k=1}^s w_k = 1$. Let $D = \{D_1, D_2, D_3, \dots, D_t\}$ be the set of t decision makers and $\lambda = \{\lambda_1, \lambda_2, \lambda_3, \dots, \lambda_t\}$ be the set of weight vector of decision makers, where $\lambda_l > 0$ and $\sum_{l=1}^t \lambda_l = 1$.

In the following sub section, we describe the TODIM based MAGDM method under bipolar neutrosophic set environment. The proposed method is described using the following steps:

Step1- Construction of the decision matrix

Assume that $M^L = (b_{ij}^L)_{r \times s}$ ($L = 1, 2, 3, \dots, t$) be the L-th decision matrix, where information about the alternative p_i provided by the decision maker D_L with respect to attribute C_j ($j = 1, 2, 3, \dots, s$). The L-th decision matrix denoted by M^L (see Equation 6) is constructed as follows:

$$M^L = \begin{pmatrix} p_1 & c_1 & c_2 & \dots & c_s \\ b_{11}^L & b_{12}^L & \dots & \dots & b_{1s}^L \\ p_2 & b_{21}^L & b_{22}^L & \dots & b_{2s}^L \\ \vdots & \vdots & \vdots & \dots & \vdots \\ p_r & b_{r1}^L & b_{r2}^L & \dots & b_{rs}^L \end{pmatrix} \quad (6)$$

where $L = 1, 2, 3, \dots, t$; $i = 1, 2, 3, \dots, r$; $j = 1, 2, 3, \dots, s$.

Step 2-Normalization of the decision matrix

In decision making situation cost criteria and benefit criteria play an important role to choose the best alternative. Cost criteria and benefit criteria exist together, so the decision matrix needs to be normalized. We use Equation 5 to normalize the cost criteria. Benefit criteria need not be normalized. Using Equation 5 the normalize decision matrix (see Equation 6) is represented below (see Equation 7).

$$M^L = \begin{pmatrix} p_1 & \tilde{c}_1 & \tilde{c}_2 & \dots & \tilde{c}_s \\ \tilde{b}_{11}^L & \tilde{b}_{12}^L & \dots & \dots & \tilde{b}_{1s}^L \\ p_2 & \tilde{b}_{21}^L & \tilde{b}_{22}^L & \dots & \tilde{b}_{2s}^L \\ \vdots & \vdots & \vdots & \dots & \vdots \\ p_r & \tilde{b}_{r1}^L & \tilde{b}_{r2}^L & \dots & \tilde{b}_{rs}^L \end{pmatrix} \quad (7)$$

Here $L = 1, 2, 3, \dots, t$; $i = 1, 2, 3, \dots, r$; $j = 1, 2, 3, \dots, s$.

Step 3- Determination of the relative weight of each criterion

We find relative weight of each criterion with respect to criterion with maximum weight. Relative weight is presented as:

$$W_{RC_j} = \frac{w_{C_j}}{w_m}, \text{ where } w_m = \max \{w_1, w_2, w_3, \dots, w_s\}. \quad (8)$$

Step 4- Calculation of score values

If the criteria are benefit criteria, then score values of Equation 6 are calculated by Equation 1, otherwise score values of Equation 7 are calculated by Equation 1.

Step 5- Calculation of accuracy values

If the criteria are benefit type, then accuracy values of Equation 6 are calculated by Equation 2, otherwise score values of Equation 7 are calculated by Equation 2.

Step 6- Construction of the dominance matrix remove

We construct the dominance matrix of each alternative p_i with respect to the criterion C_j of the L -th decision maker D_L (see Equation 9).

(For cost criteria)

$$\left. \begin{aligned} \alpha_c^L(p_i, p_j) &= \sqrt{\frac{W_{RC}}{\sum_{C=1}^s W_{RC}}} d_H(\tilde{b}_{ic}^L, \tilde{b}_{jc}^L), \text{ if } \tilde{b}_{ic}^L > \tilde{b}_{jc}^L \\ &= 0, \text{ if } \tilde{b}_{ic}^L = \tilde{b}_{jc}^L \\ &= -\frac{1}{\xi} \sqrt{\frac{W_{RC}}{\sum_{C=1}^s W_{RC}}} d_H(\tilde{b}_{ic}^L, \tilde{b}_{jc}^L), \text{ if } \tilde{b}_{ic}^L < \tilde{b}_{jc}^L \end{aligned} \right\} \quad (9)$$

(For benefit criteria)

$$\left. \begin{aligned} \alpha_c^L(p_i, p_j) &= \sqrt{\frac{W_{RC}}{\sum_{C=1}^s W_{RC}}} d_H(b_{ic}^L, b_{jc}^L), \text{ if } b_{ic}^L > b_{jc}^L \\ &= 0, \text{ if } b_{ic}^L = b_{jc}^L \\ &= -\frac{1}{\xi} \sqrt{\frac{W_{RC}}{\sum_{C=1}^s W_{RC}}} d_H(b_{ic}^L, b_{jc}^L), \text{ if } b_{ic}^L < b_{jc}^L \end{aligned} \right\} \quad (9a)$$

Here, ‘ ξ ’ denotes decay factor of loss and $\xi > 0$.

Step 7- Construction of the individual final dominance matrix

Using the Equation 10, individual final dominance matrix is constructed as follows:

$$\eta_L = \sum_{c=1}^s \alpha_c^L(p_i, p_j) \quad (10)$$

Step 8- Aggregation of all dominance matrix

Using the Equation 11, the aggregated dominance matrix is obtained as:

$$\eta(p_i, p_j) = \sum_{L=1}^l \lambda_L \eta_L(p_i, p_j) \quad (11)$$

Step 9- Calculation of the global values

Using Equation 12, the global value p_i is obtained as:

$$\beta_i = \frac{\sum_{j=1}^s \eta(p_i, p_j) - \min_{l \leq i \leq r} (\sum_{j=1}^s \eta(p_i, p_j))}{\max_{l \leq i \leq r} (\sum_{j=1}^s \eta(p_i, p_j)) - \min_{l \leq i \leq r} (\sum_{j=1}^s \eta(p_i, p_j))} \quad (12)$$

Step 10- Ranking of the alternatives

Ranking of the alternatives is done based on descending order of global values. The highest global value β_i reflects the best alternative p_i .

4. ILLUSTRATIVE EXAMPLE

To demonstrate the applicability and effectiveness of the proposed method, we solve a MAGDM problem adapted from (Ye, 2014d, Zhang et al., 2016). We assume that an investment company wants to invest a sum of money in the best option. The investment company forms a decision making board involving of three members (D_1, D_2, D_3) who evaluate the four alternatives to invest money. The alternatives are:

1. Car company (p_1),
2. Food company (p_2),
3. Company (p_3), and
4. Arms company (p_4).

Decision makers take decision to evaluate alternatives based on the criteria namely, risk factor (c_1), growth factor (c_2), environment impact (c_3). We consider three criteria as benefit type based on Zhang et al. (2016). Assume that the weight vector of attributes is $W=(.37,.33,.3)^T$ and weight vector of decision makers is $\lambda=(.38,.32,.3)^T$. Now, we apply the proposed MAGDM method to solve the problem using the following steps.

Step1- Construction of the decision matrix

We construct the decision matrix based on information provided by the decision makers in terms of BNN with respect to the criteria as follows:

Decision matrix for D_1

$$M^1 = \begin{pmatrix} & \begin{matrix} c_1 & c_2 & c_3 \end{matrix} \\ \begin{matrix} p_1 \\ p_2 \\ p_3 \\ p_4 \end{matrix} & \begin{pmatrix} (.5,.6,.7,-.3,-.6,-.3) & (.8,.5,.6,-.4,-.6,-.3) & (.9,.4,.6,-.1,-.6,-.5) \\ (.6,.2,.2,-.4,-.5,-.3) & (.6,.3,.7,-.4,-.3,-.5) & (.7,.5,.3,-.4,-.3,-.3) \\ (.8,.3,.5,-.6,-.4,-.5) & (.5,.2,.4,-.1,-.5,-.3) & (.4,.2,.8,-.5,-.3,-.2) \\ (.7,.5,.3,-.6,-.3,-.3) & (.8,.7,.2,-.8,-.6,-.1) & (.6,.3,.4,-.3,-.4,-.7) \end{pmatrix} \end{pmatrix}$$

Decision matrix for D₂

$$M^2 = \begin{pmatrix} & c_1 & C_2 & C_3 \\ p_1 & (.6, .3, .4, -.5, -.3, -.7) & (.5, .3, .4, -.3, -.3, -.4) & (.1, .5, .7, -.5, -.2, -.6) \\ p_2 & (.7, .4, .5, -.3, -.2, -.1) & (.8, .4, .5, -.7, -.3, -.2) & (.6, .2, .7, -.5, -.2, -.9) \\ p_3 & (.8, .3, .2, -.5, -.2, -.6) & (.3, .2, .1, -.6, -.3, -.4) & (.7, .5, .4, -.4, -.3, -.2) \\ p_4 & (.3, .5, .2, -.5, -.5, -.2) & (.5, .6, .4, -.3, -.6, -.7) & (.4, .3, .8, -.5, -.6, -.5) \end{pmatrix}$$

Decision matrix for D₃

$$M^3 = \begin{pmatrix} & c_1 & C_2 & C_3 \\ p_1 & (.9, .6, .4, -.7, -.3, -.2) & (.7, .5, .3, -.6, -.2, -.5) & (.4, .2, .3, -.2, -.5, -.7) \\ p_2 & (.5, .3, .2, -.6, -.4, -.1) & (.5, .2, .7, -.3, -.2, -.5) & (.6, .3, .2, -.7, -.6, -.3) \\ p_3 & (.2, .5, .6, -.4, -.5, -.7) & (.3, .2, .7, -.2, -.3, -.5) & (.8, .2, .4, -.2, -.3, -.6) \\ p_4 & (.8, .5, .5, -.4, -.6, -.3) & (.9, .3, .4, -.5, -.6, -.7) & (.7, .4, .3, -.2, -.5, -.7) \end{pmatrix}$$

Step 2-Normalization of the decision matrix

Since all the criteria are considered as benefit type, we do not need to normalize the decision matrix (M¹, M², M³).

Step 3- Determination of the relative weight of each criterion

Using Equation 8, the relative weights of the criteria are obtained as:

$$W_{RC_1} = 1, W_{RC_2} = .89, W_{RC_3} = .81.$$

Step 4- Calculation of score values

Using Equation 1, we calculate the score values of each alternative with respect to each criterion (see Table 1, 2, and 3).

Table 1: Score value for M¹

	C ₁	C ₂	C ₃
p ₁	.47	.53	.70
p ₂	.60	.50	.52
p ₃	.55	.60	.40
p ₄	.48	.50	.58

Table 2: Score value for M²

	C ₁	C ₂	C ₃
p ₁	.60	.53	.37
p ₂	.47	.45	.55
p ₃	.60	.52	.48
p ₄	.46	.58	.48

Table 3: Score value for M³

	C ₁	C ₂	C ₃
p ₁	.45	.50	.55
p ₂	.48	.50	.55
p ₃	.48	.50	.65
p ₄	.55	.67	.67

Step 5-Calculate accuracy values

Using Equation 2, we calculate the accuracy values of each alternative with respect to each criterion (see Table 4, 5, and 6.)

Table 4: Accuracy value for M¹

$$\begin{pmatrix} & C_1 & C_2 & C_3 \\ p_1 & -.2 & .1 & .7 \\ p_2 & .3 & 0 & .3 \\ p_3 & .2 & .3 & -.7 \\ p_4 & .1 & -.1 & .6 \end{pmatrix}$$

Table 5: Accuracy value for M²

$$\begin{pmatrix} & C_1 & C_2 & C_3 \\ p_1 & .4 & .2 & -.5 \\ p_2 & 0 & -.2 & .3 \\ p_3 & .7 & 0 & .1 \\ p_4 & -.2 & .5 & -.4 \end{pmatrix}$$

Table 6: Accuracy value for M³

$$\begin{pmatrix} & C_1 & C_2 & C_3 \\ p_1 & 0 & .3 & .3 \\ p_2 & -.2 & 0 & 0 \\ p_3 & -.1 & -.1 & .8 \\ p_4 & .2 & .7 & .9 \end{pmatrix}$$

Step 6- Construction of the dominance matrix

Here, using Equation 9, we construct dominance matrix (Taking $\xi = 1$). The dominance matrices are represented in Table 7, 8, 9, 10, 11, 12, 13, 14, and 15.

Table 7: Dominance matrix α_1^1

$$\alpha_1^1 = \begin{pmatrix} & p_1 & p_2 & p_3 & p_4 \\ p_1 & 0 & .27 & -.82 & -.77 \\ p_2 & -.73 & 0 & -.70 & -.64 \\ p_3 & .30 & .26 & 0 & .22 \\ p_4 & .28 & .24 & -.59 & 0 \end{pmatrix}$$

Table 8: Dominance matrix α_2^1

$$\alpha_2^1 = \begin{pmatrix} & p_1 & p_2 & p_3 & p_4 \\ p_1 & 0 & .24 & .27 & .26 \\ p_2 & -.72 & 0 & -.78 & 0 \\ p_3 & -.82 & .26 & 0 & .33 \\ p_4 & -.78 & 0 & -1 & 0 \end{pmatrix}$$

Table 9: Dominance matrix α_3^1

$$\alpha_3^1 = \begin{pmatrix} & p_1 & p_2 & p_3 & p_4 \\ p_1 & 0 & -.88 & .31 & -.82 \\ p_2 & .26 & 0 & .26 & -.75 \\ p_3 & -1 & -.86 & 0 & -.91 \\ p_4 & .25 & .23 & .27 & 0 \end{pmatrix}$$

Table 10: Dominance matrix α_4^1

$$\alpha_4^1 = \begin{pmatrix} & p_1 & p_2 & p_3 & p_4 \\ p_1 & 0 & .27 & -.52 & .29 \\ p_2 & -.73 & 0 & -.73 & .29 \\ p_3 & .19 & .27 & 0 & .29 \\ p_4 & -.79 & -.79 & -.79 & 0 \end{pmatrix}$$

Table 11: Dominance matrix α_2^2

$$\alpha_2^2 = \begin{pmatrix} p_1 & p_2 & p_3 & p_4 \\ p_1 & 0 & .24 & .22 & -.67 \\ p_2 & -.74 & 0 & -.84 & -.95 \\ p_3 & -.67 & .28 & 0 & -.95 \\ p_4 & .22 & .31 & .31 & 0 \end{pmatrix}$$

Table 12: Dominance matrix α_3^2

$$\alpha_3^2 = \begin{pmatrix} p_1 & p_2 & p_3 & p_4 \\ p_1 & 0 & -.77 & -.91 & -.77 \\ p_2 & .23 & 0 & .28 & .24 \\ p_3 & .27 & -.95 & 0 & .28 \\ p_4 & .23 & -.82 & -.95 & 0 \end{pmatrix}$$

Table 13: Dominance matrix α_1^3

$$\alpha_1^3 = \begin{pmatrix} p_1 & p_2 & p_3 & p_4 \\ p_1 & 0 & -.73 & -.94 & -.68 \\ p_2 & .27 & 0 & -.90 & -.79 \\ p_3 & .35 & .33 & 0 & -.73 \\ p_4 & .25 & .29 & .27 & 0 \end{pmatrix}$$

Table 14: Dominance matrix α_2^3

$$\alpha_2^3 = \begin{pmatrix} p_1 & p_2 & p_3 & p_4 \\ p_1 & 0 & .26 & .30 & -.45 \\ p_2 & -.78 & 0 & .14 & -.91 \\ p_3 & -.91 & -.43 & 0 & -.95 \\ p_4 & .26 & .29 & .31 & 0 \end{pmatrix}$$

Table 15: Dominance matrix α_3^3

$$\alpha_3^3 = \begin{pmatrix} p_1 & p_2 & p_3 & p_4 \\ p_1 & 0 & .26 & -.91 & -.86 \\ p_2 & -.88 & 0 & -.95 & -.86 \\ p_3 & .27 & .28 & 0 & -.63 \\ p_4 & .26 & .26 & .91 & 0 \end{pmatrix}$$

Step 7- Construction of the individual final dominance matrix

Using Equation 10, the individual final dominance matrices are constructed (see Table 16, 17, and 18).

Table 16: Final dominance matrix η_1

η_2

$$\eta_1 = \begin{pmatrix} p_1 & p_2 & p_3 & p_4 \\ p_1 & 0 & -.37 & -.24 & -1.33 \\ p_2 & -1.19 & 0 & -1.22 & -1.39 \\ p_3 & -1.52 & -.34 & 0 & -.36 \\ p_4 & -.25 & .47 & -1.32 & 0 \end{pmatrix}$$

Table 17: Final dominance matrix

$$\eta_2 = \begin{pmatrix} p_1 & p_2 & p_3 & p_4 \\ p_1 & 0 & -.26 & -1.21 & -1.15 \\ p_2 & -1.24 & 0 & -1.29 & -.42 \\ p_3 & -.21 & -.40 & 0 & -.38 \\ p_4 & -.34 & -1.3 & -1.43 & 0 \end{pmatrix}$$

Table 18: Final dominance matrix η_3

$$\eta_3 = \begin{pmatrix} & p_1 & p_2 & p_3 & p_4 \\ p_1 & 0 & -.21 & -1.5 & -1.9 \\ p_2 & -1.39 & 0 & -1.7 & -2.6 \\ p_3 & -.29 & .18 & 0 & -2.3 \\ p_4 & .77 & .84 & .77 & 0 \end{pmatrix}$$

Step 8- Aggregation of all dominance matrix

Using Equation 11, the aggregated dominance matrix is represented in Table 19.

Table 19: Aggregated dominance matrix η

$$\eta = \begin{pmatrix} & p_1 & p_2 & p_3 & p_4 \\ p_1 & 0 & -.29 & -.94 & -1.47 \\ p_2 & -1.26 & 0 & -1.07 & -1.43 \\ p_3 & -.73 & -.20 & 0 & -.95 \\ p_4 & .03 & .01 & -.73 & 0 \end{pmatrix}$$

Step 9- Calculation of the global values

Using Equation 12, the global values β_i are calculated as:

$$\beta_1 = .34, \beta_2 = 0, \beta_3 = .61, \beta_4 = 1.$$

Step 10- Ranking of the alternatives

Here $\beta_4 > \beta_3 > \beta_1 > \beta_2$.

Thus the Arm company (p_4) is the best option to invest money.

Section 5. CONCLUSION

In real decision making, the evaluation information of alternatives provided by the decision maker is often incomplete, indeterminate and inconsistent. Bipolar neutrosophic set can describe this kind of information. In this paper, we have developed a new group decision making method based on TODIM under bipolar neutrosophic set environment. Finally, a numerical example is shown to demonstrate its practicality and effectiveness. We hope that the proposed method can be extended for solving multi criteria group decision making in other neutrosophic hybrid environment.

ACKNOWLEDGMENT

We thank both the editors for their useful suggestions.

REFERENCES

- Atanassov, K. (1986). Intuitionistic fuzzy sets. *Fuzzy Sets and Systems*, 20, 87-96.
- Atanassov, K. T., Pasi G., & Yager, R. R. (2005). Intuitionistic fuzzy interpretations of multi-criteria multi-person and multi-measurement tool decision making. *International Journal of Systems Science*, 36, 859-868.
- Biswas, P., Pramanik, S., & Giri, B. C. (2014a). Entropy based grey relational analysis method for multi-attribute decision making under single valued neutrosophic assessments. *Neutrosophic Sets and Systems*, 2, 102–110.
- Biswas, P., Pramanik, S., & Giri, B. C. (2014b). A new methodology for neutrosophic multi-attribute decision making with unknown weight information. *Neutrosophic Sets and Systems*, 3, 42–52.
- Biswas, P., Pramanik, S., & Giri, B. C. (2015a). TOPSIS method for multi-attribute group decision-making under single valued neutrosophic environment. *Neural Computing and Applications*, 27(3), 727- 737. doi: 10.1007/s00521-015-1891-2.
- Biswas, P., Pramanik, S., & Giri, B. C. (2015b). Cosine similarity measure based multi-attribute decision-making with trapezoidal fuzzy neutrosophic numbers. *Neutrosophic Sets and Systems*, 8, 47-57.
- Biswas, P., Pramanik, S., & Giri, B. C. (2016a). Aggregation of triangular fuzzy neutrosophic set information and its application to multi-attribute decision making. *Neutrosophic Sets and Systems*, 12, 20-40.
- Biswas, P., Pramanik, S., & Giri, B. C. (2016b). Value and ambiguity index based ranking method of single-valued trapezoidal neutrosophic numbers and its application to multi-attribute decision making. *Neutrosophic Sets and Systems*, 12, 127-138.
- Chan, J. W. K., & Tong, T. K. L. (2007). Multi-criteria material selection and end-of-life product strategy: grey relational analysis approach. *Materials and Design*, 28, 1539-1546.
- Deli, M., & Smarandache, F. (2015). Bipolar neutrosophic sets and their application based on multi-criteria decision making Proceedings of the 2015 International Conference on Advanced Mechatronic Systems, Beijing, China, August, 249-254.
- Deli, I., & Subas, Y. (2016). A ranking method of single valued neutrosophic numbers and its applications to multi-attribute decision making problems. *International Journal of Machine Learning and Cybernetics*. doi :10.1007/s13042016-0505-3.
- Dey, P.P., Pramanik, S. & Giri, B.C. (2015). Multi-criteria group decision making in intuitionistic fuzzy environment based on grey relational analysis for weaver selection in Khadi institution. *Journal of Applied and Quantitative Methods*, 10(4), 1-14.
- Dey, P. P., Pramanik, S., & Giri, B. C. (2016). TOPSIS for solving multi-attribute decision making problems under bi-polar neutrosophic environment. In F. Smarandache, & S. Pramanik (Eds), *New trends in neutrosophic theory and applications* (pp.65-77). Brussels: Pons Editions.
- Fan, Z.P., Zhang, X., Chen F.D., & Liu, Y. (2013). Extended TODIM method for hybrid multiple attribute decision making problems. *Knowledge-Based Systems*, 42, 331, 40–48.
- Gomes, L. F. A. M., & Lima, M. M. P. P. (1992). TODIM: Basics and application to multi criteria ranking of projects with environmental impacts. *Foundations of Computing and Decision Sciences*, 16, 113-127.
- Gomes, L. F. A. M., Machado, M. A. S., Costa F. F., & Rangel, L. A. D. (2013). Criteria interactions in multiple criteria decision aiding: a Choquet formulation for the TODIM method. *Procedia Computer Science* , 17, 324–331.
- Gumus, S., Kucukvar, M., & Tatari, O. (2016). Intuitionistic fuzzy multi-criteria decision making framework based on life cycle environmental, economic and social impacts: The case of U.S. wind energy. *Sustainable Production and Consumption*. doi.org/10.1016/j.spc.2016.06.006.
- Hwang, C. L., & Yoon, K. (1981). Multiple attribute decision making: methods and applications: a state-of-the-art survey. Springer-Verlag, Berlin, Heidelberg.

- Ji, P., Wang, J. Q., & Zhang, H. Y. (2016). Frank prioritized Bonferroni mean operator with single-valued neutrosophic sets and its application in selecting third-party logistics providers. *Neural Computing and Applications*. doi:10.1007/s00521-016-2660-6.
- Ji, P., Zhang, H., & Wang, J. (2016). A projection-based TODIM method under multi-valued neutrosophic environments and its application in personnel selection. *Neural Computing and Applications*, DOI 10.1007/s00521-016-2436-z.
- Kahneman, D., & Tversky, A. (1979). Prospect theory: an analysis of decision under risk. *Economic Journal of Economic Society*, 47(2), 263–292.
- Kharal, A. (2014). A neutrosophic multi-criteria decision making method. *New Mathematics and Natural Computation*, 10, 143–162.
- Krohling, R. A., & Pacheco, A. G. C. (2014). Interval-valued intuitionistic fuzzy TODIM. *Information Technology and Quantitative Management*, 31, 236-244.
- Krohling, R. A., Pacheco A. G. C., & Siviero, A. L. T. (2013). IF-TODIM: an intuitionistic fuzzy TODIM to multi-criteria decision making. *Knowledge- Based Systems*, 53, 142-146.
- Krohling, R. A., & De Souza, T. T. M. (2012). Combining prospect theory and fuzzy numbers to multi-criteria decision making. *Expert Systems with Application*, 39(13), 11487–11493.
- Liu, P., & Teng, F. (2014). An extended TODIM method for multiple attribute group decision-making based on 2-dimension uncertain linguistic variable. *Complexity*, 21(5), 20–30.
- Lourenzutti, R. & Krohling, R. A. (2013). A study of TODIM in a intuitionistic fuzzy and random environment. *Expert Systems with Applications*, 40, 6459–6468.
- Mondal, K., & Pramanik, S. (2014a). A study on problems of Hijras in West Bengal based on neutrosophic cognitive maps. *Neutrosophic Sets and Systems*, 5, 21-26.
- Mondal, K., & Pramanik, S., (2014b). Multi-criteria group decision making approach for teacher recruitment in higher education under simplified neutrosophic environment. *Neutrosophic Sets and Systems*, 6, 28-34.
- Mondal, K., & Pramanik, S., (2014c). Intuitionistic fuzzy multicriteria group decision making approach to quality clay-brick selection problems based on grey relational analysis. *Journal of Applied Quantitative Methods*, 9(2), 35-50.
- Mondal, K., & Pramanik, S. (2015a). Intuitionistic fuzzy similarity measure based on tangent function and its application to multi-attribute decision. *Global journal of advanced research*, 2 (2), 464-471.
- Mondal, K., & Pramanik, S., (2015b). Neutrosophic decision making model of school choice. *Neutrosophic Sets and Systems*, 7, 62-68.
- Pramanik, S. (2016a). Neutrosophic multi-objective linear programming. *Global Journal of Engineering Science and Research Management*, 3(8), 36-46.
- Pramanik, S. (2016b). Neutrosophic linear goal programming. *Global Journal of Engineering Science and Research Management*, 3(7), 01-11.
- Pramanik, S., Banerjee, D., & Giri, B. C. (2016). Multi – criteria group decision making model in neutrosophic refined set and its application. *Global Journal of Engineering Science and Research Management*, 3(6), 12-18.
- Pramanik, S., Biswas, P., & Giri, B. C. (2017). Hybrid vector similarity measures and their applications to multi-attribute decision making under neutrosophic environment. *Neural Computing and Applications*, 28(5), 1163-76.
- Pramanik, S., & Chakrabarti, S. (2013). A study on problems of construction workers in West Bengal based on neutrosophic cognitive maps. *International Journal of Innovative Research in Science, Engineering and Technology*, 2(11), 6387-6394.
- Pramanik, S., Dalapati, S., & Roy, T. K. (2016). Logistics center location selection approach based on neutrosophic multi-criteria decision making. In F. Smarandache, & S. Pramanik (Eds.), *New trends in neutrosophic theory and applications* (pp. 161-174). Brussels: Pons Editions.

- Pramanik, S., & Mondal, K. (2016). Rough bipolar neutrosophic set. *Global Journal of Engineering Science and Research Management*, 3(6), 71-81.
- Pramanik, S. & Mukhopadhyaya, D. (2011). Grey relational analysis based intuitionistic fuzzy multi-criteria group decision-making approach for teacher selection in higher education. *International Journal of Computer Applications*, 34 (10), 21-29.
- Pramanik, S., & Roy, T. K. (2014). Neutrosophic game theoretic approach to Indo-Pak conflict over Jammu-Kashmir. *Neutrosophic Sets and Systems*, 2, 82-101.
- Rao, R. V., & Davim, J. P. A. (2008). Decision-making framework model for material selection using a combined multiple attributes decision-making method. *International Journal of Advanced Manufacturing Technology*, 35, 751-760.
- Roy, R. & Das, P. (2015). A multi-objective production planning problem based on neutrosophic linear programming approach. *International Journal of Fuzzy Mathematical Archive*, 8(2), 81-91.
- Shanian, A., & Savadogo, O. (2009). A methodological concept for material selection of highly sensitive components based on multiple criteria decision analysis. *Expert Systems with Applications*, 36, 1362–1370.
- Smarandache, F. (1998). *A unifying field in logics: neutrosophic logic. Neutrosophy, neutrosophic set, neutrosophic probability, and neutrosophic statistics*. Rehoboth: American Research Press.
- Tosun, O. & Akyu, G. (2015). A fuzzy TODIM approach for the supplier selection problem. *International Journal of Computational Intelligence Systems*, 8(2), 317–329.
- Triantaphyllou, E. (2000). Multi-criteria decision making methods. In *Multi-criteria decision making methods: A comparative study* (pp. 5-21). Springer US.
- Wang, J.Q. (2008). Overview on fuzzy multi-criteria decision making approach. *Control and Decision*, 23, 601–606.
- Wang, J. Q., & Li, X. E. (2015). TODIM method with multi-valued neutrosophic set. *Control and Decision*, 30 (6), 1139-1142.
- Wang, H. Smarandache, F. Zhang, Y. Q., & Sunderraman, R. (2010). Single valued neutrosophic sets. *Multispace and Multistructure*, 4, 410–413.
- Wei, G. (2010). Some induced geometric aggregation operators with intuitionistic fuzzy information and their application to group decision making. *Applied Soft Computing*, 10, 423-431.
- Xu, Z. (2007). Some similarity measures of intuitionistic fuzzy sets and their applications to multiple attribute decision making. *Fuzzy Optimization and Decision Making*, 6, 109-121.
- Xu, Z., & Yager, R. R. (2008). Dynamic intuitionistic fuzzy multi-attribute making. *International Journal of Approximate Reasoning*, 48, 246-262.
- Ye, J. (2013a). Multi criteria decision-making method using the correlation coefficient under single-valued neutrosophic environment. *International Journal of General Systems* 42, 386–394.
- Ye, J. (2013b). Single valued neutrosophic cross-entropy for multi criteria decision making problems. *Applied Mathematical Modelling*, 38 (3), 1170–1175.
- Ye, J. (2014a). Single valued neutrosophic minimum spanning tree and its clustering method. *Journal of Intelligent Systems*, 23, 311–324.
- Ye, J. (2014b). Clustering methods using distance-based similarity measures of single-valued neutrosophic sets. *Journal of Intelligent Systems*, 23, 379–389.
- Ye, J. (2014c). A multi criteria decision-making method using aggregation operators for simplified neutrosophic sets. *Journal of Intelligent and Fuzzy Systems*, 26, 2459–2466.
- Ye, J. (2014d). Similarity measures between interval neutrosophic sets and their multi criteria decision-making method. *Journal of Intelligent and Fuzzy systems*, 26, 165-172.
- Ye, J. (2015a). Trapezoidal neutrosophic set and its application to multiple attribute decision-making. *Neural Computing and Applications*, 26, 1157–1166.
- Ye, J. (2015b). Bidirectional projection method for multiple attribute group decision making with neutrosophic number. *Neural Computing and Applications*. doi: 10.1007/s00521-015-2123-5.

- Ye, J. (2015c). Improved cosine similarity measures of simplified neutrosophic sets for medical diagnoses. *Artificial Intelligence in Medicine*, 63, 171–179.
- Ye, J. (2017). Projection and bidirectional projection measures of single valued neutrosophic sets and their decision – making method for mechanical design scheme. *Journal of Experimental and Theoretical Artificial Intelligence*, 29(4), 731-740.
- Zadeh, L. A. (1965). Fuzzy sets. *Information and Control*, 8(3), 338-353.
- Zhang, M., Liu, P., & Shi, L. (2016). An extended multiple attribute group decision-making TODIM method based on the neutrosophic numbers. *Journal of Intelligent and Fuzzy Systems*, 30, 1773–1781.
- Abdel-Basset, M., Mohamed, M., Smarandache, F., & Chang, V. (2018). Neutrosophic Association Rule Mining Algorithm for Big Data Analysis. *Symmetry*, 10(4), 106.
- Abdel-Basset, M., & Mohamed, M. (2018). The Role of Single Valued Neutrosophic Sets and Rough Sets in Smart City: Imperfect and Incomplete Information Systems. *Measurement*. Volume 124, August 2018, Pages 47-55
- Abdel-Basset, M., Gunasekaran, M., Mohamed, M., & Smarandache, F. A novel method for solving the fully neutrosophic linear programming problems. *Neural Computing and Applications*, 1-11.
- Abdel-Basset, M., Manogaran, G., Gamal, A., & Smarandache, F. (2018). A hybrid approach of neutrosophic sets and DEMATEL method for developing supplier selection criteria. *Design Automation for Embedded Systems*, 1-22.
- Abdel-Basset, M., Mohamed, M., & Chang, V. (2018). NMCDA: A framework for evaluating cloud computing services. *Future Generation Computer Systems*, 86, 12-29.
- Abdel-Basset, M., Mohamed, M., Zhou, Y., & Hezam, I. (2017). Multi-criteria group decision making based on neutrosophic analytic hierarchy process. *Journal of Intelligent & Fuzzy Systems*, 33(6), 4055-4066.
- Abdel-Basset, M.; Mohamed, M.; Smarandache, F. An Extension of Neutrosophic AHP–SWOT Analysis for Strategic Planning and Decision-Making. *Symmetry* 2018, 10, 116.