

Refined neutrosophic quadruple (po-)hypergroups and their fundamental group

M. Al-Tahan¹, B. Davvaz^{2,*}

¹Lebanese International University, Bekaa, 961, Lebanon.

E-mail: madeline.tahan@liu.edu.lb

²Yazd University, Yazd, 98, Iran.

E-mail: davvaz@yazd.ac.ir

*Correspondence: B. Davvaz (davvaz@yazd.ac.ir)

Abstract: After introducing the notion of hyperstructures about 80 years ago by F. Marty, a number of researches on its theory, generalization, and its applications have been done. On the other hand, the theory of Neutrosophy, the study of neutralities, was developed in 1995 by F. Smarandache as an extension of dialectics. This paper aims at finding a connection between refined neutrosophy of sets and hypergroups. In this regard, we define refined neutrosophic quadruple hypergroups, study their properties, and find their fundamental refined neutrosophic quadruple groups. Moreover, some results related to refined neutrosophic quadruple po-hypergroups are obtained.

Keywords: H_v -group, po-hypergroup, refined neutrosophic quadruple number, refined neutrosophic quadruple hypergroup, fundamental group.

1 Introduction

In 1934, Marty [19] introduced the concept of hypergroups by considering the quotient of a group by its subgroup. And this was the birth of an interesting new branch of Mathematics known as “Algebraic hyperstructures” which is considered as a generalization of algebraic structures. In algebraic structure, the composition of two elements is an element whereas in algebraic hyperstructure, the composition of two elements is a non-empty set. Since then, many different kinds of hyperstructures (hyperrings, hypermodules, hypervector spaces, ...) were introduced. And many studies were done on the theory of algebraic hyperstructures as well on their applications to various subjects of Sciences (see [12, 13, 30]). Later, in 1991, Vougioklis [28] generalized hyperstructures by introducing a larger class known as weak hyperstructures or H_v -structures. For more details about H_v -structures, see [28, 29, 30, 31].

In 1965, Zadeh [32] extended the classical notion of sets by introducing the notion of Fuzzy sets whose elements have degrees of membership. The theory of fuzzy sets is mainly concerned with the measurement of the degree of membership and non-membership of a given abstract situation. Despite its wide range of real life applications, fuzzy set theory can not be applied to models or problems that contain indeterminacy. This is the reason that arose the importance of introducing a new logic known as neutrosophic logic that contains the concept of indeterminacy. It was introduced by F. Smarandache in 1995, studied and developed by him

and by other authors. For more details about neutrosophic theory, we refer to [17, 22, 23, 25]. Recently, many authors are working on the applications of this important concept. For example in [2], Abdel-Baset et al. offered a novel approach for estimating the smart medical devices selection process in a group decision making in a vague decision environment and used neutrosophics in their methodology. Moreover, in [7], R. Alhabib et al. worked on some neutrosophic probability distribution. Other interesting applications of it are found in [1, 3, 4, 15, 20].

In 2015, Smarandache [22] introduced the concept of neutrosophic quadruple numbers and presented some basic operations on the set of neutrosophic quadruple numbers such as, addition, subtraction, multiplication, and scalar multiplication. After that, a connection between neutrosophy and algebraic structures was established where Agboola et al. [5] considered the set of neutrosophic quadruple numbers and used the defined operations on it to discuss neutrosophic quadruple algebraic structures. More results about neutrosophic algebraic structures are found in [11, 26]. A generalization of the latter work was done in 2016 where Akinleye et al. [6] considered the set of neutrosophic quadruple numbers and defined some hyperoperations on it and discussed neutrosophic quadruple hyperstructures. More specifically, the latter papers introduced the notions of neutrosophic groups, neutrosophic rings, neutrosophic hypergroups and neutrosophic hyperrings on a set of real numbers and studied their basic properties.

The authors in [9] discussed neutrosophic quadruple H_v -groups and studied their properties. Then in [10], they found the fundamental group of neutrosophic quadruple H_v -groups and proved that it is a neutrosophic quadruple group. This paper is an extension to the above mentioned results. In Section 2, some definitions related to weak hyperstructures have been presented while section 3 involves the refined neutrosophic quadruple hypergroup and the studying of its properties. As for section 4, an order on refined neutrosophic quadruple hypergroups is defined and some examples on refined neutrosophic quadruple po-hypergroups are presented. Finally, in section 5, the fundamental refined neutrosophic quadruple group of refined neutrosophic quadruple hypergroups with some important theorems, corollaries and propositions have been submitted.

2 Preliminaries

In this section, some definitions and theorems related to both: hyperstructure theory and neutrosophic theory are presented. (See [12, 13, 30].)

2.1 Basic notions of hypergroups

Definition 2.1. Let H be a non-empty set. Then, a mapping $\circ : H \times H \rightarrow \mathcal{P}^*(H)$ is called a *binary hyperoperation* on H , where $\mathcal{P}^*(H)$ is the family of all non-empty subsets of H . The couple (H, \circ) is called a *hypergroupoid*.

In this definition, if A and B are two non-empty subsets of H and $x \in H$, then:

$$A \circ B = \bigcup_{\substack{a \in A \\ b \in B}} a \circ b, \quad x \circ A = \{x\} \circ A \quad \text{and} \quad A \circ x = A \circ \{x\}.$$

Definition 2.2. A hypergroupoid (H, \circ) is called a:

1. *semihypergroup* if for every $x, y, z \in H$, we have $x \circ (y \circ z) = (x \circ y) \circ z$;

2. *quasi-hypergroup* if for every $x \in H, x \circ H = H = H \circ x$ (The latter condition is called the reproduction axiom);
3. *hypergroup* if it is a semihypergroup and a quasi-hypergroup.

Definition 2.3. [13] Let (H, \star) and (K, \star') be two hypergroups. Then $f : H \rightarrow K$ is said to be *hypergroup homomorphism* if $f(x \star y) = f(x) \star' f(y)$ for all $x, y \in H$. (H, \star) and (K, \star') are called *isomorphic H_v -groups*, and written as $H \cong K$, if there exists a bijective function $f : R \rightarrow S$ that is also a homomorphism. The set of all isomorphism of (H, \star) is denoted as $Aut(H)$.

T. Vougiouklis, the pioneer of H_v -structures, generalized the concept of algebraic hyperstructures to weak algebraic hyperstructures [28]. The latter concept is known as “weak” since the equality sign in the definitions of H_v -structures is more likely to be replaced by non-empty intersection. The concepts in H_v -structures are mostly used in representation theory [29].

A hypergroupoid (H, \circ) is called an *H_v -semigroup* if $(x \circ (y \circ z)) \cap ((x \circ y) \circ z) \neq \emptyset$ for all $x, y, z \in H$. An element $0 \in H$ is called an *identity* if $x \in (0 \circ x \cap x \circ 0)$ for all $x \in H$ and it is called a *scalar identity* if $x = 0 \circ x = x \circ 0$ for all $x \in H$. If the scalar identity exists then it is unique. A hypergroupoid (H, \circ) is called an *H_v -group* if it is a quasi-hypergroup and an H_v -semigroup. A non empty subset S of an H_v -group (H, \circ) is called *H_v -subgroup of H* if (S, \circ) is an H_v -group.

Definition 2.4. [27] A hypergroup is called *cyclic* if there exist $h \in H$ such that $H = h \cup h^2 \cup \dots \cup h^i \cup \dots$ with $i \in \mathbb{N}$. If there exists $s \in \mathbb{N}$ such that $H = h \cup h^2 \cup \dots \cup h^s$ then H is a cyclic hypergroup with finite period. Otherwise, H is called cyclic hypergroup with infinite period. Here, $h^s = \underbrace{h \star h \star \dots \star h}_{s \text{ times}}$.

Definition 2.5. [27] A hypergroup is called a *single power cyclic hypergroup* if there exist $h \in H$ and $s \in \mathbb{N}$ such that $H = h \cup h^2 \cup \dots \cup h^s \cup \dots$ and $h \cup h^2 \cup \dots \cup h^{m-1} \subset h^m$ for every $m \geq 1$. In this case, h is called a *generator of H* .

2.2 Refined neutrosophic quadruple hypergroups

Let T, I, F , represent the neutrosophic components truth, indeterminacy, and falsehood respectively. Symbolic (or Literal) Neutrosophic theory is referring to the use of these symbols in neutrosophics. In 2013, F. Smarandache [24] introduced the refined neutrosophic components. Where the neutrosophic literal components T, I, F can be split into respectively the following neutrosophic literal subcomponents:

$$T_1, \dots, T_p; I, \dots, I_r; F_1, \dots, F_s,$$

where p, r, s are positive integers with $\max\{p, r, s\} \geq 2$.

Definition 2.6. [25] Let X be a nonempty set and $p, r, s \in \mathbb{N}$ with $(p, r, s) \neq (1, 1, 1)$. A refined neutrosophic quadruple X -number is a number having the following form:

$$a + \sum_{i=1}^p b_i T_i + \sum_{j=1}^r c_j I_j + \sum_{k=1}^s b_k F_k,$$

where $a, b_i, c_j, d_k \in X$ and T, I, F have their usual neutrosophic logic meanings, and T_i, I_j, F_k are refinements of T, I, F respectively.

The set of all refined neutrosophic quadruple X -numbers is denoted by $RNQ(X)$, that is,

$$RNQ(X) = \{a + \sum_{i=1}^p b_i T_i + \sum_{j=1}^r c_j I_j + \sum_{k=1}^s d_k F_k : a, b_i, c_j, d_k \in X\}.$$

For simplicity, we write $a + \sum_{i=1}^p b_i T_i + \sum_{j=1}^r c_j I_j + \sum_{k=1}^s d_k F_k$ as

$$(a, \sum_{i=1}^p b_i T_i, \sum_{j=1}^r c_j I_j, \sum_{k=1}^s d_k F_k).$$

In what follows, T_i, I_j, F_k are refinements of T, I, F respectively with $1 \leq i \leq p, 1 \leq j \leq r$ and $1 \leq k \leq s$. Let $(H, +)$ be a hypergroupoid with identity “0” and $0 + 0 = 0$ and define “ \oplus ” on $RNQ(H)$ as follows:

$$\begin{aligned} & (a, \sum_{i=1}^p b_i T_i, \sum_{j=1}^r c_j I_j, \sum_{k=1}^s d_k F_k) \oplus (a', \sum_{i=1}^p b'_i T_i, \sum_{j=1}^r c'_j I_j, \sum_{k=1}^s d'_k F_k) \\ & = \{(x, \sum_{i=1}^p y_i T_i, \sum_{j=1}^r z_j I_j, \sum_{k=1}^s w_k F_k) : x \in a + a', y_i \in b_i + b'_i, z_j \in c_j + c'_j, w_k \in d_k + d'_k\}. \end{aligned}$$

3 New properties of refined neutrosophic quadruple hypergroups

In this section, refined neutrosophic quadruple hypergroups are defined and their properties are studied.

Proposition 3.1. *Let $(H, +)$ be a hypergroupoid with $0 \in H$ and T_i, I_j, F_k are refinements of T, I, F respectively. Then $(RNQ(H), \oplus)$ is a quasi-hypergroup with identity $\bar{0} = (0, \sum_{i=1}^p 0T_i, \sum_{j=1}^r 0I_j, \sum_{k=1}^s 0F_k)$ if and only if $(H, +)$ is a quasi-hypergroup with identity 0.*

Proof. Let $(H, +)$ be a quasi-hypergroup. We prove now that $(RNQ(H), \oplus)$ satisfies the reproduction axiom. That is, $\bar{x} \oplus RNQ(H) = RNQ(H) \oplus \bar{x} = RNQ(H)$ for all $\bar{x} = (a, \sum_{i=1}^p b_i T_i, \sum_{j=1}^r c_j I_j, \sum_{k=1}^s d_k F_k) \in RNQ(H)$. We prove $\bar{x} \oplus RNQ(H) = RNQ(H)$ and the proof of $RNQ(H) \oplus \bar{x} = RNQ(H)$ is done in a similar manner. Let $\bar{y} = (a', \sum_{i=1}^p b'_i T_i, \sum_{j=1}^r c'_j I_j, \sum_{k=1}^s d'_k F_k) \in RNQ(H)$, we have $\bar{x} \oplus \bar{y} = (a + a', \sum_{i=1}^p (b_i + b'_i) T_i, \sum_{j=1}^r (c_j + c'_j) I_j, \sum_{k=1}^s (d_k + d'_k) F_k) \subseteq RNQ(H)$ as $(a + a') \cup (b_i + b'_i) \cup (c_j + c'_j) \cup (d_k + d'_k) \subseteq H$. Thus $\bar{x} \oplus RNQ(H) \subseteq RNQ(H)$. Let $\bar{y} = (a', \sum_{i=1}^p b'_i T_i, \sum_{j=1}^r c'_j I_j, \sum_{k=1}^s d'_k F_k) \in RNQ(H)$. Since $(H, +)$ satisfies the reproduction axiom and $a', b'_i, c'_j, d'_k \in H$, it follows that $a' \in a + H, b'_i \in b_i + H, c'_j \in c_j + H$ and $d'_k \in d_k + H$. The latter implies that there exist $a^*, b_i^*, c_j^*, d_k^* \in H$ such that $a' \in a + a^*, b'_i \in b_i + b_i^*, c'_j \in c_j + c_j^*$ and $d'_k \in d_k + d_k^*$. It is clear that $\bar{y} \in \bar{x} \oplus \bar{z}$ where $\bar{z} = (a^*, \sum_{i=1}^p b_i^* T_i, \sum_{j=1}^r c_j^* I_j, \sum_{k=1}^s d_k^* F_k) \in RNQ(H)$. Thus, $(RNQ(H), \oplus)$ satisfies the reproduction axiom.

Conversely, let $(RNQ(H), \oplus)$ be a quasi-hypergroup and $a \in H$. Since $0 \in H$, it follows that $\bar{a} = (a, \sum_{i=1}^p 0T_i, \sum_{j=1}^r 0I_j, \sum_{k=1}^s 0F_k) \in RNQ(H)$. Having $(RNQ(H), \oplus)$ a quasi-hypergroup implies that $\bar{a} \oplus RNQ(H) =$

$RNQ(H) \oplus \bar{a} = RNQ(H)$. The latter implies that $a + H = H + a = H$. □

Proposition 3.2. *Let $(H, +)$ be a hypergroupoid with $0 \in H$. Then $(RNQ(H), \oplus)$ is a semi-hypergroup with identity element $\bar{0} = (0, \sum_{i=1}^p 0T_i, \sum_{j=1}^r 0I_j, \sum_{k=1}^s 0F_k)$ if and only if $(H, +)$ is a semi-hypergroup with identity element 0.*

Proof. Let $(H, +)$ be a semi-hypergroup and $\bar{x}, \bar{y}, \bar{z} \in RNQ(H)$ with $\bar{x} = (a, \sum_{i=1}^p b_i T_i, \sum_{j=1}^r c_j I_j, \sum_{k=1}^s d_k F_k)$, $\bar{y} = (a', \sum_{i=1}^p b'_i T_i, \sum_{j=1}^r c'_j I_j, \sum_{k=1}^s d'_k F_k)$ and $\bar{z} = (a'', \sum_{i=1}^p b''_i T_i, \sum_{j=1}^r c''_j I_j, \sum_{k=1}^s d''_k F_k)$. Having $a + (a' + a'') = (a + a') + a''$, $b_i + (b'_i + b''_i) = (b_i + b'_i) + b''_i$, $c_j + (c'_j + c''_j) = (c_j + c'_j) + c''_j$ and $d_k + (d'_k + d''_k) = (d_k + d'_k) + d''_k$ implies that $\bar{x} \oplus (\bar{y} \oplus \bar{z}) = (\bar{x} \oplus \bar{y}) \oplus \bar{z}$.

Let $(RNQ(H), \oplus)$ be a semi-hypergroup and $a, b, c \in H$. Then $\bar{a}, \bar{b}, \bar{c} \in RNQ(H)$ with $\bar{a} = (a, \sum_{i=1}^p 0T_i, \sum_{j=1}^r 0I_j, \sum_{k=1}^s 0F_k)$,

$\bar{b} = (b, \sum_{i=1}^p 0T_i, \sum_{j=1}^r 0I_j, \sum_{k=1}^s 0F_k)$ and $\bar{c} = (c, \sum_{i=1}^p 0T_i, \sum_{j=1}^r 0I_j, \sum_{k=1}^s 0F_k)$. Having $\bar{a} \oplus (\bar{b} \oplus \bar{c}) = (\bar{a} \oplus \bar{b}) \oplus \bar{c}$ implies that $a + (b + c) = (a + b) + c$. □

Proposition 3.3. *Let $(H, +)$ be a hypergroupoid with $0 \in H$. Then $(RNQ(H), \oplus)$ is an H_v -semigroup with identity element $\bar{0} = (0, \sum_{i=1}^p 0T_i, \sum_{j=1}^r 0I_j, \sum_{k=1}^s 0F_k)$ if and only if $(H, +)$ is an H_v -semigroup with identity element 0.*

Proof. The proof is similar to that of Proposition 3.2 but instead of equality we have non-empty intersection. □

Theorem 3.4. *Let $(H, +)$ be a hypergroupoid. Then $(RNQ(H), \oplus)$ is a hypergroup with identity element $\bar{0} = (0, \sum_{i=1}^p 0T_i, \sum_{j=1}^r 0I_j, \sum_{k=1}^s 0F_k)$ if and only if $(H, +)$ is a hypergroup with identity element 0.*

Proof. The proof is direct from Propositions 3.1 and 3.2. □

Theorem 3.5. *Let $(H, +)$ be a hypergroupoid with $0 \in H$. Then $(RNQ(H), \oplus)$ is an H_v -group with identity element $\bar{0} = (0, \sum_{i=1}^p 0T_i, \sum_{j=1}^r 0I_j, \sum_{k=1}^s 0F_k)$ if and only if $(H, +)$ is an H_v -group with an identity element 0.*

Proof. The proof follows from Propositions 3.1 and 3.3. □

Theorem 3.6. *Let $(H, +)$ be a hypergroupoid. Then $(RNQ(H), \oplus)$ is a commutative hypergroup (H_v -group) with identity element $\bar{0} = (0, \sum_{i=1}^p 0T_i, \sum_{j=1}^r 0I_j, \sum_{k=1}^s 0F_k)$ if and only if $(H, +)$ is a commutative hypergroup (H_v -group) with an identity 0.*

Proof. The proof is straightforward. □

NOTATION 1. *Let $(H, +)$ be a hypergroup (H_v -group) with identity “0” satisfying $0+0 = 0$. Then $(RNQ(H), \oplus)$ is called a refined neutrosophic quadruple hypergroup (refined neutrosophic quadruple H_v -group).*

Corollary 3.7. Let $(H, +)$ be a hypergroup (H_v -group) containing an identity element 0 with the property that $0 + 0 = 0$. Then there are infinite number of refined neutrosophic quadruple hypergroups (H_v -groups).

Proof. Let $(H, +)$ be a hypergroup (H_v -group). Theorem 3.4 and Theorem 3.5 implies that $(RNQ(H), \oplus)$ is a neutrosophic quadruple hypergroup (H_v -group) with identity $\bar{0}$ and $\bar{0} \oplus \bar{0} = \bar{0}$. Applying Theorem 3.4 and Theorem 3.5 on $(RNQ(H), \oplus)$, we get $RNQ(RNQ(H))$ is a neutrosophic quadruple hypergroup (H_v -group). Continuing on this pattern, we get $RNQ(RNQ(\dots (RNQ(H)) \dots))$ is a neutrosophic quadruple hypergroup (H_v -group). \square

Proposition 3.8. Let X be any set with a hyperoperation “+”. Then $RNQ(X)$ is a cyclic refined neutrosophic quadruple hypergroup if and only if X is a cyclic hypergroup with an identity element “ $0 \in X$ ” and $0 + 0 = 0$.

Proof. Let X be a cyclic hypergroup with identity “ $0 \in X$ ” and $0 + 0 = 0$. Then there exist $a \in X$ such that a is a generator of X . It is clear that \bar{a} is a generator of $RNQ(X)$ where $\bar{a} = (a, \sum_{i=1}^p aT_i, \sum_{j=1}^r aI_j, \sum_{k=1}^s aF_k) \in RNQ(X)$.

Let $RNQ(X)$ be a cyclic quadruple hypergroup. Then there exist $\bar{x} \in RNQ(X)$ such that $\bar{x} = (a, \sum_{i=1}^p b_i T_i, \sum_{j=1}^r c_j I_j, \sum_{k=1}^s d_k F_k)$ is a generator of $RNQ(X)$. It is clear that a is a generator of X . \square

Example 3.9. Let T_1, T_2 be refinements of T , I_1, F_1 be refinements of I, F respectively, $H_1 = \{0, 1\}$ and define $(H_1, +_1)$ as follows:

$+_1$	0	1
0	0	1
1	1	H_1

Since (H_1, \oplus) is a commutative hypergroup with an identity 0 , it follows by Theorem 3.6 that $(RNQ(H_1), \oplus)$ is a commutative refined neutrosophic quadruple hypergroup with 32 elements and identity $\bar{0} = (0, 0T_1 + 0T_2, 0I_1, 0F_1)$. Moreover, having $H_1 = 1 +_1 1$ implies that 1 is a generator of $(H_1, +)$ and $(H_1, +)$ is a single-power cyclic hypergroup of period 2. Theorem 3.8 asserts that $(RNQ(H_1), \oplus)$ is a single power cyclic hypergroup of period 2 and the generator element is $(1, 1T_1 + 1T_2, 1I_1, 1F_1)$.

It is clear that $(1, 0T_1 + 0T_2, 1I_1, 1F_1) \oplus (1, 0T_1 + 1T_2, 0I_1, 1F_1) = \{(1, 0T_1 + 1T_2, 1I_1, 0F_1), (1, 0T_1 + 1T_2, 1I_1, 1F_1)\}$.

Definition 3.10. Let $(H, +)$ be a hypergroup (H_v -group). A subset X of $RNQ(H)$ with the property that $\bar{0} = (0, \sum_{i=1}^p 0T_i, \sum_{j=1}^r 0I_j, \sum_{k=1}^s 0F_k) \in X$ is called a refined neutrosophic subhypergroup (H_v -subgroup) of $RNQ(H)$ if there exists $S \subseteq H$ such that $X = RNQ(S)$ and (X, \oplus) is a refined neutrosophic quadruple hypergroup (H_v -group).

Proposition 3.11. Let $(H, +)$ be a hypergroup (H_v -group) and $S \subseteq H$. A subset $X = RNQ(S) \subseteq RNQ(H)$ is a refined neutrosophic subhypergroup (H_v -subgroup) of $RNQ(H)$ if the following conditions are satisfied:

1. $\bar{0} = (0, \sum_{i=1}^p 0T_i, \sum_{j=1}^r 0I_j, \sum_{k=1}^s 0F_k) \in X$;
2. $\bar{x} \oplus X = X \oplus \bar{x} = X$ for all $\bar{x} \in X$.

Proof. The proof is straightforward. □

Theorem 3.12. Let $(H, +)$ be a hypergroup (H_v -group) with identity “0”, $S \subseteq H$ and $0 \in S$. Then $(RNQ(S), \oplus)$ is a refined neutrosophic quadruple subhypergroup (H_v -subgroup) of $(RNQ(H), \oplus)$ if and only if $(S, +)$ is a subhypergroup (H_v -subgroup) of $(H, +)$.

Proof. The proof is straightforward by applying Proposition 3.11. □

Example 3.13. Since $(H_1, +_1)$ in Example 3.9 has only two subhypergroups ($\{0\}$ and H_1), it follows by applying Theorem 3.12 that $(RNQ(H_1), \oplus)$ has only two refined neutrosophic quadruple subhypergroups: $(\{\bar{0}\}, \oplus) = (RNQ(\{0\}), \oplus)$ and $(RNQ(H_1), \oplus)$.

Example 3.14. Let $H_2 = \{0, 1, 2, 3\}$ and define $(H_2, +_2)$ as follows:

$+_2$	0	1	2	3
0	0	1	2	3
1	1	2	3	0
2	2	3	$\{0, 2\}$	1
3	3	0	1	2

It is clear that $(H_2, +_2)$ is a commutative H_v -group that has exactly three non-isomorphic H_v -subgroups containing 0: $\{0\}$, $\{0, 2\}$ and H_2 . We can deduce by Theorem 3.12 that $(RNQ(H_2), \oplus)$ is a commutative refined neutrosophic quadruple H_v -group and has three non-isomorphic refined neutrosophic quadruple H_v -subgroups: $RNQ(\{0\}) = \{\bar{0}\}$, $RNQ(\{0, 2\})$ and $RNQ(H_2)$.

Proposition 3.15. Let $(H, +)$ be a hypergroup and $(S, +)$ be a subhypergroup of $(H, +)$ containing 0. Then $RNQ(S) \oplus RNQ(S) = RNQ(S)$.

Proof. The proof is straightforward. □

Definition 3.16. Let $(RNQ(H), \oplus_1)$ and $(RNQ(J), \oplus_2)$ be refined neutrosophic quadruple hypergroups with $0_H \in H$ and $0_J \in J$. A function $\phi : RNQ(H) \rightarrow RNQ(J)$ is called *refined neutrosophic homomorphism* if the following conditions are satisfied:

1. $\phi(0_H, \sum_{i=1}^p 0_H T_i, \sum_{j=1}^r 0_H I_j, \sum_{k=1}^s 0_H F_k) = (0_J, \sum_{i=1}^p 0_J T_i, \sum_{j=1}^r 0_J I_j, \sum_{k=1}^s 0_J F_k)$;
2. $\phi(x \oplus_1 y) = \phi(x) \oplus_2 \phi(y)$ for all $x, y \in RNQ(H)$.

If ϕ is a refined neutrosophic bijective homomorphism then it is called *refined neutrosophic isomorphism* and we write $RNQ(H) \cong RNQ(J)$.

Example 3.17. Let $(H, +)$ be a hypergroup. Then the function $f : RNQ(H) \rightarrow RNQ(H)$ is an isomorphism, where $f(x) = x$ for all $x \in RNQ(H)$.

Example 3.18. Let $(H, +)$ be a hypergroup and $0 \in H$ and $f : RNQ(H) \rightarrow RNQ(H)$ be defined as follows:

$$f\left(\left(a, \sum_{i=1}^p b_i T_i, \sum_{j=1}^r c_j I_j, \sum_{k=1}^s d_k F_k\right)\right) = \left(a, \sum_{i=1}^p 0T_i, \sum_{j=1}^r 0I_j, \sum_{k=1}^s 0F_k\right).$$

Then f is a refined neutrosophic homomorphism.

Proposition 3.19. Let $(H, +_1)$ and $(J, +_2)$ be hypergroups with $0_H \in H, 0_J \in J$. If there exist a homomorphism $f : H \rightarrow J$ with $f(0_H) = 0_J$ then there exist a refined neutrosophic homomorphism from $(RNQ(H), \oplus_1)$ to $(RNQ(J), \oplus_2)$.

Proof. Let $\phi : RNQ(H) \rightarrow RNQ(J)$ be defined as follows:

$$\phi\left(\left(a, \sum_{i=1}^p b_i T_i, \sum_{j=1}^r c_j I_j, \sum_{k=1}^s d_k F_k\right)\right) = \left(f(a), \sum_{i=1}^p f(b_i)T_i, \sum_{j=1}^r f(c_j)I_j, \sum_{k=1}^s f(d_k)F_k\right).$$

It is clear that ϕ is a refined neutrosophic homomorphism. □

Corollary 3.20. Let $(H, +_1)$ and $(J, +_2)$ be isomorphic hypergroups with $0_H \in H, 0_J \in J$. Then $(RNQ(H), \oplus_1)$ and $(RNQ(J), \oplus_2)$ are isomorphic refined neutrosophic quadruple hypergroups.

Proof. The proof is straightforward by using Proposition 3.19. □

Definition 3.21. Let $(H, +)$ be a commutative hypergroup with an identity element “0” and $S \subseteq H$ be a subhypergroup of H . Then $(H/S, +')$ is a hypergroup with: S as an identity element and $S +' S = S$. Here “+’” is defined as follows: For all $x, y \in H$,

$$(x + S) +' (y + S) = (x + y) + S.$$

Proposition 3.22. Let $(S, +)$ be a subhypergroup of a commutative hypergroup $(H, +)$. Then $(RNQ(H/S), \oplus)$ is a hypergroup.

Proof. Since $(H, +)$ is commutative, it follows that “+’” is well defined. The proof follows from having $(H/S, +')$ a hypergroup with S as an identity, $S +' S = S$ and from Theorem 3.4. □

Proposition 3.23. Let $(S, +)$ be a subhypergroup of a commutative hypergroup $(H, +)$. Then $(RNQ(H/S), \oplus) \cong (RNQ(H)/RNQ(S), \oplus')$.

Proof. Let $g : RNQ(H)/RNQ(S) \rightarrow RNQ(H/S)$ be defined as follows:

$$\begin{aligned} g\left(\left(a, \sum_{i=1}^p b_i T_i, \sum_{j=1}^r c_j I_j, \sum_{k=1}^s d_k F_k\right) \oplus RNQ(S)\right) \\ = \left(a + S, \sum_{i=1}^p (b_i + S)T_i, \sum_{j=1}^r (c_j + S)I_j, \sum_{k=1}^s (d_k + S)F_k\right). \end{aligned}$$

Then g is a hypergroup isomorphism. This can be proved easily by applying a similar proof to that of Proposition 3.27 that was done by the authors in [9]. □

Example 3.24. Let $H_3 = \{0, 1, 2, 3, 4\}$ and define “+” on H_3 as follows: $x + y = \{x, y\}$ for all $x, y \in H_3$. It is clear that $\{0\}$, $\{0, 1\}$, $\{0, 1, 2\}$, $\{0, 1, 2, 3\}$ and H_3 are the only non-isomorphic subhypergroups of H_3 . By applying Proposition 3.23, we get $RNQ(H_3/\{0, 1\}) \cong RNQ(H_3)/RNQ(\{0, 1\})$, $RNQ(H_3/\{0, 1, 2\}) \cong RNQ(H_3)/RNQ(\{0, 1, 2\})$ and $RNQ(H_3/\{0, 1, 2, 3\}) \cong RNQ(H_3)/RNQ(\{0, 1, 2, 3\})$.

4 Ordered refined neutrosophic quadruple hypergroups

In this section, an order on refined neutrosophic quadruple hypergroups is defined and some examples and results on refined neutrosophic quadruple partially ordered hypergroups (po-hypergroups) are presented.

A partial order relation on a set X (Poset) is a binary relation “ \leq ” on X which satisfies conditions reflexivity, antisymmetry and transitivity.

Let (H, \leq) be a partial ordered set and define $(RNQ(H), \trianglelefteq)$ as follows:

$$(a, \sum_{i=1}^p b_i T_i, \sum_{j=1}^r c_j I_j, \sum_{k=1}^s d_k F_k) \trianglelefteq (a', \sum_{i=1}^p b'_i T_i, \sum_{j=1}^r c'_j I_j, \sum_{k=1}^s d'_k F_k)$$

if and only if $a \leq a', b_i \leq b'_i, c_j \leq c'_j$ and $d_k \leq d'_k$. It is clear that $(RNQ(H), \trianglelefteq)$ is a partial ordered set.

Definition 4.1. [16] An algebraic hyperstructure (H, \circ, \leq) is called a *partially ordered hypergroup* or *po-hypergroup*, if (H, \circ) is a hypergroup and \leq is a partial order relation on H such that the monotone condition holds as follows:

$$x \leq y \Rightarrow a \circ x \leq a \circ y \text{ for all } a, x, y \in H.$$

Let A, B be non-empty subsets of (H, \leq) . The inequality $A \leq B$ means that for any $a \in A$, there exist $b \in B$ such that $a \leq b$.

Theorem 4.2. Let $(H, +)$ be a hypergroupoid. Then $(RNQ(H), \oplus, \trianglelefteq)$ is a refined neutrosophic quadruple po-hypergroup with identity element $\bar{0} = (0, \sum_{i=1}^p 0T_i, \sum_{j=1}^r 0I_j, \sum_{k=1}^s 0F_k)$ if and only if $(H, +, \leq)$ is a po-hypergroup with identity element 0.

Proof. Let $(H, +, \leq)$ be a po-hypergroup, $\bar{e} = (e, \sum_{i=1}^p f_i T_i, \sum_{j=1}^r c_j I_j, \sum_{k=1}^s h_k F_k) \in RNQ(H)$ and

$(a, \sum_{i=1}^p b_i T_i, \sum_{j=1}^r c_j I_j, \sum_{k=1}^s d_k F_k) \trianglelefteq (a', \sum_{i=1}^p b'_i T_i, \sum_{j=1}^r c'_j I_j, \sum_{k=1}^s d'_k F_k)$. We need to show that:

$$\bar{e} \oplus (a, \sum_{i=1}^p b_i T_i, \sum_{j=1}^r c_j I_j, \sum_{k=1}^s d_k F_k) \trianglelefteq \bar{e} \oplus (a', \sum_{i=1}^p b'_i T_i, \sum_{j=1}^r c'_j I_j, \sum_{k=1}^s d'_k F_k).$$

Having $a \leq a', b_i \leq b'_i, c_j \leq c'_j, d_k \leq d'_k$ and $(H, +, \leq)$ a po-hypergroup implies that $e + a \leq e + a', f_i + b_i \leq f_i + b'_i, g_j + c_j \leq g_j + c'_j$ and $h_k + d_k \leq h_k + d'_k$. Let $\bar{a}^* = (a^*, \sum_{i=1}^p b_i^* T_i, \sum_{j=1}^r c_j^* I_j, \sum_{k=1}^s d_k^* F_k) \in$

$\bar{e} \oplus (a, \sum_{i=1}^p b_i T_i, \sum_{j=1}^r c_j I_j, \sum_{k=1}^s d_k F_k)$. Then $a^* \in e + a, b_i^* \in f_i + b_i, c_j^* \in g_j + c_j$ and $d_k^* \in h_k + d_k$. Having $e + a \leq e + a', f_i + b_i \leq f_i + b'_i, g_j + c_j \leq g_j + c'_j$ and $h_k + d_k \leq h_k + d'_k$ implies that there exist $a^{*'} \in e + a', b_i^{*'} \in f_i + b'_i, c_j^{*'} \in g_j + c'_j$ and $d_k^{*'} \in h_k + d'_k$ such that $a^* \leq a^{*'}, b_i^* \leq b_i^{*'}, c_j^* \leq c_j^{*'}$ and $d_k^* \leq d_k^{*'}$. We get now that $\bar{a}^* \trianglelefteq \bar{a}^{*'}$ where $\bar{a}^{*' } = (a^{*' }, \sum_{i=1}^p b_i^{*' } T_i, \sum_{j=1}^r c_j^{*' } I_j, \sum_{k=1}^s d_k^{*' } F_k)$ and $\bar{a}^{*' } \in \bar{e} \oplus (a', \sum_{i=1}^p b'_i T_i, \sum_{j=1}^r c'_j I_j, \sum_{k=1}^s d'_k F_k)$.

Let $a, b, e \in H$ and $a \leq b$. Having $0 \leq 0$ implies that

$$(a, \sum_{i=1}^p 0T_i, \sum_{j=1}^r 0I_j, \sum_{k=1}^s 0F_k) \trianglelefteq (b, \sum_{i=1}^p 0T_i, \sum_{j=1}^r 0I_j, \sum_{k=1}^s 0F_k).$$

Since $(RNQ(H), \oplus, \trianglelefteq)$ is a refined neutrosophic quadruple po-hypergroup, it follows that for

$$\bar{e} = (e, \sum_{i=1}^p 0T_i, \sum_{j=1}^r 0I_j, \sum_{k=1}^s 0F_k),$$

$$\bar{e} \oplus (a, \sum_{i=1}^p 0T_i, \sum_{j=1}^r 0I_j, \sum_{k=1}^s 0F_k) \trianglelefteq \bar{e} \oplus (b, \sum_{i=1}^p 0T_i, \sum_{j=1}^r 0I_j, \sum_{k=1}^s 0F_k).$$

It is clear that $e + a \leq e + b$. □

Corollary 4.3. *Let $(H, +, \leq)$ be a po-hypergroup containing an identity element 0 with the property that $0 + 0 = 0$. Then there is infinite number of refined neutrosophic quadruple po-hypergroups.*

Proof. The proof is straightforward using Theorem 4.2. □

Example 4.4. Let $H_1 = \{0, 1\}$ and define $(H_1, +_1)$ as in Example 3.9. It is clear that $(H_1, +_1, \leq)$ is a po-hypergroup. Here, the partial order relation “ \leq ” is directed to the set $\{(0, 0), (1, 1)\}$. By using Theorem 4.2, we get $(RNQ(H_1), \oplus, \trianglelefteq)$ is a refined neutrosophic quadruple po-hypergroup.

Example 4.5. Let (H, \leq) be any poset and define $(H, +)$ as the biset hypergroup, i.e. $x + y = \{x, y\}$ for all $x, y \in H$. Then $(RNQ(H), \oplus, \trianglelefteq)$ is a refined neutrosophic quadruple po-hypergroup.

Theorem 4.6. [16] *Let (H, \circ) be a hypergroup such that there exists an element $0 \in H$ and the following conditions hold:*

1. $0 \circ 0 = 0$;
2. $\{0, x\} \subseteq 0 \circ x$ for all $x \in H$;
3. If $x \circ 0 = y \circ 0$ then $x = y$ for all $x, y \in H$.

Then there exist a relation “ \leq ” on H such that (H, \circ, \leq) is a po-hypergroup.

Heidari et al. [16], in their proof of Theorem 4.6, defined the binary relation “ \leq ” on H as follows:
 $x \leq y \iff x \in y \circ 0$, for all $x, y \in H$.

Corollary 4.7. *Let $(H, +)$ be a hypergroup satisfying conditions of Theorem 4.6. Then there exist a relation “ \trianglelefteq ” on $RNQ(H)$ such that $(RNQ(H), \oplus, \trianglelefteq)$ is a refined neutrosophic quadruple po-hypergroup.*

Proof. The proof follows from Theorems 4.2 and 4.6. □

Example 4.8. Let $H = \{0, x, y\}$ and define “+” by the following table:

+	0	x	y
0	0	$\{0, x\}$	H
c	$\{0, x\}$	H	H
y	H	H	H

Then $(RNQ(H), \oplus, \trianglelefteq)$ is a refined neutrosophic quadruple po-hypergroup. Here the partial order relation “ \leq ” is directed to the set $\{(0, 0), (x, x), (y, y), (x, y), (0, x), (0, y)\}$ and \trianglelefteq is defined in the usual way on $RNQ(H)$.

Definition 4.9. Let $(RNQ(H), \oplus_1, \trianglelefteq_1)$ and $(RNQ(J), \oplus_2, \trianglelefteq_2)$ be refined neutrosophic quadruple po-hypergroups. A function $\phi : RNQ(H) \rightarrow RNQ(J)$ is called an *ordered refined neutrosophic homomorphism* if the following conditions hold:

1. $\phi(0_H, \sum_{i=1}^p 0_H T_i, \sum_{j=1}^r p 0_H I_j, \sum_{k=1}^s 0_H F_k) = (0_J, \sum_{i=1}^p 0_J T_i, \sum_{j=1}^r 0_J I_j, \sum_{k=1}^s 0_J F_k)$;
2. $\phi(x \oplus_1 y) = \phi(x) \oplus_2 \phi(y)$ for all $x, y \in RNQ(H)$;
3. if $x \trianglelefteq_1 y$ then $\phi(x) \trianglelefteq_2 \phi(y)$ for all $x, y \in RNQ(H)$.

If ϕ is an ordered refined neutrosophic homomorphism and is bijective then it is called an *ordered refined neutrosophic isomorphism* and we say $RNQ(H)$ and $RNQ(J)$ are isomorphic refined neutrosophic quadruple po-hypergroups.

Example 4.10. Let $(H, +, \leq)$ be a po-hypergroup. Then $f : RNQ(H) \rightarrow RNQ(H)$ is an ordered refined neutrosophic isomorphism, where $f(x) = x$ for all $x \in RNQ(H)$.

Example 4.11. Let $(H, +, \leq)$ be a po-hypergroup, $0 \in H$ and $f : RNQ(H) \rightarrow RNQ(H)$ be defined as follows:

$$f((a, \sum_{i=1}^p b_i T_i, \sum_{j=1}^r c_j I_j, \sum_{k=1}^s d_k F_k)) = (a, \sum_{i=1}^p 0 T_i, \sum_{j=1}^r 0 I_j, \sum_{k=1}^s 0 F_k).$$

Then f is an ordered refined neutrosophic homomorphism.

Example 4.12. Let $(H, +_1, \leq_1)$ and $(J, +_2, \leq_2)$ be po-hypergroups, $0_H \in H, 0_J \in J$ and $g : H \rightarrow J$ be an ordered homomorphism. Then $f : RNQ(H) \rightarrow RNQ(J)$ is an ordered refined neutrosophic homomorphism. Here, f is defined as follows:

$$f((a, \sum_{i=1}^p b_i T_i, \sum_{j=1}^r c_j I_j, \sum_{k=1}^s d_k F_k)) = (g(a), \sum_{i=1}^p 0_J T_i, \sum_{j=1}^r 0_J I_j, \sum_{k=1}^s 0_J F_k).$$

Proposition 4.13. Let $(H, +_1, \leq_1)$ and $(J, +_2, \leq_2)$ be po-hypergroups with $0_H \in H, 0_J \in J$. If there exist an ordered homomorphism $f : H \rightarrow J$ with $f(0_H) = 0_J$ then there exist an ordered refined neutrosophic homomorphism from $(RNQ(H), \oplus_1, \trianglelefteq_1)$ to $(RNQ(J), \oplus_2, \trianglelefteq_2)$.

Proof. Let $\phi : RNQ(H) \rightarrow RNQ(J)$ be defined as follows:

$$\phi((a, \sum_{i=1}^p b_i T_i, \sum_{j=1}^r c_j I_j, \sum_{k=1}^s d_k F_k)) = (f(a), \sum_{i=1}^p f(b_i) T_i, \sum_{j=1}^r f(c_j) I_j, \sum_{k=1}^s f(d_k) F_k).$$

It is clear that ϕ is an ordered refined neutrosophic homomorphism. □

Corollary 4.14. Let $(H, +_1, \leq_1)$ and $(J, +_2, \leq_2)$ be isomorphic po-hypergroup with $0_H \in H, 0_J \in J$. Then $(RNQ(H), \oplus_1, \trianglelefteq_1)$ and $(RNQ(J), \oplus_2, \trianglelefteq_2)$ are isomorphic refined neutrosophic quadruple po-hypergroups.

Proof. The proof is straightforward by using Proposition 4.13. □

5 Fundamental group of refined neutrosophic quadruple hypergroups

This section presents the study of fundamental relation on refined neutrosophic quadruple hypergroups and finds their fundamental refined quadruple neutrosophic groups.

Theorem 5.1. *Let $(G, +)$ be a groupoid. Then $(RNQ(G), \oplus)$ is a group with identity element*

$\bar{0} = (0, \sum_{i=1}^p 0T_i, \sum_{j=1}^r 0I_j, \sum_{k=1}^s 0F_k)$ *if and only if $(G, +)$ is a group with identity element 0.*

Proof. It is clear that $\bar{0} = (0, \sum_{i=1}^p 0T_i, \sum_{j=1}^r 0I_j, \sum_{k=1}^s 0F_k)$ is the identity of $(RNQ(G), \oplus)$ if and only if 0 is the

identity of G . Let $\bar{x} = (a, \sum_{i=1}^p b_i T_i, \sum_{j=1}^r c_j I_j, \sum_{k=1}^s d_k F_k) \in RNQ(G)$. Then the inverse

$-\bar{x} = (a, \sum_{i=1}^p (-b_i) T_i, \sum_{j=1}^r (-c_j) I_j, \sum_{k=1}^s (-d_k) F_k)$ of \bar{x} exists if and only if the inverse $-y$ of y exists in G for all

$y \in G$. The proof of $(RNQ(G), \oplus)$ is binary closed if and only if $(G, +)$ is binary closed is similar to that of Proposition 3.1. And the proof of $(RNQ(G), \oplus)$ is associative if and only if $(G, +)$ is associative is similar to that of Proposition 3.2. \square

NOTATION 2. *Let $(G, +)$ be a group with identity element "0". Then $(RNQ(G), \oplus)$ is called refined neutrosophic quadruple group.*

Proposition 5.2. *Let G, G' be isomorphic groups. Then $RNQ(G)$ and $RNQ(G')$ are isomorphic neutrosophic quadruple groups.*

Definition 5.3. [14] For all $n > 1$, we define the relation β_n on a semihypergroup (H, \circ) as follows:

$$x\beta_n y \text{ if there exist } a_1, \dots, a_n \text{ in } H \text{ such that } \{x, y\} \subseteq \prod_{i=1}^n a_i$$

Here, $\prod_{i=1}^n a_i = a_1 \circ a_2 \dots \circ a_n$. And we set $\beta = \bigcup_{n \geq 1} \beta_n$, where $\beta_1 = \{(x, x) \mid x \in H\}$ is the diagonal relation on H .

Koskas [18] introduced this relation as an important tool to connect hypergroups with groups. And due to it's importance in connecting algebraic hyperstructures with algebraic structures, different researchers studied it on various hypergroups and some extended this definition to cover other types of hyperstructures.

Clearly, the relation β is reflexive and symmetric. Denote by β^* the transitive closure of β . Then β^* is called the *fundamental equivalence relation* on H and it is the smallest strongly regular relation on H . If H is a hypergroup then $\beta = \beta^*$ and H/β^* is called the *fundamental group*.

Throughout this section, β and β^* are the relation on H and β_N and β_N^* are the relations on $RNQ(H)$.

Theorem 5.4. *Let $(H, +)$ be a hypergroup with identity element element "0" and $0 + 0 = 0$ and let $a, a', b_i, b'_i, c_j, c'_j, d_k, d'_k \in H$. Then*

$(a, \sum_{i=1}^p b_i T_i, \sum_{j=1}^r c_j I_j, \sum_{k=1}^s d_k F_k) \beta_N (a', \sum_{i=1}^p b'_i T_i, \sum_{j=1}^r c'_j I_j, \sum_{k=1}^s d'_k F_k)$ *if and only if $a\beta a', b_i\beta b'_i, c_j\beta c'_j$ and $d_k\beta d'_k$.*

Proof. Let $(a, \sum_{i=1}^p b_i T_i, \sum_{j=1}^r c_j I_j, \sum_{k=1}^s d_k F_k) \beta_N (a', \sum_{i=1}^p b'_i T_i, \sum_{j=1}^r c'_j I_j, \sum_{k=1}^s d'_k F_k)$. Then there exist

$(a_t, \sum_{i=1}^p b_{it} T_i, \sum_{j=1}^r c_{jt} I_j, \sum_{k=1}^s d_{kt} F_k)$ with $t = 1, \dots, n$ such that

$$\{(a, \sum_{i=1}^p b_i T_i, \sum_{j=1}^r c_j I_j, \sum_{k=1}^s d_k F_k), (a', \sum_{i=1}^p b'_i T_i, \sum_{j=1}^r c'_j I_j, \sum_{k=1}^s d'_k F_k)\}$$

is a subset of

$$(a_1, \sum_{i=1}^p b_{i1} T_i, \sum_{j=1}^r c_{j1} I_j, \sum_{k=1}^s d_{k1} F_k) \oplus \dots \oplus (a_n, \sum_{i=1}^p b_{in} T_i, \sum_{j=1}^r c_{jn} I_j, \sum_{k=1}^s d_{kn} F_k).$$

The latter implies that $a, a' \in a_1 + \dots + a_n$, $b_i, b'_i \in b_{i1} + \dots + b_{in}$, $c_j, c'_j \in c_{j1} + \dots + c_{jn}$ and $d_k, d'_k \in d_{k1} + \dots + d_{kn}$. Thus, $a\beta a'$, $b_i\beta b'_i$, $c_j\beta c'_j$ and $d_k\beta d'_k$.

Conversely, let $a\beta a'$, $b_i\beta b'_i$, $c_j\beta c'_j$ and $d_k\beta d'_k$. Then there exist $t_1, t_2, t_3, t_4 \in \mathbb{N}$ and $x_1, \dots, x_{t_1}, y_{i1}, \dots, y_{it_2}, z_{j1}, \dots, z_{jt_3}, w_{k1}, \dots, w_{kt_4} \in H$ such that $a, a' \in x_1 + \dots + x_{t_1}$, $b_i, b'_i \in y_{i1} + \dots + y_{it_2}$, $c_j, c'_j \in z_{j1} + \dots + z_{jt_3}$ and $d_k, d'_k \in w_{k1} + \dots + w_{kt_4}$. By setting $t = \max\{t_1, t_2, t_3, t_4\}$ and $x_m = 0$ for $t_1 < m \leq t$, $y_{im} = 0$ for $t_2 < m \leq t$, $z_{jm} = 0$ for $t_3 < m \leq t$ and $w_{km} = 0$ for $t_4 < m \leq t$ and using the fact that $e \in 0 + e \cap e + 0$ for all $e \in H$, we get $a, a' \in x_1 + \dots + x_{t_1}$, $b_i, b'_i \in y_{i1} + \dots + y_{it}$, $c_j, c'_j \in z_{j1} + \dots + z_{jt}$ and $d_k, d'_k \in w_{k1} + \dots + w_{kt}$. The latter implies that $\{(a, \sum_{i=1}^p b_i T_i, \sum_{j=1}^r c_j I_j, \sum_{k=1}^s d_k F_k), (a', \sum_{i=1}^p b'_i T_i, \sum_{j=1}^r c'_j I_j, \sum_{k=1}^s d'_k F_k)\}$ is a subset of $(x_1, \sum_{i=1}^p y_{i1} T_i, \sum_{j=1}^r z_{j1} I_j, \sum_{k=1}^s w_{k1} F_k) \oplus \dots \oplus (x_t, \sum_{i=1}^p y_{it} T_i, \sum_{j=1}^r z_{jt} I_j, \sum_{k=1}^s w_{kt} F_k)$. Thus,

$$(a, \sum_{i=1}^p b_i T_i, \sum_{j=1}^r c_j I_j, \sum_{k=1}^s d_k F_k) \beta_N (a', \sum_{i=1}^p b'_i T_i, \sum_{j=1}^r c'_j I_j, \sum_{k=1}^s d'_k F_k).$$

□

Theorem 5.5. Let $(H, +)$ be a hypergroup with identity “0” and $0 + 0 = 0$. Then $RNQ(H)/\beta_N \cong RNQ(H/\beta)$.

Proof. Let $\phi : RNQ(H)/\beta_N \rightarrow RNQ(H/\beta)$ be defined as

$$\phi(\beta_N((a, \sum_{i=1}^p b_i T_i, \sum_{j=1}^r c_j I_j, \sum_{k=1}^s d_k F_k))) = (\beta(a), \sum_{i=1}^p \beta(b_i) T_i, \sum_{j=1}^r \beta(c_j) I_j, \sum_{k=1}^s \beta(d_k) F_k).$$

Theorem 5.4 asserts that ϕ is well-defined and one-to-one. Also, it is clear that ϕ is onto. We need to show that ϕ is a group homomorphism. Let

$\bar{a} = (a, \sum_{i=1}^p b_i T_i, \sum_{j=1}^r c_j I_j, \sum_{k=1}^s d_k F_k)$ and $\bar{a}' = (a', \sum_{i=1}^p b'_i T_i, \sum_{j=1}^r c'_j I_j, \sum_{k=1}^s d'_k F_k)$. Since $\beta_N(\bar{a}) \boxplus \beta_N(\bar{a}') = \beta_N(\bar{x})$ where $\bar{x} = (x, \sum_{i=1}^p y_i T_i, \sum_{j=1}^r z_j I_j, \sum_{k=1}^s w_k F_k) \in (a, \sum_{i=1}^p b_i T_i, \sum_{j=1}^r c_j I_j, \sum_{k=1}^s d_k F_k) \oplus (a', \sum_{i=1}^p b'_i T_i, \sum_{j=1}^r c'_j I_j, \sum_{k=1}^s d'_k F_k) =$

$(a + a', \sum_{i=1}^p (b_i + b'_i)T_i, \sum_{j=1}^r (c_j + c'_j)I_j, \sum_{k=1}^s (d_k + d'_k)F_k)$, it follows that

$$\phi(\beta_N(\bar{a}) \boxplus \beta_N(\bar{a}')) = \phi(\bar{x}) = (\beta(x), \sum_{i=1}^p \beta(y_i)T_i, \sum_{j=1}^r \beta(z_j)I_j, \sum_{k=1}^s \beta(w_k)F_k).$$

Having $\beta(x) = \beta(a) \oplus \beta(a')$, $\beta(y_i) = \beta(b_i) \oplus \beta(b'_i)$, $\beta(z_j) = \beta(c_j) \oplus \beta(c'_j)$ and $\beta(w_k) = \beta(d_k) \boxplus \beta(d'_k)$ imply that $\phi(\beta_N(\bar{a}) \boxplus \beta_N(\bar{a}')) = \phi(\beta_N(\bar{a})) \oplus' \phi(\beta_N(\bar{a}'))$. \square

Corollary 5.6. *Let $(H, +)$ be a hypergroup with identity element "0" and $0 + 0 = 0$. If G is the fundamental group of H (up to isomorphism) then $RNQ(G)$ is the fundamental group of $RNQ(H)$ (up to isomorphism).*

Proof. The proof follows from Proposition 5.2 and Theorem 5.5. \square

Corollary 5.7. *Let $(H, +)$ be a hypergroup with identity element "0" and $0 + 0 = 0$. If H has a trivial fundamental group then $RNQ(H)$ has a trivial fundamental group.*

Proof. The proof is straightforward by applying Corollary 5.6. \square

Theorem 5.8. [8] *Every single power cyclic hypergroup has a trivial fundamental group.*

Corollary 5.9. *Let $(H, +)$ be a single power cyclic hypergroup with $0 \in H$ and $0 + 0 = 0$. Then $RNQ(H)$ has a trivial fundamental group.*

Proof. The proof follows from Corollary 5.7 and Theorem 5.8. \square

6 Conclusion

This paper contributed to the study of neutrosophic hyperstructures by introducing refined neutrosophic quadruple hypergroups (po-hypergroups) and determining their fundamental refined neutrosophic quadruple groups. Several interesting results related to these new hypergroups were obtained. For future work, it will be interesting to study new properties of other types of refined neutrosophic quadruple hyperstructures.

References

- [1] M. Abdel-Basset, M. Saleh, A. Gamal and F. Smarandache, *An approach of TOPSIS technique for developing supplier selection with group decision making under type-2 neutrosophic number*, Applied Soft Computing, 77(2019), 438-452.
- [2] M. Abdel-Baset, G. Manogaran, A. Gamal and F. Smarandache, *A group decision making framework based on neutrosophic TOPSIS approach for smart medical device selection*, Journal of medical systems, 43(2)(2019), 38.
- [3] M. Abdel-Baset, V. Chang, A. Gamal, *Evaluation of the green supply chain management practices: A novel neutrosophic approach*, Computers in Industry, 108(2019), 210-220.
- [4] M. Abdel-Baset, G. Manogaran, A. Gamal and F. Smarandache, *A hybrid approach of neutrosophic sets and DEMATEL method for developing supplier selection criteria*, Design Automation for Embedded Systems, 2018, 1-22.
- [5] A. A. A. Agboola, B. Davvaz and F. Smarandache, *Neutrosophic quadruple algebraic hyperstructures*, Ann. Fuzzy Math. Inform. 14(2017), 29-42.

- [6] S. A. Akinleye, F. Smarandache and A. A. A. Agboola, *On neutrosophic quadruple algebraic structures*, Neutrosophic Sets Syst. 12(2016), 122-126.
- [7] R. Alhabib, M. M. Ranna, H. Farah, A.A. Salama, *Some Neutrosophic Probability Distributions*, Neutrosophic Sets and Systems, 22(2018), 30-38.
- [8] M. Al-Tahan and B. Davvaz, *On some properties of single power cyclic hypergroups and regular relations*, Journal of Algebra and Its Applications 16(11) (2017).
- [9] M. Al-Tahan and B. Davvaz, *On Neutrosophic Quadruple H_v -groups and their properties*, Southeast Asian Bulletin of Mathematics, accepted.
- [10] M. Al-Tahan and B. Davvaz, *Fundamental group and complete parts of Neutrosophic Quadruple H_v -groups*, submitted.
- [11] T. Chalapathi, R. V. Kiran Kumar, *Neutrosophic Units of Neutrosophic Rings and Fields*, Neutrosophic Sets and Systems, 21(2018), 5-12.
- [12] P. Corsini and V. Leoreanu, *Applications of hyperstructures theory*, Advances in Mathematics, Kluwer Academic Publisher, 2003.
- [13] B. Davvaz, *Polygroup Theory and Related Systems*, World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2013. viii+200 pp.
- [14] B. Davvaz, *Semihypergroup Theory*, Elsevier/Academic Press, London, 2016. viii+156 pp.
- [15] A. Elhassouny, F. Smarandache, *Neutrosophic modifications of Simplified TOPSIS for Imperfect Information (nS-TOPSIS)*, Neutrosophic Sets and Systems, 24(2019),100-113.
- [16] D. Heidari and B. Davvaz, *On ordered hyperstructures*, U.P.B. Sci. Bull. Series A 73(2)(2011), 85-96.
- [17] Y. B. Jun , S-Z. Song, F. Smarandache and H.Bordbar, *Neutrosophic Quadruple BCK/BCI-Algebras*, Axioms 7(41)(2018), doi:10.3390/axioms7020041.
- [18] M. Koskas, *Groupoides, demi-hypergroupes et hypergroupes*, J. Math. Pure Appl.,9(49)(1970), 155-192.
- [19] F. Marty, *Sur une generalization de la notion de group*, In 8th Congress Math. Scandenaves, (1934), 45-49.
- [20] N. A. Nabeeh, F. Smarandache, M. Abdel-Basset, H. A. El-Ghareeb and A. Aboelfetouh, *An Integrated Neutrosophic-TOPSIS Approach and Its Application to Personnel Selection: A New Trend in Brain Processing and Analysis*, IEEE Access, 7(2019), 29734-29744.
- [21] S. Omid, B. Davvaz and C. Abdioglu, *Some properties of quasi- Γ -hyperideals and hyperfilters in ordered Γ -semihypergroups*, Southeast Asian Bull. Math., 42(2) (2018), 223–242.
- [22] F. Smarandache, *Neutrosophic quadruple numbers, refined neutrosophic quadruple numbers, absorbance law, and the multiplication of neutrosophic quadruple numbers*, Neutrosophic Sets and Systems 10 (2015), 96-98.
- [23] F. Smarandache, *Neutrosophic set- A generalization of the intuitionistic fuzzy set*. Int. J. Pure Appl. Math. 24 (2005), 287-297.
- [24] F. Smarandache, *Neutrosophy. Neutrosophic Probability, Set, and Logic*, Amer. Res. Press, Rehoboth, USA, 105 p., 1998.
- [25] F. Smarandache, *Symbolic Neutrosophic Theory*, Chapter 7, Europa Nova, Bruxelles, 194 p., 2015; <https://arxiv.org/ftp/arxiv/papers/1512/1512.00047.pdf>
- [26] W. B. Vasantha, I. Kandasamy and F. Smarandache, *Algebraic Structure of Neutrosophic Duplets in Neutrosophic Rings $\langle Z \cup I \rangle$, $\langle Q \cup I \rangle$, and $\langle R \cup I \rangle$* , Neutrosophic Sets and Systems, 23(2018), 85-95.
- [27] T. Vougiouklis, *Cyclicity in a special class of hypergroups*, Acta Univ. Carolin. Math. Phys. 22(1) (1981), 3-6.

- [28] T. Vougiouklis, *The fundamental relation in hyperrings. The general hyperfield*, Algebraic hyperstructures and applications (Xnithi, 1990), 203-211, World Sci. Publ., Teaneck, NJ, 1991.
- [29] T. Vougiouklis, *Hyperstructures and their representations*, Hadronic Press Monographs in Mathematics, Hadronic Press, Inc., Palm Harbor, FL, 1994.
- [30] T. Vougiouklis, *From H_v -rings to H_v -fields*, Int. J. Algebraic Hyperstructures Appl. 1(1)(2014), 1-13.
- [31] T. Vougiouklis, *On the hyperstructure theory*, Southeast Asian Bull. Math., 40(4)(2016), 603–620.
- [32] L. A. Zadeh, *Fuzzy sets*, Inform and control, 8(1965), 338-353.

Received: April 26, 2019.

Accepted: June 04, 2019.