



Special equiform Smarandache curves in Minkowski space-time



E.M. Solouma^{a,b,*}

^aAl Imam Mohammad Ibn Saud Islamic University, College of Science, Department of Mathematics and Statistics, KSA, P. O. Box 90950, Riyadh 11623, Saudi Arabia

^bDepartment of Mathematics, Faculty of Science, Beni-Suef University, Egypt

ARTICLE INFO

Article history:

Received 10 March 2017

Revised 5 April 2017

Accepted 30 April 2017

Available online 23 May 2017

MSC:

53B30

53C40

53C50

Keywords:

Smarandache curve

Equiform Frenet frame

Minkowski space-time

ABSTRACT

In this paper, we introduce special equiform Smarandache curves reference to the equiform Frenet frame of a curve ζ on a spacelike surface M in Minkowski 3-space E_1^3 . Also, we study the equiform Frenet invariants of the spacial equiform Smarandache curves in E_1^3 . Moreover, we give some properties to these curves when the curve ζ has constant curvature or it is a circular helix. Finally, we give an example to illustrate these curves.

© 2017 Egyptian Mathematical Society. Production and hosting by Elsevier B.V.
This is an open access article under the CC BY-NC-ND license.
(<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

1. Introduction

A regular non-null curve in Minkowski space-time, whose position vector is composed by Frenet frame vectors on another regular curve, is called a Smarandache curve [1]. Recently special Smarandache curves have been studied by some authors [2–5].

In this work, we study special equiform Smarandache curves with reference to the equiform Frenet frame of a curve ζ on a spacelike surface M in Minkowski 3-space E_1^3 . In Section 2, we clarify the basic conceptions of Minkowski 3-space E_1^3 and give of equiform Frenet frame that will be used during this work. Section 3 is delicate to the study of the special four equiform Smarandache curves, $T\eta$, $T\xi$, $\eta\xi$ and $T\eta\xi$ -equiform Smarandache curves by being the connection with the first and second equiform curvature $k_1(\theta)$, and $k_2(\theta)$ of the equiform spacelike curve ζ in E_1^3 . Furthermore, we present some properties on the curves when the curve ζ has constant curvature or it is a circular helix. Finally, we give an example to clarify these curves. We hope these results will be helpful to mathematicians who are specialized on mathematical modeling.

2. Preliminaries

The Minkowski 3-space E_1^3 is the Euclidean 3-space E^3 provided with the metric

$$\mathcal{G} = -dz_1^2 + dz_2^2 + dz_3^2,$$

where (z_1, z_2, z_3) is a rectangular coordinate system of E_1^3 . Any arbitrary vector $v \in E_1^3$ can have one of three Lorentzian clause depicts; it can be timelike if $\mathcal{G}(v, v) < 0$, spacelike if $\mathcal{G}(v, v) > 0$ or $v = 0$, and lightlike if $\mathcal{G}(v, v) = 0$ and $v \neq 0$. Similarly, any arbitrary curve $\zeta = \zeta(s)$ can be timelike, spacelike or lightlike if all of its velocity vectors $\zeta'(s)$ are timelike, spacelike or lightlike, respectively.

Let $\zeta = \zeta(s)$ be a regular non-null curve parametrized by arc-length in E_1^3 and $\{t, n, b, \kappa, \tau\}$ be its Frenet invariants where $\{t, n, b\}$, κ and τ are the moving Frenet frame and the natural curvature functions respectively. If ζ is a spacelike curve with spacelike principal normal vector, then the Frenet formulas of the curve ζ can be given as [6–8]:

$$\begin{pmatrix} \dot{t}(s) \\ \dot{n}(s) \\ \dot{b}(s) \end{pmatrix} = \begin{pmatrix} 0 & \kappa(s) & 0 \\ -\kappa(s) & 0 & \tau(s) \\ 0 & \tau(s) & 0 \end{pmatrix} \begin{pmatrix} t(s) \\ n(s) \\ b(s) \end{pmatrix}, \quad (1)$$

where $\left(\cdot = \frac{d}{ds}\right)$, $\mathcal{G}(t, t) = \mathcal{G}(n, n) = -\mathcal{G}(b, b) = 1$, and $\mathcal{G}(t, n) = \mathcal{G}(t, b) = \mathcal{G}(n, b) = 0$.

* Corresponding author at: Al Imam Mohammad Ibn Saud Islamic University, College of Science, Department of Mathematics and Statistics, KSA, P. O. Box 90950, Riyadh 11623, Saudi Arabia.

E-mail address: emadms74@gmail.com

Definition 2.1. A surface M in the Minkowski 3-space E_1^3 is said to be timelike, spacelike surface if, respectively the induced metric on the surface is a Lorentz metrica, positive definite Riemannian metric. In other words, the normal vector on the timelike(spacelike) surface is a spacelike(timelike) vector [8].

Let $\zeta : I \rightarrow E_1^3$ be a spacelike curve in Minkowski space E_1^3 . We define the equiform parameter of ζ by $\theta = \int \kappa ds$. Then, we have $\rho = \frac{ds}{d\theta}$, where $\rho = \frac{1}{\kappa}$ is the radius of curvature of the curve ζ . Let \mathcal{F} be a homothety with the center in the origin and the coefficient μ . If we put $\bar{\zeta} = \mathcal{F}(\zeta)$, then it follows

$$\bar{s} = \mu s \text{ and } \bar{\rho} = \mu \rho,$$

where \bar{s} is the arc-length parameter of $\bar{\zeta}$ and $\bar{\rho}$ the radius of curvature of this curve. Therefore, θ is an equiform invariant parameter of ζ [9]. From that point, we recall $\{T, \eta, \xi, \}$ be the moving equiform Frenet frame where $T(\theta) = \rho t(s)$, $\eta(\theta) = \rho n(s)$ and $\xi(\theta) = \rho b(s)$ are the equiform tangent vector, equiform principal normal vector and equiform binormal vector respectively. Additionally, the first and second equiform curvature of the curve $\zeta = \zeta(\theta)$ are defined by $k_1(\theta) = \dot{\rho} = \frac{d\rho}{ds}$ and $k_2(\theta) = \frac{\tau}{\kappa}$. So, the moving equiform Frenet frame of $\zeta = \zeta(\theta)$ is given as [10]:

$$\begin{pmatrix} T'(\theta) \\ \eta'(\theta) \\ \xi'(\theta) \end{pmatrix} = \begin{pmatrix} k_1(\theta) & 1 & 0 \\ -1 & k_1(\theta) & k_2(\theta) \\ 0 & k_2(\theta) & k_1(\theta) \end{pmatrix} \begin{pmatrix} T(\theta) \\ \eta(\theta) \\ \xi(\theta) \end{pmatrix}, \tag{2}$$

where $\left(' = \frac{d}{d\theta} \right)$, $\mathcal{G}(T, T) = \mathcal{G}(\eta, \eta) = -\mathcal{G}(\xi, \xi) = \rho^2$, and $\mathcal{G}(T, \eta) = \mathcal{G}(T, \xi) = \mathcal{G}(\eta, \xi) = 0$.

The pseudo-Riemannian sphere with center at the origin and of radius $r = 1$ in the Minkowski 3-space E_1^3 is a quadric defined by

$$S_1^2 = \{ \bar{u} \in E_1^3 : -u_1^2 + u_2^2 + u_3^2 = 1. \}$$

Let $\zeta = \zeta(\theta)$ be a regular non-null curve parametrized by arc-length in Minkowski 3-space E_1^3 with its moving equiform Frenet frame $\{T, \eta, \xi, \}$. Then $T\eta$, $T\xi$, $\eta\xi$ and $T\eta\xi$ -equiform Smarandache curves of ζ are defined, respectively as follows [11]:

$$\mathfrak{S} = \mathfrak{S}(\theta^*) = \frac{1}{\sqrt{2}}(T(\theta) + \eta(\theta)),$$

$$\mathfrak{S} = \mathfrak{S}(\theta^*) = \frac{1}{\sqrt{2}}(T(\theta) + \xi(\theta)),$$

$$\mathfrak{S} = \mathfrak{S}(\theta^*) = \frac{1}{\sqrt{2}}(\eta(\theta) + \xi(\theta)),$$

$$\mathfrak{S} = \mathfrak{S}(\theta^*) = \frac{1}{\sqrt{3}}(T(\theta) + \eta(\theta) + \xi(\theta)).$$

3. Special equiform Smarandache curves in E_1^3

In this section, we define the special equiform Smarandache curves reference to the equiform Frenet frame of a curve ζ in Minkowski 3-space E_1^3 . Furthermore, we obtain the natural equiform curvature functions of the equiform Smarandache curves lying completely on pseudo-sphere S_1^2 and give some properties on the curves when the curve ζ has constant curvature or it is a circular helix

Definition 3.1. A curve in Minkowski space-time, whose position vector is composed by Frenet frame vectors on another curve, is called a Smarandache curve.

As consequence with the above definition, we introduce a special form of the equiform Smarandache curves in E_1^3 in the following subsection

3.1. $T\eta$ -equiform Smarandache curves in E_1^3

Definition 3.2. Let $\zeta = \zeta(\theta)$ be a regular equiform spacelike curve lying completely on a spacelike surface M in E_1^3 with moving equiform Frenet frame $\{T, \eta, \xi, \}$. Then $T\eta$ -equiform Smarandache curves are defined by

$$\mathfrak{S} = \mathfrak{S}(\theta^*) = \frac{1}{\sqrt{2}}(T(\theta) + \eta(\theta)). \tag{3}$$

Theorem 3.1. Let $\zeta = \zeta(s)$ be a spacelike curve with spacelike principal normal vector in E_1^3 . If ζ is a circular helix with $\kappa > 0$, then $T\eta$ -equiform Smarandache curve is also circular helix and its the natural curvature functions are satisfied the following equation,

$$\begin{aligned} \kappa_{\mathfrak{S}}(\theta^*) &= \frac{\sqrt{2}}{\rho(k_2^2 - 2)} : k_2 \neq \pm\sqrt{2}, \\ \tau_{\mathfrak{S}}(\theta^*) &= \frac{\sqrt{2} k_2(2k_2 + 1) - (k_2^2 - 1)[k_2 + k_2^2(k_2 + 2)]}{\rho^2(2(k_2^2 + 2))}, \\ k_2 &\neq -\sqrt{2}. \end{aligned} \tag{4}$$

Proof. Let $\mathfrak{S} = \mathfrak{S}(\theta^*)$ be a $T\eta$ -equiform Smarandache curves reference to the equiform spacelike curve $\zeta = \zeta(\theta)$. From Eq. (3) and using Eq. (2), we get

$$\mathfrak{S}'(\theta^*) = \frac{d\mathfrak{S}}{d\theta^*} \frac{d\theta^*}{d\theta} = \frac{1}{\sqrt{2}}((k_1 - 1)T(\theta) + (k_1 + 1)\eta(\theta) + k_2\xi(\theta)), \tag{5}$$

hence

$$T_{\mathfrak{S}}(\theta^*) = \frac{1}{\rho\sqrt{2k_1^2 - k_2^2 - 2}}((k_1 - 1)T(\theta) + (k_1 + 1)\eta(\theta) + k_2\xi(\theta)), \tag{6}$$

where

$$\frac{d\theta^*}{d\theta} = \frac{\rho\sqrt{2k_1^2 - k_2^2 - 2}}{\sqrt{2}}. \tag{7}$$

Now

$$\frac{dT_{\mathfrak{S}}}{d\theta^*} = \frac{\sqrt{2}}{\rho^2[2k_1^2 - k_2^2 - 2]^2}(\lambda_1 T(\theta) + \lambda_2 \eta(\theta) + \lambda_3 \xi(\theta)),$$

where

$$\begin{cases} \lambda_1 = (k_1 - 1)(2k_1k_1' - k_2k_2') + (2k_1^2 - k_2^2 - 2)(k_1' + k_1^2 - 3k_1), \\ \lambda_2 = (k_1 + 1)(2k_1k_1' - k_2k_2') + (2k_1^2 - k_2^2 - 2)(k_1' + k_1^2 + k_2^2 + k_1 - 2), \\ \lambda_3 = k_2(2k_1k_1' - k_2k_2') + (2k_1^2 - k_2^2 - 2)(k_2' + 2k_1k_2). \end{cases}$$

Then

$$\kappa_{\mathfrak{S}}(\theta^*) = \left\| \frac{dT_{\mathfrak{S}}}{d\theta^*} \right\| = \frac{\sqrt{2(\lambda_1^2 + \lambda_2^2 - \lambda_3^2)}}{\rho[2k_1^2 - k_2^2 - 2]^2}, \tag{8}$$

and

$$N_{\mathfrak{S}}(\theta^*) = \frac{\lambda_1 T(\theta) + \lambda_2 \eta(\theta) + \lambda_3 \xi(\theta)}{\rho\sqrt{\lambda_1^2 + \lambda_2^2 - \lambda_3^2}}.$$

Also

$$B_{\mathfrak{S}}(\theta^*) = \frac{1}{p_1}\{m_1 T(\theta) + m_2 \eta(\theta) + m_3 \xi(\theta)\},$$

where

$$m_1 = \lambda_2 k_2 - \lambda_3 (k_1 + 1),$$

$$m_2 = \lambda_2 k_2 - \lambda_3 (k_1 - 1),$$

$$m_3 = \lambda_2 (k_1 - 1) - \lambda_1 (k_1 + 1)$$

$$\text{and } p_1 = \rho \sqrt{2k_1^2 - k_2^2 - 2\sqrt{\lambda_1^2 + \lambda_2^2 - \lambda_3^2}}.$$

Now, from Eq. (5)

$$\mathfrak{S}''(\theta^*) = \frac{1}{\sqrt{2}} \left\{ [k'_1 + k_1^2 - 2k_1 - 1]T(\theta) + [k'_1 + k_1^2 + k_2^2 + 2k_1 - 1]\eta(\theta) + [k'_2 + 2k_1 k_2 + k_2]\xi(\theta) \right\},$$

and thus

$$\mathfrak{S}'''(\theta^*) = \frac{1}{\sqrt{2}} (\beta_1 T(\theta) + \beta_2 \eta(\theta) + \beta_3 \xi(\theta)),$$

where

$$\begin{cases} \beta_1 = k'_1 + 3k'_1(k_1 - 1) + k_1^2(k_1 - 3) - k_2(k_2 + 2), \\ \beta_2 = k'_1 + k_1 + 3(k_1 k'_1 + k_2 k'_2) + 3k_1(k_1 - 1) + k_1^2(3k_1 + 1) - 1, \\ \beta_3 = k'_2 + k_2 + 3(k_1 k'_1 + k_2 k'_2) + 3k_1 k_2(k_1 + 1) - k_2. \end{cases}$$

Hence, we have

$$\tau_{\mathfrak{S}}(\theta^*) = \frac{\sqrt{2}}{\rho^2} \left\{ \frac{w_1 + w_2 + w_3}{\ell_1^2 + \ell_2^2 - \ell_3^2} \right\}, \tag{9}$$

where

$$w_1 = (k'_1 + k_1^2 + k_2^2 + 2k_1 - 1)[\beta_3(k_1 - 1) - \beta_1 k_2],$$

$$w_2 = (k'_2 + 2k_1 k_2 + k_2)[\beta_1(k_1 + 1) - \beta_2(k_1 - 1)],$$

$$w_3 = (k'_1 + k_1^2 - 2k_1 - 1)[\beta_2 k_2 - \beta_3(k_1 + 1)],$$

$$\ell_1 = k'_1 k_2 - k'_2(k_1 + 1) + k_2(k_2 - k_1^2 - k_1 - 2),$$

$$\ell_2 = k'_1 k_2 - k'_2(k_1 - 1) - k_1 k_2(k_1 + 1),$$

$$\ell_3 = k_1(2k_1 + 1) - 2k'_1 + k_2^2(k_1 + 1) + 2.$$

Now, if κ and τ are non-zero constants, then the natural curvature functions $\kappa_{\mathfrak{S}}$, $\tau_{\mathfrak{S}}$ are also non-zero constants and satisfying Eq. (4) which means that the $T\eta$ -equiform Smarandache curve is circular helix. \square

3.2. $T\xi$ -equiform Smarandache curves in E_1^3

Definition 3.3. Let $\zeta = \zeta(\theta)$ be a regular equiform spacelike curve lying completely on a spacelike surface M in E_1^3 with moving equiform Frenet frame $\{T, \eta, \xi\}$. Then $T\xi$ -equiform Smarandache curves are defined by

$$\mathfrak{S} = \mathfrak{S}(\theta^*) = \frac{1}{\sqrt{2}} (T(\theta) + \xi(\theta)). \tag{10}$$

Theorem 3.2. Let $\zeta = \zeta(s)$ be a spacelike curve with spacelike principal normal vector in E_1^3 . If ζ is a circular helix with $\kappa > 0$, then $T\xi$ -equiform Smarandache curve is contained in a plane and its curvature is satisfied the following equation,

$$\kappa_{\mathfrak{S}}(\theta^*) = \frac{\sqrt{2}\sqrt{(1 - k_2^2)(k_2 + 1)^2 - k_2^2(3k_2^2 - 2)}}{\rho(k_2 + 1)^2} : k_2 \neq -1. \tag{11}$$

Proof. Let $\mathfrak{S} = \mathfrak{S}(\theta^*)$ be a $T\xi$ -equiform Smarandache curves of $\zeta = \zeta(\theta)$. Then from Eq. (10), we have

$$\mathfrak{S}'(\theta^*) = \frac{1}{\sqrt{2}} (k_1 T(\theta) + (k_2 + 1)\eta(\theta) + k_1 \xi(\theta)). \tag{12}$$

$$T_{\mathfrak{S}}(\theta^*) = \frac{1}{\rho(k_2 + 1)} (k_1 T(\theta) + (k_2 + 1)\eta(\theta) + k_1 \xi(\theta)), \tag{13}$$

where

$$\frac{d\theta^*}{d\theta} = \frac{\rho(k_2 + 1)}{\sqrt{2}}. \tag{14}$$

Now

$$\frac{dT_{\mathfrak{S}}}{d\theta^*} = \frac{\sqrt{2}}{\rho^2(k_2 + 1)^3} (\varepsilon_1 T(\theta) + \varepsilon_2 \eta(\theta) + \varepsilon_3 \xi(\theta)),$$

where

$$\begin{cases} \varepsilon_1 = (k_2 + 1)(k'_1 - k_2 - 1) - k_1 k_2, \\ \varepsilon_2 = k_1(k_2 + 1)^2, \\ \varepsilon_3 = (k_2 + 1)[k'_1 + k_2(k_2 + 1) + k_2^2 - k_1^2] - k_1 k'_2. \end{cases}$$

Then

$$\kappa_{\mathfrak{S}}(\theta^*) = \frac{\sqrt{2}\sqrt{\varepsilon_1^2 + \varepsilon_2^2 - \varepsilon_3^2}}{\rho(k_2 + 1)^3}, \tag{15}$$

and

$$N_{\mathfrak{S}}(\theta^*) = \frac{\varepsilon_1 T(\theta) + \varepsilon_2 \eta(\theta) + \varepsilon_3 \xi(\theta)}{\rho\sqrt{\varepsilon_1^2 + \varepsilon_2^2 - \varepsilon_3^2}}.$$

Also

$$B_{\mathfrak{S}}(\theta^*) = \frac{1}{p_2} \left\{ [\varepsilon_2 k_1 - \varepsilon_3(k_2 + 1)]T(\theta) + k_1(\varepsilon_1 - \varepsilon_3)\eta(\theta) + [\varepsilon_2 k_1 - \varepsilon_1(k_2 + 1)]\xi(\theta) \right\},$$

where $p_2 = \rho(k_2 + 1)\sqrt{\varepsilon_1^2 + \varepsilon_2^2 - \varepsilon_3^2}$.

Now, from Eq. (12) we have

$$\mathfrak{S}''(\theta^*) = \frac{1}{\sqrt{2}} \left\{ [k'_1 + k_1^2 - k_2 + 1]T(\theta) + [k'_2 + 2k_1(k_2 + 2)]\eta(\theta) + [k'_1 + k_2(2k_2 + 1)]\xi(\theta) \right\},$$

and

$$\mathfrak{S}'''(\theta^*) = \frac{1}{\sqrt{2}} (\delta_1 T(\theta) + \delta_2 \eta(\theta) + \delta_3 \xi(\theta)),$$

where

$$\begin{cases} \delta_1 = k'_1 - k'_2 + 3k_1(k'_1 - k_2) + k_1(k_1^2 - 1), \\ \delta_2 = k'_2 + k'_1 + 3(k'_1 k_2 + k_1 k'_2) + k_1^2(2k_2 + 3) + k_2(2k_2^2 + k_2 - 1) + 1, \\ \delta_3 = k'_1 + k'_2 + k_1(k'_1 + 3k_2) + k_2(5k'_2 + 4k_1 k_2). \end{cases}$$

Hence, we have

$$\tau_{\mathfrak{S}}(\theta^*) = \frac{\sqrt{2}}{\rho^2} \left\{ \frac{(k'_1 + 2k_2^2 + k_2)[\delta_1(k_2 + 1) - \delta_2 k_1] + k_1(\delta_3 - \delta_1)}{(k'_2 + 2k_1 k_2 + 2k_1) + (k'_1 + k_1^2 - k_2 + 1)[\delta_2 k_1 - \delta_3(k_2 + 1)]}{\left\{ k_1 k'_2 + (k_2 + 1)(2k_1^2 - k_1^2) - k_2(2k_2^2 + 3k_2 + 1) \right\}^2 + \left\{ k_1[k_1^2 - 2k_2(k_2 + 1) + 1] \right\}^2 - \left\{ k_1 k'_2 + k_2^2 - k_1^2 - k'_1(k_2 + 1) + k_2^2(k_2 + 1) - 1 \right\}^2} \right\}. \tag{16}$$

So, if κ and τ are non-zero constants, then $\kappa_{\mathfrak{S}}$ is non-zero constant and satisfying Eq. (11), also $\tau_{\mathfrak{S}} = 0$ which means that the $T\xi$ -equiform Smarandache curve is contained in a plane. \square

3.3. $\eta\xi$ -equiform Smarandache curves in E_1^3

Definition 3.4. Let $\zeta = \zeta(\theta)$ be a regular equiform spacelike curve lying completely on a spacelike surface M in E_1^3 with moving equiform Frenet frame $\{T, \eta, \xi\}$. Then $\eta\xi$ -equiform Smarandache curves are defined by

$$\mathfrak{S} = \mathfrak{S}(\theta^*) = \frac{1}{\sqrt{2}} (\eta(\theta) + \xi(\theta)). \tag{17}$$

Theorem 3.3. Let $\zeta = \zeta(s)$ be a spacelike curve with spacelike principal normal vector in E_1^3 . If ζ is a circular helix with $\kappa > 0$, then

$\eta\xi$ -equiform Smarandache curve is also circular helix and its the natural curvature functions are satisfied the following equation,

$$\begin{aligned} \kappa_{\mathfrak{S}}(\theta^*) &= \frac{\sqrt{2}(k_2^2 - 1)}{\rho}, \\ \tau_{\mathfrak{S}}(\theta^*) &= \frac{\sqrt{2}}{\rho^2 k_2} : k_2 \neq 0. \end{aligned} \tag{18}$$

Proof. Let $\mathfrak{S} = \mathfrak{S}(\theta^*)$ be a $\eta\xi$ -equiform Smarandache curves of the curve $\zeta = \zeta(\theta)$. From Eq. (17), we get

$$\mathfrak{S}'(\theta^*) = \frac{1}{\sqrt{2}}(-T(\theta) + (k_1 + k_2)\eta(\theta) + (k_1 + k_2)\xi(\theta)), \tag{19}$$

hence

$$T_{\mathfrak{S}}(\theta^*) = \frac{1}{\rho}(-T(\theta) + (k_1 + k_2)\eta(\theta) + (k_1 + k_2)\xi(\theta)), \tag{20}$$

where

$$\frac{d\theta^*}{d\theta} = \frac{\rho}{\sqrt{2}}. \tag{21}$$

Now

$$\frac{dT_{\mathfrak{S}}}{d\theta^*} = \frac{\sqrt{2}}{\rho^2}(\gamma_1 T(\theta) + \gamma_2 \eta(\theta) + \gamma_3 \xi(\theta)),$$

where

$$\begin{aligned} \gamma_1 &= -(k_1 + k_2), \\ \gamma_2 &= k'_1 + k'_2 + k_2(k_1 + k_2) - 1, \\ \gamma_3 &= k'_1 + k'_2 + k_2(k_1 + k_2). \end{aligned}$$

Then

$$\kappa_{\mathfrak{S}}(\theta^*) = \frac{\sqrt{2(\gamma_1^2 + \gamma_2^2 - \gamma_3^2)}}{\rho}, \tag{22}$$

and

$$N_{\mathfrak{S}}(\theta^*) = \frac{\gamma_1 T(\theta) + \gamma_2 \eta(\theta) + \gamma_3 \xi(\theta)}{\rho \sqrt{\gamma_1^2 + \gamma_2^2 - \gamma_3^2}}.$$

Also

$$B_{\mathfrak{S}}(\theta^*) = \frac{1}{\rho \sqrt{\gamma_1^2 + \gamma_2^2 - \gamma_3^2}} \{[(\gamma_2 - \gamma_3)(k_1 + k_2)]T(\theta) + [\gamma_3 + \gamma_1(k_1 + k_2)]\eta(\theta) - [\gamma_2 + \gamma_1(k_1 + k_2)]\xi(\theta)\}.$$

From Eq. (19), we have

$$\mathfrak{S}''(\theta^*) = \frac{1}{\sqrt{2}}\{-[2k_1 + k_2]T(\theta) + [k'_1 + k'_2 + (k_1 + k_2)^2 - 1]\eta(\theta) + [k'_1 + k'_2 + (k_1 + k_2)^2]\xi(\theta)\},$$

and

$$\mathfrak{S}'''(\theta^*) = \frac{1}{\sqrt{2}}(\omega_1 T(\theta) + \omega_2 \eta(\theta) + \omega_3 \xi(\theta)),$$

where

$$\begin{cases} \omega_1 = -[3k'_1 + 2k'_2 + k_1(2k_1 + k_2) + (k_1 + k_2)^2 - 1], \\ \omega_2 = k''_1 + k''_2 - 3k_1 - k_2 + 3(k_1 + k_2)(k'_1 + k'_2) + (k_1 + k_2)^3, \\ \omega_3 = k''_1 + k''_2 + 3(k_1 + k_2)(k'_1 + k'_2) + (k_1 + k_2)^3. \end{cases}$$

Hence, we have

$$\tau_{\mathfrak{S}}(\theta^*) = \frac{\sqrt{2} \left\{ \begin{aligned} &[k'_1 + k'_2 + (k_1 + k_2)^2 - 1][\omega_3 + \omega_1(k_1 + k_2)] \\ &- [k'_1 + k'_2 + (k_1 + k_2)^2][\omega_2 + \omega_1(k_1 + k_2)] \\ &- (\omega_3 - \omega_2)(k_1 + k_2)(2k_1 + k_2) \end{aligned} \right\}}{\rho^2 \{k_1^2 - k_2^2 + 2(k'_1 + k'_2)\}}. \tag{23}$$

Then, if κ and τ are non-zero constants, then the natural curvature functions $\kappa_{\mathfrak{S}}$, $\tau_{\mathfrak{S}}$ are also non-zero constants and satisfying Eq. (18) which means that the $\eta\xi$ -equiform Smarandache curve is circular helix. \square

3.4. $T\eta\xi$ -equiform Smarandache curves in E_1^3

Definition 3.5. Let $\zeta = \zeta(\theta)$ be a regular equiform spacelike curve lying completely on a spacelike surface M in E_1^3 with moving equiform Frenet frame $\{T, \eta, \xi\}$. Then $T\eta\xi$ -equiform Smarandache curves are defined by

$$\mathfrak{S} = \mathfrak{S}(\theta^*) = \frac{1}{\sqrt{3}}(T(\theta) + \eta(\theta) + \xi(\theta)). \tag{24}$$

Theorem 3.4. Let $\zeta = \zeta(s)$ be a spacelike curve with spacelike principal normal vector in E_1^3 . If ζ is a circular helix with $\kappa > 0$, then $T\eta\xi$ -equiform Smarandache curve is also circular helix and its the natural curvature functions are satisfied the following equation,

$$\begin{aligned} \kappa_{\mathfrak{S}}(\theta^*) &= \frac{\sqrt{3}\sqrt{2(1-k_2)}}{2\rho} : |k_1| < 1, \\ \tau_{\mathfrak{S}}(\theta^*) &= \frac{\sqrt{3}}{3\rho^2} \frac{k_2(k_2^2 + 1)}{(k_2 - 3)(k_2 + 1)^2} : k_2 \neq -1, 3. \end{aligned} \tag{25}$$

Proof. Let $\mathfrak{S} = \mathfrak{S}(\theta^*)$ be a $T\eta\xi$ -equiform Smarandache curves of the curve $\zeta = \zeta(\theta)$. Then from Eq. (24), we get

$$\mathfrak{S}'(\theta^*) = \frac{1}{\sqrt{3}}((k_1 - 1)T(\theta) + (k_1 + k_2 + 1)\eta(\theta) + (k_1 + k_2)\xi(\theta)). \tag{26}$$

$$T_{\mathfrak{S}}(\theta^*) = \frac{1}{\rho \sqrt{k_1^2 + 2k_2 + 2}} ((k_1 - 1)T(\theta) + (k_1 + k_2 + 1)\eta(\theta) + (k_1 + k_2)\xi(\theta)), \tag{27}$$

where

$$\frac{d\theta^*}{d\theta} = \frac{\rho \sqrt{k_1^2 + 2k_2 + 2}}{\sqrt{3}}. \tag{28}$$

Now

$$\frac{dT_{\mathfrak{S}}}{d\theta^*} = \frac{\sqrt{3}}{\rho^2 [k_1^2 + 2k_2 + 2]^2} (\chi_1 T(\theta) + \chi_2 \eta(\theta) + \chi_3 \xi(\theta)),$$

where

$$\begin{cases} \chi_1 = (k_1 - 1)(k_1 k'_1 + k'_2) - (k_2 + 1)(k_1^2 + 2k_2 + 2), \\ \chi_2 = (k_1^2 + 2k_2 + 2)[k'_1 + k'_2 + k_1 + k_2(k_1 + k_2) - 1] \\ \quad + (k_1 k'_1 + k'_2)(k_1 + k_2 + 1), \\ \chi_3 = (k_1^2 + 2k_2 + 2)[k'_1 + k'_2 + k_2(k_1 + k_2 + 1)] \\ \quad + (k_1 + k_2)(k_1 k'_1 + k'_2). \end{cases}$$

Then

$$\kappa_{\mathfrak{S}}(\theta^*) = \frac{\sqrt{3(\chi_1^2 + \chi_2^2 - \chi_3^2)}}{\rho [k_1^2 + 2k_2 + 2]^2}, \tag{29}$$

and

$$N_{\mathfrak{S}}(\theta^*) = \frac{\chi_1 T(\theta) + \chi_2 \eta(\theta) + \chi_3 \xi(\theta)}{\rho \sqrt{\chi_1^2 + \chi_2^2 - \chi_3^2}}.$$

Also

$$\begin{aligned} B_{\mathfrak{S}}(\theta^*) &= \frac{1}{\rho \sqrt{k_1^2 + 2k_2 + 2} \sqrt{\chi_1^2 + \chi_2^2 - \chi_3^2}} \\ &\times \left\{ [-\chi_3 + (\chi_2 - \chi_3)(k_1 + k_2)]T(\theta) \right. \\ &+ [\chi_1(k_1 + k_2) - \chi_1(k_1 - 1)]\eta(\theta) + [\chi_2(k_2 - 1) \\ &\left. - \chi_1(k_1 + k_2 + 1)]\xi(\theta) \right\}, \end{aligned}$$




Fig. 1. Spacelike curve $\zeta = \zeta(s)$ on S_1^2 .

From Eq. (26), we have

$$\begin{aligned} \mathfrak{S}''(\theta^*) &= \frac{1}{\sqrt{3}} \{ [k_1^2 - k_1 - k_2 - 1]T(\theta) \\ &\quad + [k_1' + k_2' + 2k_1 + (k_1 + k_2)^2 - 1]\eta(\theta) \\ &\quad + [k_1' + k_2' + k_2 + (k_1 + k_2)^2]\xi(\theta) \}, \end{aligned}$$

and thus

$$\mathfrak{S}'''(\theta^*) = \frac{1}{\sqrt{3}} (\phi_1 T(\theta) + \phi_2 \eta(\theta) + \phi_3 \xi(\theta)),$$

where

$$\begin{cases} \phi_1 = 2k_1'(k_1 + 1) - 2k_2' + k_1(k_1^2 - k_1 - k_2 - 2) + 1, \\ \phi_2 = k_1' + k_2' + 2k_1' + k_2(k_2 - 1) + 3(k_1 + k_2)(k_1' + k_2') \\ \quad + k_1(2k_1^2 + k_1 - 1) + (k_1 + k_2)^3 - 1, \\ \phi_3 = k_1' + k_2' + k_2' + k_2(3k_1 - 1) + 3(k_1 + k_2)(k_1' + k_2') \\ \quad + (k_1 + k_2)^3. \end{cases}$$

Hence, we have

$$\tau_{\mathfrak{S}}(\theta^*) = \frac{\sqrt{3}}{\rho^2} \left\{ \frac{v_1 + v_2 + v_3}{q^2 + q_2^2 - q_3^2} \right\}, \tag{30}$$

where

$$\begin{aligned} v_1 &= (k_1^2 - k_1 - k_2 - 1)[\phi_2(k_1 + k_2) - \phi_3(k_1 + k_2 - 1)], \\ v_2 &= [k_1' + k_2' + 2k_1 + (k_1 + k_2)^2 - 1][\phi_3(k_1 - 1) - \phi_1(k_1 + k_2)], \\ v_3 &= [k_1' + k_2' + k_2 + (k_1 + k_2)^2][\phi_1(k_1 + k_2 - 1) - \phi_2(k_1 - 1)], \\ q_1 &= (k_1 + k_2)(2k_1 - k_2 - 1) - [k_1' + k_2' + 2k_1 + (k_1 + k_2)^2], \\ q_2 &= -(k_1 + k_2)(k_1 k_2 + 1) - (k_1 - 1)(k_1' + k_2' + k_2), \\ q_3 &= (k_1 - 1)[k_1' + k_2' + 2k_1 + (k_1 + k_2)^2] + k_1(2k_1 - 3) \\ &\quad + (k_1 + k_2 + 1)[2(k_2 + 1) - k_1(k_1 + 1)] + 1. \end{aligned}$$

Now, if κ and τ are non-zero constants, then the natural curvature functions $\kappa_{\mathfrak{S}}$, $\tau_{\mathfrak{S}}$ are also non-zero constants and satisfying Eq. (25) which means that the $T\eta$ -equiform Smarandache curve is circular heilx. \square




Fig. 2. Equiform spacelike curve $\zeta = \zeta(\theta)$ on S_1^2 .




Fig. 3. The $T\eta$ -equiform Smarandache curve $\mathfrak{S}(\theta^*)$ on S_1^2 .

4. Example

Let $\zeta(s) = (\sqrt{3}s, s \sin(\sqrt{3} \ln s), s \cos(\sqrt{3} \ln s))$ be a unit speed spacelike curve parametrized by arc-length s with spacelike principal normal vector in E_1^3 (see Fig. 1). Then it is easy to show that

$$\begin{cases} t(s) = (\sqrt{3}, \sin(\sqrt{3} \ln s) + \sqrt{3} \cos(\sqrt{3} \ln s), \\ \quad \cos(\sqrt{3} \ln s) - \sqrt{3} \sin(\sqrt{3} \ln s)), \\ n(s) = \frac{1}{\sqrt{2}} (0, \cos(\sqrt{3} \ln s) - \sqrt{3} \sin(\sqrt{3} \ln s), \\ \quad -\sin(\sqrt{3} \ln s) - \sqrt{3} \cos(\sqrt{3} \ln s)), \\ \kappa = \frac{2\sqrt{3}}{s}, \quad \rho = \frac{s}{2\sqrt{3}}, \quad k_1 = \frac{1}{2\sqrt{3}}, \\ b(s) = (2, \frac{\sqrt{3}}{2} \sin(\sqrt{3} \ln s) + \frac{3}{2} \cos(\sqrt{3} \ln s), \\ \quad \frac{\sqrt{3}}{2} \cos(\sqrt{3} \ln s) - \frac{3}{2} \sin(\sqrt{3} \ln s)), \\ \tau = \frac{3}{s}, \quad k_2 = \frac{\sqrt{3}}{2}. \end{cases}$$




Fig. 4. The $T\xi$ -equiiform Smarandache curve $\mathfrak{S}(\theta^*)$ on S^2_1 .

Hence, the equiform parameter is $\theta = \int \kappa ds = 2\sqrt{3}s + c$. Here we take $c = 0$, then we have $s = e^{\theta/2\sqrt{3}}$ and $\rho = \frac{e^{\theta/2\sqrt{3}}}{2\sqrt{3}}$. So the equiform spacelike curve ζ is define as (see Fig. 2)

$$\zeta(\theta) = \left(\sqrt{3} e^{\theta/2\sqrt{3}}, e^{\theta/2\sqrt{3}} \sin\left(\frac{\theta}{2}\right), e^{\theta/2\sqrt{3}} \cos\left(\frac{\theta}{2}\right) \right).$$

It easy to show that

$$T(\theta) = \frac{e^{\theta/2\sqrt{3}}}{2} \left(1, \frac{1}{\sqrt{3}} \sin\left(\frac{\theta}{2}\right) + \cos\left(\frac{\theta}{2}\right), \frac{1}{\sqrt{3}} \cos\left(\frac{\theta}{2}\right) - \sin\left(\frac{\theta}{2}\right) \right).$$

It is clear that T is an equiform spacelike vector. Also

$$\eta(\theta) = \frac{e^{\theta/2\sqrt{3}}}{4} \left(0, \frac{1}{\sqrt{3}} \cos\left(\frac{\theta}{2}\right) - \sin\left(\frac{\theta}{2}\right), \frac{-1}{\sqrt{3}} \sin\left(\frac{\theta}{2}\right) - \cos\left(\frac{\theta}{2}\right) \right),$$

and

$$\xi(\theta) = \frac{e^{\theta/2\sqrt{3}}}{4} \left(\frac{4}{\sqrt{3}}, \sin\left(\frac{\theta}{2}\right) + \sqrt{3} \cos\left(\frac{\theta}{2}\right), \cos\left(\frac{\theta}{2}\right) - \sqrt{3} \sin\left(\frac{\theta}{2}\right) \right).$$

Then η is an equiform spacelike vector and ξ is an equiform time-like vector.

The $T\eta$ -equiiform Smarandache curve $\mathfrak{S}(\theta^*)$ of the curve $\zeta(\theta)$ is given by (see Fig. 3)

$$\mathfrak{S}(\theta^*) = \frac{\sqrt{6} e^{\theta/2\sqrt{3}}}{24} \left(2\sqrt{3}, (2\sqrt{3} + 1) \cos\left(\frac{\theta}{2}\right) + (2 - \sqrt{3}) \sin\left(\frac{\theta}{2}\right), (2 - \sqrt{3}) \cos\left(\frac{\theta}{2}\right) - (2\sqrt{3} + 1) \sin\left(\frac{\theta}{2}\right) \right).$$

The $T\xi$ -equiiform Smarandache curve $\mathfrak{S}(\theta^*)$ of the curve $\zeta(\theta)$ is given by (see Fig. 4)

$$\mathfrak{S}(\theta^*) = \frac{\sqrt{6} e^{\theta/2\sqrt{3}}}{24} \left(2(2 + \sqrt{3}), (2 + \sqrt{3}) \sin\left(\frac{\theta}{2}\right) + (2\sqrt{3} + 3) \cos\left(\frac{\theta}{2}\right), (2 + \sqrt{3}) \cos\left(\frac{\theta}{2}\right) - (2\sqrt{3} + 3) \sin\left(\frac{\theta}{2}\right) \right).$$




Fig. 5. The $\eta\xi$ -equiiform Smarandache curve $\mathfrak{S}(\theta^*)$ on S^2_1 .

The $\eta\xi$ -equiiform Smarandache curve $\mathfrak{S}(\theta^*)$ of the curve $\zeta(\theta)$ is given by (see Fig. 5)

$$\mathfrak{S}(\theta^*) = \frac{\sqrt{6} e^{\theta/2\sqrt{3}}}{6} \left(1, \cos\left(\frac{\theta}{2}\right), -\sin\left(\frac{\theta}{2}\right) \right).$$

The $T\xi$ -equiiform Smarandache curve $\mathfrak{S}(\theta^*)$ of the curve $\zeta(\theta)$ is given by (see Fig. 6)




Fig. 6. The $T\eta\xi$ -equiform Smarandache curve $\mathfrak{S}(\theta^*)$ on S_1^2 .

$$\mathfrak{S}(\theta^*) = \frac{\sqrt{9}e^{\theta/2\sqrt{3}}}{18} \left(2 + \sqrt{3} \sin\left(\frac{\theta}{2}\right) + (2 + \sqrt{3}) \cos\left(\frac{\theta}{2}\right), \cos\left(\frac{\theta}{2}\right), -(2 + \sqrt{3}) \sin\left(\frac{\theta}{2}\right) \right).$$

Acknowledgment

The author is grateful to the Editor in Chief and the referees for their efforts and constructive criticism which have improved the manuscript

References

- [1] C. Ashbacher, Smarandache geometries, *Smarandache Notions J.* 8 (1-3) (1997) 212–215.
- [2] Ö. Bektaş, S. Yüce, Special Smarandache curves according to Darboux frame in euclidean 3- space, *Romanian J. Math. Comput. Sci.* 3 (2013) 48–59.
- [3] M. Çetin, Y. Tunçer, M.K. Karacan, Smarandache curves according to bishop frame in euclidean 3- space, *Gen. Math. Notes* 20 (2) (2014) 50–66.
- [4] E.M. Solouma, Special Smarandache curves recording by curves on a space-like surface in Minkowski space-time, *PONTE J.* 73 (2) (2017) 251–263, doi:10.21506/j.ponte.2017.2.20.
- [5] K. Taşköprü, M. Tosun, Smarandache curves according to Sabban frame on, *Boletim da Sociedade Paraneense de Matematica* 32 (1) (2014) 51–59.
- [6] M.P.D. Carmo, *Differential geometry of curves and surfaces*, Prentice Hall, Englewood Cliffs, NJ, 1976.
- [7] R. López, Differential geometry of curves and surfaces in Lorentz–Minkowski space, *Int. Electron. J. Geometry* 7 (7) (2014) 44–107.
- [8] B. O'Neill, *Semi-Riemannian geometry with applications to relativity*, Academic press, New York, 1983.
- [9] M.E. Aydın, Mahmut Ergut the equiform differential geometry of curves in 4-dimensional galilean space g_4 , *Stud. Univ. Babeş-Bolyai Math.* 58 (3) (2013) 399–406.
- [10] H.K. Elsayied, M. Elzawy, A. Elsharkawy, Equiform spacelike normal curves in Minkowski 3-space, in: *Proceedings of the 5th International Conference of Mathematics & Information Sciences*, Zewail City, Egypt, 2016, pp. 11–13.
- [11] M. Turgut, S. Yılmaz, Smarandache curves in Minkowski space-time, *Int. J. Math. Comb.* 3 (2008) 51–55.