





Hashem Bordbar* 
Rajab Ali Borzooei 
Florentin Smarandache 
Young Bae Jun 

A GENERAL MODEL OF NEUTROSOPHIC IDEALS IN BCK/BCI -ALGEBRAS BASED ON NEUTROSOPHIC POINTS

Abstract

More general form of $(\in, \in \vee q)$ -neutrosophic ideal is introduced, and their properties are investigated. Relations between (\in, \in) -neutrosophic ideal and $(\in, \in \vee q_{(k_T, k_I, k_F)})$ -neutrosophic ideal are discussed. Characterizations of $(\in, \in \vee q_{(k_T, k_I, k_F)})$ -neutrosophic ideal are discussed, and conditions for a neutrosophic set to be an $(\in, \in \vee q_{(k_T, k_I, k_F)})$ -neutrosophic ideal are displayed.

Keywords: Ideal, neutrosophic \in -subset, neutrosophic q_k -subset, neutrosophic $\in \vee q_k$ -subset, $(\in, \in \vee q_{(k_T, k_I, k_F)})$ -neutrosophic ideal.

2010 Mathematical Subject Classification: 06F35, 03G25, 03B52.

1. Introduction

Smarandache [23, 24] introduced the concept of neutrosophic sets which is a more general platform to extend the notions of the classical set and

*Corresponding author.

(intuitionistic, interval valued) fuzzy set. Neutrosophic set theory is applied to several parts which are referred to the site <http://fs.gallup.unm.edu/neutrosophy.htm>. Jun [10] introduced the notion of neutrosophic subalgebras in BCK/BCI -algebras based on neutrosophic points. Borumand and Jun [22] studied several properties of $(\in, \in \vee q)$ -neutrosophic subalgebras and $(q, \in \vee q)$ -neutrosophic subalgebras in BCK/BCI -algebras. Jun et al. [11] discussed neutrosophic \mathcal{N} -structures with an application in BCK/BCI -algebras, and in [13, 14] introduced neutrosophic quadruple numbers based on a set and construct neutrosophic quadruple BCK/BCI -algebras.

Song et al. [25] introduced the notion of commutative \mathcal{N} -ideal in BCK -algebras and investigated several properties. Bordbar, Jun and et al. [21] and [17] introduced the notion of $(q, \in \vee q)$ -neutrosophic ideal, and $(\in, \in \vee q)$ -neutrosophic ideal in BCK/BCI -algebras, and investigated related properties. Also in [7, 26], they discussed the notion of $BMBJ$ -neutrosophic sets, subalgebra and ideals, as a generalisation of neutrosophic set, and investigated it's application and related properties to BCI/BCK -algebras.

For more information about the mentioned topics, please refer to [3, 4, 8, 12, 16, 18, 19, 20].

In this paper, we introduce a more general form of $(\in, \in \vee q)$ -neutrosophic ideal, and investigate their properties. We discuss relations between (\in, \in) -neutrosophic ideal and $(\in, \in \vee q_{(k_T, k_I, k_F)})$ -neutrosophic ideal. We consider characterizations of $(\in, \in \vee q_{(k_T, k_I, k_F)})$ -neutrosophic ideal. We investigate conditions for a neutrosophic set to be an $(\in, \in \vee q_{(k_T, k_I, k_F)})$ -neutrosophic ideal. We find conditions for an $(\in, \in \vee q_{(k_T, k_I, k_F)})$ -neutrosophic ideal to be an (\in, \in) -neutrosophic ideal.

2. Preliminaries

By a BCI -algebra we mean a set X with a binary operation $*$ and the special element 0 satisfying the axioms:

$$(a1) \quad ((x * y) * (x * z)) * (z * y) = 0,$$

$$(a2) \quad (x * (x * y)) * y = 0,$$

$$(a3) \quad x * x = 0,$$

$$(a4) \quad x * y = y * x = 0 \Rightarrow x = y,$$

for all $x, y, z \in X$. If a BCI-algebra X satisfies the axiom

$$(a5) \quad 0 * x = 0 \text{ for all } x \in X,$$

then we say that X is a BCK-algebra. A subset I of a BCK/BCI-algebra X is called an ideal of X (see [9, 15]) if it satisfies:

$$0 \in I, \tag{2.1}$$

$$(\forall x, y \in X) (x * y \in I, y \in I \Rightarrow x \in I). \tag{2.2}$$

The collection of all BCK-algebras and all BCI-algebras are denoted by $\mathcal{B}_K(X)$ and $\mathcal{B}_I(X)$, respectively. Also $\mathcal{B}(X) := \mathcal{B}_K(X) \cup \mathcal{B}_I(X)$.

We refer the reader to the books [9] and [15] for further information regarding BCK/BCI-algebras.

For any family $\{a_i \mid i \in \Lambda\}$ of real numbers, we define

$$\bigvee \{a_i \mid i \in \Lambda\} = \sup \{a_i \mid i \in \Lambda\}$$

and

$$\bigwedge \{a_i \mid i \in \Lambda\} = \inf \{a_i \mid i \in \Lambda\}.$$

If $\Lambda = \{1, 2\}$, we will also use $a_1 \vee a_2$ and $a_1 \wedge a_2$ instead of $\bigvee \{a_i \mid i \in \{1, 2\}\}$ and $\bigwedge \{a_i \mid i \in \{1, 2\}\}$, respectively.

Let X be a non-empty set. A neutrosophic set (NS) in X (see [23]) is a structure of the form:

$$A := \{\langle x; A_T(x), A_I(x), A_F(x) \rangle \mid x \in X\}$$

where $A_T : X \rightarrow [0, 1]$ is a truth membership function, $A_I : X \rightarrow [0, 1]$ is an indeterminate membership function, and $A_F : X \rightarrow [0, 1]$ is a false membership function. For the sake of simplicity, we shall use the symbol $A = (A_T, A_I, A_F)$ for the neutrosophic set

$$A := \{\langle x; A_T(x), A_I(x), A_F(x) \rangle \mid x \in X\}.$$

Given a neutrosophic set $A = (A_T, A_I, A_F)$ in a set X , $\alpha, \beta \in (0, 1]$ and $\gamma \in [0, 1)$, we consider the following sets (see [10]):

$$T_\epsilon(A; \alpha) := \{x \in X \mid A_T(x) \geq \alpha\},$$

$$I_{\in}(A; \beta) := \{x \in X \mid A_I(x) \geq \beta\},$$

$$F_{\in}(A; \gamma) := \{x \in X \mid A_F(x) \leq \gamma\}.$$

We say $T_{\in}(A; \alpha)$, $I_{\in}(A; \beta)$ and $F_{\in}(A; \gamma)$ are *neutrosophic \in -subsets*.

3. Generalizations of neutrosophic ideals based on neutrosophic points

In what follows, let k_T , k_I and k_F denote arbitrary elements of $[0, 1)$ unless otherwise specified. If k_T , k_I and k_F are the same number in $[0, 1)$, then it is denoted by k , i.e., $k = k_T = k_I = k_F$.

Given a neutrosophic set $A = (A_T, A_I, A_F)$ in a set X , $\alpha, \beta \in (0, 1]$ and $\gamma \in [0, 1)$, we consider the following sets:

$$T_{q_{k_T}}(A; \alpha) := \{x \in X \mid A_T(x) + \alpha + k_T > 1\},$$

$$I_{q_{k_I}}(A; \beta) := \{x \in X \mid A_I(x) + \beta + k_I > 1\},$$

$$F_{q_{k_F}}(A; \gamma) := \{x \in X \mid A_F(x) + \gamma + k_F < 1\},$$

$$T_{\in \vee q_{k_T}}(A; \alpha) := \{x \in X \mid A_T(x) \geq \alpha \text{ or } A_T(x) + \alpha + k_T > 1\},$$

$$I_{\in \vee q_{k_I}}(A; \beta) := \{x \in X \mid A_I(x) \geq \beta \text{ or } A_I(x) + \beta + k_I > 1\},$$

$$F_{\in \vee q_{k_F}}(A; \gamma) := \{x \in X \mid A_F(x) \leq \gamma \text{ or } A_F(x) + \gamma + k_F < 1\}.$$

We say $T_{q_{k_T}}(A; \alpha)$, $I_{q_{k_I}}(A; \beta)$ and $F_{q_{k_F}}(A; \gamma)$ are *neutrosophic q_k -subsets*; and $T_{\in \vee q_{k_T}}(A; \alpha)$, $I_{\in \vee q_{k_I}}(A; \beta)$ and $F_{\in \vee q_{k_F}}(A; \gamma)$ are *neutrosophic $\in \vee q_k$ -subsets*. For $\psi \in \{\in, q, q_k, q_{k_T}, q_{k_I}, q_{k_F}, \in \vee q, \in \vee q_k, \in \vee q_{k_T}, \in \vee q_{k_I}, \in \vee q_{k_F}\}$, the element of $T_{\psi}(A; \alpha)$ (resp., $I_{\psi}(A; \beta)$ and $F_{\psi}(A; \gamma)$) is called a *neutrosophic T_{ψ} -point* (resp., *neutrosophic I_{ψ} -point* and *neutrosophic F_{ψ} -point*) with value α (resp., β and γ).

It is clear that

$$T_{\in \vee q_{k_T}}(A; \alpha) = T_{\in}(A; \alpha) \cup T_{q_{k_T}}(A; \alpha), \quad (3.1)$$

$$I_{\in \vee q_{k_I}}(A; \beta) = I_{\in}(A; \beta) \cup I_{q_{k_I}}(A; \beta), \quad (3.2)$$

$$F_{\in \vee q_{k_F}}(A; \gamma) = F_{\in}(A; \gamma) \cup F_{q_{k_F}}(A; \gamma). \quad (3.3)$$

THEOREM 3.1. *Given a neutrosophic set $A = (A_T, A_I, A_F)$ in $X \in \mathcal{B}(X)$, the following assertions are equivalent.*

(1) The nonempty neutrosophic \in -subsets $T_{\in}(A; \alpha)$, $I_{\in}(A; \beta)$ and $F_{\in}(A; \gamma)$ are ideals of X for all $\alpha \in (\frac{1-k_T}{2}, 1]$, $\beta \in (\frac{1-k_I}{2}, 1]$ and $\gamma \in [0, \frac{1-k_F}{2})$.

(2) $A = (A_T, A_I, A_F)$ satisfies the following assertion.

$$(\forall x \in X) \begin{pmatrix} A_T(x) \leq A_T(0) \vee \frac{1-k_T}{2} \\ A_I(x) \leq A_I(0) \vee \frac{1-k_I}{2} \\ A_F(x) \geq A_F(0) \wedge \frac{1-k_F}{2} \end{pmatrix} \quad (3.4)$$

and

$$(\forall x, y \in X) \begin{pmatrix} A_T(x) \vee \frac{1-k_T}{2} \geq A_T(x * y) \wedge A_T(y) \\ A_I(x) \vee \frac{1-k_I}{2} \geq A_I(x * y) \wedge A_I(y) \\ A_F(x) \wedge \frac{1-k_F}{2} \leq A_F(x * y) \vee A_F(y) \end{pmatrix} \quad (3.5)$$

PROOF: Assume that the nonempty neutrosophic \in -subsets $T_{\in}(A; \alpha)$, $I_{\in}(A; \beta)$ and $F_{\in}(A; \gamma)$ are ideals of X for all $\alpha \in (\frac{1-k_T}{2}, 1]$, $\beta \in (\frac{1-k_I}{2}, 1]$ and $\gamma \in [0, \frac{1-k_F}{2})$. If there are $a, b \in X$ such that $A_T(a) > A_T(0) \vee \frac{1-k_T}{2}$, then $a \in T_{\in}(A; \alpha_a)$ and $0 \notin T_{\in}(A; \alpha_a)$ for $\alpha_a := A_T(a) \in (\frac{1-k_T}{2}, 1]$. This is a contradiction, and so $A_T(x) \leq A_T(0) \vee \frac{1-k_T}{2}$ for all $x \in X$. We also know that $A_I(x) \leq A_I(0) \vee \frac{1-k_I}{2}$ for all $x \in X$ by the similar way. Now, let $x \in X$ be such that $A_F(x) < A_F(0) \wedge \frac{1-k_F}{2}$. If we take $\gamma_x := A_F(x)$, then $\gamma_x \in [0, \frac{1-k_F}{2})$ and so $0 \in F_{\in}(A; \gamma_x)$ since $F_{\in}(A; \gamma_x)$ is an ideal of X . Hence $A_F(0) \leq \gamma_x = A_F(x)$, which is a contradiction. Hence $A_F(x) \geq A_F(0) \wedge \frac{1-k_F}{2}$ for all $x \in X$. Suppose that $A_I(x) \vee \frac{1-k_I}{2} < A_I(x * y) \wedge A_I(y)$ for some $x, y \in X$ and take $\beta := A_I(x * y) \wedge A_I(y)$. Then $\beta \in (\frac{1-k_I}{2}, 1]$ and $x * y, y \in I_{\in}(A; \beta)$. But $x \notin I_{\in}(A; \beta)$ which is a contradiction. Thus $A_I(x) \vee \frac{1-k_I}{2} \geq A_I(x * y) \wedge A_I(y)$ for all $x, y \in X$. Similarly, we have $A_T(x) \vee \frac{1-k_T}{2} \geq A_T(x * y) \wedge A_T(y)$ for all $x, y \in X$. Suppose that there exist $x, y \in X$ such that $A_F(x) \wedge \frac{1-k_F}{2} > A_F(x * y) \vee A_F(y)$. Taking $\gamma := A_F(x * y) \vee A_F(y)$ implies that $\gamma \in [0, \frac{1-k_F}{2})$, $x * y \in F_{\in}(A; \gamma)$ and $y \in F_{\in}(A; \gamma)$, but $x \notin F_{\in}(A; \gamma)$. This is a contradiction, and so $A_F(x) \wedge \frac{1-k_F}{2} \leq A_F(x * y) \vee A_F(y)$ for all $x, y \in X$.

Conversely, suppose that $A = (A_T, A_I, A_F)$ satisfies two conditions (3.4) and (3.5). Let $\alpha \in (\frac{1-k_T}{2}, 1]$, $\beta \in (\frac{1-k_I}{2}, 1]$ and $\gamma \in [0, \frac{1-k_F}{2})$ be such that $T_{\in}(A; \alpha)$, $I_{\in}(A; \beta)$ and $F_{\in}(A; \gamma)$ are nonempty. For any $x \in T_{\in}(A; \alpha)$, $y \in I_{\in}(A; \beta)$ and $z \in F_{\in}(A; \gamma)$, we get

$$\begin{aligned} A_T(0) \vee \frac{1-k_T}{2} &\geq A_T(x) \geq \alpha > \frac{1-k_T}{2}, \\ A_I(0) \vee \frac{1-k_I}{2} &\geq A_I(y) \geq \beta > \frac{1-k_I}{2}, \\ A_F(0) \wedge \frac{1-k_F}{2} &\leq A_F(z) \leq \gamma < \frac{1-k_F}{2}, \end{aligned}$$

and so $A_T(0) \geq \alpha$, $A_I(0) \geq \beta$ and $A_F(0) \leq \gamma$. Hence $0 \in T_{\in}(A; \alpha)$, $0 \in I_{\in}(A; \beta)$ and $0 \in F_{\in}(A; \gamma)$. Let $a, b, x, y, u, v \in X$ be such that $a * b \in T_{\in}(A; \alpha)$, $b \in T_{\in}(A; \alpha)$, $x * y \in I_{\in}(A; \beta)$, $y \in I_{\in}(A; \beta)$, $u * v \in F_{\in}(A; \gamma)$, and $v \in F_{\in}(A; \gamma)$. It follows from (3.5) that

$$\begin{aligned} A_T(a) \vee \frac{1-k_T}{2} &\geq A_T(a * b) \wedge A_T(b) \geq \alpha > \frac{1-k_T}{2}, \\ A_I(x) \vee \frac{1-k_I}{2} &\geq A_I(x * y) \wedge A_I(y) \geq \beta > \frac{1-k_I}{2}, \\ A_F(u) \wedge \frac{1-k_F}{2} &\leq A_F(u * v) \vee A_F(v) \leq \gamma < \frac{1-k_F}{2}. \end{aligned}$$

Hence $A_T(a) \geq \alpha$, $A_I(x) \geq \beta$ and $A_F(u) \leq \gamma$, that is, $a \in T_{\in}(A; \alpha)$, $x \in I_{\in}(A; \beta)$ and $u \in F_{\in}(A; \gamma)$. Therefore $T_{\in}(A; \alpha)$, $I_{\in}(A; \beta)$ and $F_{\in}(A; \gamma)$ are ideals of X for all $\alpha \in (\frac{1-k_T}{2}, 1]$, $\beta \in (\frac{1-k_I}{2}, 1]$ and $\gamma \in [0, \frac{1-k_F}{2})$. \square

COROLLARY 3.2 ([21]). Given a neutrosophic set $A = (A_T, A_I, A_F)$ in $X \in \mathcal{B}(X)$, the following assertions are equivalent.

(1) The nonempty neutrosophic \in -subsets $T_{\in}(A; \alpha)$, $I_{\in}(A; \beta)$ and $F_{\in}(A; \gamma)$ are ideals of X for all $\alpha, \beta \in (0.5, 1]$ and $\gamma \in [0, 0.5)$.

(2) $A = (A_T, A_I, A_F)$ satisfies the following assertion.

$$(\forall x \in X) \begin{pmatrix} A_T(x) \leq A_T(0) \vee 0.5 \\ A_I(x) \leq A_I(0) \vee 0.5 \\ A_F(x) \geq A_F(0) \wedge 0.5 \end{pmatrix}$$

and

$$(\forall x, y \in X) \begin{pmatrix} A_T(x) \vee 0.5 \geq A_T(x * y) \wedge A_T(y) \\ A_I(x) \vee 0.5 \geq A_I(x * y) \wedge A_I(y) \\ A_F(x) \wedge 0.5 \leq A_F(x * y) \vee A_F(y) \end{pmatrix}$$

DEFINITION 3.3. A neutrosophic set $A = (A_T, A_I, A_F)$ in $X \in \mathcal{B}(X)$ is called an $(\in, \in \vee q_{(k_T, k_I, k_F)})$ -neutrosophic ideal of X if the following assertions are valid.

$$(\forall x \in X) \left(\begin{array}{l} x \in T_{\in}(A; \alpha_x) \Rightarrow 0 \in T_{\in \vee q_{k_T}}(A; \alpha_x) \\ x \in I_{\in}(A; \beta_x) \Rightarrow 0 \in I_{\in \vee q_{k_I}}(A; \beta_x) \\ x \in F_{\in}(A; \gamma_x) \Rightarrow 0 \in F_{\in \vee q_{k_F}}(A; \gamma_x) \end{array} \right), \quad (3.6)$$

$$(\forall x, y \in X) \left(\begin{array}{l} x * y \in T_{\in}(A; \alpha_x), y \in T_{\in}(A; \alpha_y) \Rightarrow x \in T_{\in \vee q_{k_T}}(A; \alpha_x \wedge \alpha_y) \\ x * y \in I_{\in}(A; \beta_x), y \in I_{\in}(A; \beta_y) \Rightarrow x \in I_{\in \vee q_{k_I}}(A; \beta_x \wedge \beta_y) \\ x * y \in F_{\in}(A; \gamma_x), y \in F_{\in}(A; \gamma_y) \Rightarrow x \in F_{\in \vee q_{k_F}}(A; \gamma_x \vee \gamma_y) \end{array} \right) \quad (3.7)$$

for all $\alpha_x, \alpha_y, \beta_x, \beta_y \in (0, 1]$ and $\gamma_x, \gamma_y \in [0, 1]$.

Example 3.4. Let $X = \{0, 1, 2, 3, 4\}$ be a set with the binary operation $*$ which is given in Table 1.

Table 1: Cayley table for the binary operation “ $*$ ”

$*$	0	1	2	3	4
0	0	0	0	0	0
1	1	0	1	0	1
2	2	2	0	2	0
3	3	1	3	0	3
4	4	4	4	4	0

Then $(X, *, 0)$ is a BCK-algebra (see [15]). Consider a neutrosophic set $A = (A_T, A_I, A_F)$ in X which is given by Table 2.

Table 2: Tabular representation of $A = (A_T, A_I, A_F)$

X	$A_T(x)$	$A_I(x)$	$A_F(x)$
0	0.6	0.5	0.45
1	0.5	0.3	0.93
2	0.3	0.7	0.67
3	0.4	0.3	0.93
4	0.1	0.2	0.74

Routine calculations show that $A = (A_T, A_I, A_F)$ is an $(\in, \in \vee q_{(k_T, k_I, k_F)})$ -neutrosophic ideal of X for $k_T = 0.24$, $k_I = 0.08$ and $k_F = 0.16$.

THEOREM 3.5. *A neutrosophic set $A = (A_T, A_I, A_F)$ in $X \in \mathcal{B}(X)$ is an $(\in, \in \vee q_{(k_T, k_I, k_F)})$ -neutrosophic ideal of $X \in \mathcal{B}(X)$ if and only if $A = (A_T, A_I, A_F)$ satisfies the following assertions.*

$$(\forall x \in X) \left(\begin{array}{l} A_T(0) \geq A_T(x) \wedge \frac{1-k_T}{2} \\ A_I(0) \geq A_I(x) \wedge \frac{1-k_I}{2} \\ A_F(0) \leq A_F(x) \vee \frac{1-k_F}{2} \end{array} \right), \quad (3.8)$$

$$(\forall x, y \in X) \left(\begin{array}{l} A_T(x) \geq \wedge \{A_T(x * y), A_T(y), \frac{1-k_T}{2}\} \\ A_I(x) \geq \wedge \{A_I(x * y), A_I(y), \frac{1-k_I}{2}\} \\ A_F(x) \leq \vee \{A_F(x * y), A_F(y), \frac{1-k_F}{2}\} \end{array} \right). \quad (3.9)$$

PROOF: Assume that $A = (A_T, A_I, A_F)$ in $X \in \mathcal{B}(X)$ is an $(\in, \in \vee q_{(k_T, k_I, k_F)})$ -neutrosophic ideal of $X \in \mathcal{B}(X)$. If $A_T(0) < A_T(a) \wedge \frac{1-k_T}{2}$ for some $a \in X$, then there exists $\alpha_a \in (0, 1]$ such that $A_T(0) < \alpha_a \leq A_T(a) \wedge \frac{1-k_T}{2}$. It follows that $\alpha_a \in (0, \frac{1-k_T}{2}]$, $a \in T_{\in}(A; \alpha_a)$ and $0 \notin T_{\in}(A; \alpha_a)$. Also, $A_T(0) + \alpha_a + k_T < 2\alpha_a + k_T \leq 1$, i.e., $0 \notin T_{q_{k_T}}(A; \alpha_a)$. Hence $0 \notin T_{\in \vee q_{k_T}}(A; \alpha_a)$, a contradiction. Thus $A_T(0) \geq A_T(x) \wedge \frac{1-k_T}{2}$ for all $x \in X$. Similarly, we have $A_I(0) \geq A_I(x) \wedge \frac{1-k_I}{2}$ for all $x \in X$. Suppose that $A_F(0) > A_F(z) \vee \frac{1-k_F}{2}$ for some $z \in X$ and take $\gamma_z := A_F(z) \vee \frac{1-k_F}{2}$. Then $\gamma_z \geq \frac{1-k_F}{2}$, $z \in F_{\in}(A; \gamma_z)$ and $0 \notin F_{\in}(A; \gamma_z)$. Also $A_F(0) + \gamma_z + k_F \geq 1$, that is, $0 \notin F_{q_{k_F}}(A; \gamma_z)$. This is a contradiction, and thus $A_F(0) \leq A_F(x) \vee \frac{1-k_F}{2}$ for all $x \in X$. Suppose that $A_I(a) < \wedge \{A_I(a * b), A_I(b), \frac{1-k_I}{2}\}$ for some $a, b \in X$ and take $\beta := \wedge \{A_I(a * b), A_I(b), \frac{1-k_I}{2}\}$. Then $\beta \leq \frac{1-k_I}{2}$, $a * b \in I_{\in}(A; \beta)$, $b \in I_{\in}(A; \beta)$ and $a \notin I_{\in}(A; \beta)$. Also, we have $A_I(a) + \beta + k_I \leq 1$, i.e., $a \notin I_{q_{k_F}}(A; \beta)$. This is impossible, and therefore $A_I(x) \geq \wedge \{A_I(x * y), A_I(y), \frac{1-k_I}{2}\}$ for all $x, y \in X$. By the similar way, we can verify that $A_T(x) \geq \wedge \{A_T(x * y), A_T(y), \frac{1-k_T}{2}\}$ for all $x, y \in X$. Now assume that $A_F(a) > \vee \{A_F(a * b), A_F(b), \frac{1-k_F}{2}\}$ for some $a, b \in X$. Then there exists $\gamma \in [0, 1)$ such that $A_F(a) > \gamma \geq \vee \{A_F(a * b), A_F(b), \frac{1-k_F}{2}\}$. Then $\gamma \geq \frac{1-k_F}{2}$, $a * b \in F_{\in}(A; \gamma)$, $b \in F_{\in}(A; \gamma)$ and $a \notin F_{\in}(A; \gamma)$. Also, $A_F(a) + \gamma + k_F \geq 1$, i.e., $a \notin F_{q_{k_F}}(A; \gamma)$. Thus $a \notin F_{\in \vee q_{k_F}}(A; \gamma)$, which is a contradiction. Hence $A_F(x) \leq \vee \{A_F(x * y), A_F(y), \frac{1-k_F}{2}\}$ for all $x, y \in X$.

Conversely, suppose that $A = (A_T, A_I, A_F)$ satisfies two conditions (3.8) and (3.9). For any $x, y, z \in X$, let $\alpha_x, \beta_y \in (0, 1]$ and $\gamma_z \in [0, 1]$ be such that $x \in T_{\in}(A; \alpha_x)$, $y \in I_{\in}(A; \beta_y)$ and $z \in F_{\in}(A; \gamma_z)$. Then $A_T(x) \geq \alpha_x$, $A_I(y) \geq \beta_y$ and $A_F(z) \leq \gamma_z$. Assume that $A_T(0) < \alpha_x$, $A_I(0) < \beta_y$ and $A_F(0) > \gamma_z$. If $A_T(x) < \frac{1-k_T}{2}$, then

$$A_T(0) \geq A_T(x) \wedge \frac{1-k_T}{2} = A_T(x) \geq \alpha_x,$$

a contradiction. Hence $A_T(x) \geq \frac{1-k_T}{2}$, and so

$$A_T(0) + \alpha_x + k_T > 2A_T(0) + k_T \geq 2 \left(A_T(x) \wedge \frac{1-k_T}{2} \right) + k_T = 1.$$

Hence $0 \in T_{q_{k_T}}(A; \alpha_x) \subseteq T_{\in \vee q_{k_T}}(A; \alpha_x)$. Similarly, we get $0 \in I_{q_{k_I}}(A; \beta_y) \subseteq I_{\in \vee q_{k_I}}(A; \beta_y)$. If $A_F(z) > \frac{1-k_F}{2}$, then $A_F(0) \leq A_F(z) \vee \frac{1-k_F}{2} = A_F(z) \leq \gamma_z$ which is a contradiction. Hence $A_F(z) \leq \frac{1-k_F}{2}$, and thus

$$A_F(0) + \gamma_z + k_F < 2A_F(0) + k_F \leq 2 \left(A_F(z) \vee \frac{1-k_F}{2} \right) + k_F = 1.$$

Hence $0 \in F_{q_{k_F}}(A; \gamma_z) \subseteq F_{\in \vee q_{k_F}}(A; \gamma_z)$. For any $a, b, p, q, x, y \in X$, let $\alpha_a, \alpha_b, \beta_p, \beta_q \in (0, 1]$ and $\gamma_x, \gamma_y \in [0, 1]$ be such that $a * b \in T_{\in}(A; \alpha_a)$, $b \in T_{\in}(A; \alpha_b)$, $p * q \in I_{\in}(A; \beta_p)$, $q \in I_{\in}(A; \beta_q)$, $x * y \in F_{\in}(A; \gamma_x)$, and $y \in F_{\in}(A; \gamma_y)$. Then $A_T(a * b) \geq \alpha_a$, $A_T(b) \geq \alpha_b$, $A_I(p * q) \geq \beta_p$, $A_I(q) \geq \beta_q$, $A_F(x * y) \leq \gamma_x$, and $A_F(y) \leq \gamma_y$. Suppose that $a \notin T_{\in}(A; \alpha_a \wedge \alpha_b)$. Then $A_T(a) < \alpha_a \wedge \alpha_b$. If $A_T(a * b) \wedge A_T(b) < \frac{1-k_T}{2}$, then

$$A_T(a) \geq \bigwedge \{ A_T(a * b), A_T(b), \frac{1-k_T}{2} \} = A_T(a * b) \wedge A_T(b) \geq \alpha_a \wedge \alpha_b.$$

This is a contradiction, and so $A_T(a * b) \wedge A_T(b) \geq \frac{1-k_T}{2}$. Thus

$$\begin{aligned} A_T(a) + (\alpha_a \wedge \alpha_b) + k_T &> 2A_T(a) + k_T \\ &\geq 2 \left(\bigwedge \{ A_T(a * b), A_T(b), \frac{1-k_T}{2} \} \right) + k_T = 1, \end{aligned}$$

which induces $a \in T_{q_{k_T}}(A; \alpha_a \wedge \alpha_b) \subseteq T_{\in \vee q_{k_T}}(A; \alpha_a \wedge \alpha_b)$. By the similarly way, we get $p \in I_{\in \vee q_{k_I}}(A; \beta_p \wedge \beta_q)$. Suppose that $x \notin F_{\in}(A; \gamma_x \vee \gamma_y)$, that is, $A_F(x) > \gamma_x \vee \gamma_y$. If $A_F(x * y) \vee A_F(y) > \frac{1-k_F}{2}$, then

$$A_F(x) \leq \bigvee \{ A_F(x * y), A_F(y), \frac{1-k_F}{2} \} = A_F(x * y) \vee A_F(y) \leq \gamma_x \vee \gamma_y,$$

which is impossible. Thus $A_F(x * y) \vee A_F(y) \leq \frac{1-k_F}{2}$, and so

$$\begin{aligned}
 A_F(x) + (\gamma_x \vee \gamma_y) + k_F &< 2A_F(x) \\
 &\leq 2 \left(\bigvee \{A_F(x * y), A_F(y), \frac{1-k_F}{2}\} \right) + k_F = 1.
 \end{aligned}$$

This implies that $x \in F_{q_{k_F}}(A; \gamma_x \vee \gamma_y) \subseteq F_{\in \vee q_{k_F}}(A; \gamma_x \vee \gamma_y)$. Consequently, $A = (A_T, A_I, A_F)$ is an $(\in, \in \vee q_{(k_T, k_I, k_F)})$ -neutrosophic ideal of $X \in \mathcal{B}(X)$. \square

COROLLARY 3.6 ([21]). For a neutrosophic set $A = (A_T, A_I, A_F)$ in $X \in \mathcal{B}(X)$, the following are equivalent.

- (1) $A = (A_T, A_I, A_F)$ is an $(\in, \in \vee q)$ -neutrosophic ideal of $X \in \mathcal{B}(X)$.
- (2) $A = (A_T, A_I, A_F)$ satisfies the following assertions.

$$\begin{aligned}
 (\forall x \in X) &\left(\begin{array}{l} A_T(0) \geq A_T(x) \wedge 0.5 \\ A_I(0) \geq A_I(x) \wedge 0.5 \\ A_F(0) \leq A_F(x) \vee 0.5 \end{array} \right), \\
 (\forall x, y \in X) &\left(\begin{array}{l} A_T(x) \geq \bigwedge \{A_T(x * y), A_T(y), 0.5\} \\ A_I(x) \geq \bigwedge \{A_I(x * y), A_I(y), 0.5\} \\ A_F(x) \leq \bigvee \{A_F(x * y), A_F(y), 0.5\} \end{array} \right).
 \end{aligned}$$

THEOREM 3.7. *A neutrosophic set $A = (A_T, A_I, A_F)$ in $X \in \mathcal{B}(X)$ is an $(\in, \in \vee q_{(k_T, k_I, k_F)})$ -neutrosophic ideal of $X \in \mathcal{B}(X)$ if and only if the nonempty neutrosophic \in -subsets $T_{\in}(A; \alpha)$, $I_{\in}(A; \beta)$ and $F_{\in}(A; \gamma)$ are ideals of X for all $\alpha \in (0, \frac{1-k_T}{2}]$, $\beta \in (0, \frac{1-k_I}{2}]$ and $\gamma \in [\frac{1-k_F}{2}, 1)$.*

PROOF: Suppose that $A = (A_T, A_I, A_F)$ is an $(\in, \in \vee q_{(k_T, k_I, k_F)})$ -neutrosophic ideal of $X \in \mathcal{B}(X)$ and let $\alpha \in (0, \frac{1-k_T}{2}]$, $\beta \in (0, \frac{1-k_I}{2}]$ and $\gamma \in [\frac{1-k_F}{2}, 1)$ be such that $T_{\in}(A; \alpha)$, $I_{\in}(A; \beta)$ and $F_{\in}(A; \gamma)$ are nonempty. Using (3.8), we get $A_T(0) \geq A_T(x) \wedge \frac{1-k_T}{2}$, $A_I(0) \geq A_I(y) \wedge \frac{1-k_I}{2}$, and $A_F(0) \leq A_F(z) \vee \frac{1-k_F}{2}$ for all $x \in T_{\in}(A; \alpha)$, $y \in I_{\in}(A; \beta)$ and $z \in F_{\in}(A; \gamma)$. It follows that $A_T(0) \geq \alpha \wedge \frac{1-k_T}{2} = \alpha$, $A_I(0) \geq \beta \wedge \frac{1-k_I}{2} = \beta$, and $A_F(0) \leq \gamma \vee \frac{1-k_F}{2} = \gamma$, that is, $0 \in T_{\in}(A; \alpha)$, $0 \in I_{\in}(A; \beta)$ and $0 \in F_{\in}(A; \gamma)$. Let $x, y, a, b, u, v \in X$ be such that $x * y \in T_{\in}(A; \alpha)$, $y \in T_{\in}(A; \alpha)$, $a * b \in I_{\in}(A; \beta)$, $b \in I_{\in}(A; \beta)$, $u * v \in F_{\in}(A; \gamma)$, and $v \in F_{\in}(A; \gamma)$ for $\alpha \in (0, \frac{1-k_T}{2}]$, $\beta \in (0, \frac{1-k_I}{2}]$ and $\gamma \in [\frac{1-k_F}{2}, 1)$. Then $A_T(x * y) \geq \alpha$, $A_T(y) \geq \alpha$, $A_I(a * b) \geq \beta$, $A_I(b) \geq \beta$, $A_F(u * v) \leq \gamma$, and $A_F(v) \leq \gamma$. It follows from (3.9) that

A general model of neutrosophic ideals in BCK/BCI-algebras...

$$A_T(x) \geq \bigwedge \{A_T(x * y), A_T(y), \frac{1-k_T}{2}\} \geq \alpha \wedge \frac{1-k_T}{2} = \alpha,$$

$$A_I(a) \geq \bigwedge \{A_I(a * b), A_I(b), \frac{1-k_I}{2}\} \geq \beta \wedge \frac{1-k_I}{2} = \beta,$$

$$A_F(u) \leq \bigvee \{A_F(u * v), A_F(v), \frac{1-k_F}{2}\} \leq \gamma \vee \frac{1-k_F}{2} = \gamma$$

and so that $x \in T_\infty(A; \alpha)$, $a \in I_\infty(A; \beta)$ and $u \in F_\infty(A; \gamma)$. Therefore $T_\infty(A; \alpha)$, $I_\infty(A; \beta)$ and $F_\infty(A; \gamma)$ are ideals of X for all $\alpha \in (0, \frac{1-k_T}{2}]$, $\beta \in (0, \frac{1-k_I}{2}]$ and $\gamma \in [\frac{1-k_F}{2}, 1)$.

Conversely, let $A = (A_T, A_I, A_F)$ be a neutrosophic set in $X \in \mathcal{B}(X)$ such that the nonempty neutrosophic \in -subsets $T_\infty(A; \alpha)$, $I_\infty(A; \beta)$ and $F_\infty(A; \gamma)$ are ideals of X for all $\alpha \in (0, \frac{1-k_T}{2}]$, $\beta \in (0, \frac{1-k_I}{2}]$ and $\gamma \in [\frac{1-k_F}{2}, 1)$. If there exist $x, y, z \in X$ such that $A_T(0) < A_T(x) \wedge \frac{1-k_T}{2}$, $A_I(0) < A_I(y) \wedge \frac{1-k_I}{2}$, and $A_F(0) > A_F(z) \vee \frac{1-k_F}{2}$, then $0 \notin T_\infty(A; \alpha_x)$, $0 \notin I_\infty(A; \beta_y)$ and $0 \notin F_\infty(A; \gamma_z)$ by taking $\alpha_x := A_T(x) \wedge \frac{1-k_T}{2}$, $\beta_y := A_I(y) \wedge \frac{1-k_I}{2}$, and $\gamma_z := A_F(z) \vee \frac{1-k_F}{2}$. This is a contradiction, and so $A_T(0) \geq A_T(x) \wedge \frac{1-k_T}{2}$, $A_I(0) \geq A_I(y) \wedge \frac{1-k_I}{2}$, and $A_F(0) \leq A_F(x) \vee \frac{1-k_F}{2}$ for all $x \in X$. Now, suppose that there $x, y, a, b, u, v \in X$ be such that $A_T(x) < \bigwedge \{A_T(x * y), A_T(y), \frac{1-k_T}{2}\}$, $A_I(a) < \bigwedge \{A_I(a * b), A_I(b), \frac{1-k_I}{2}\}$, and $A_F(u) > \bigvee \{A_F(u * v), A_F(v), \frac{1-k_F}{2}\}$. If we take $\alpha := \bigwedge \{A_T(x * y), A_T(y), \frac{1-k_T}{2}\}$, $\beta := \bigwedge \{A_I(a * b), A_I(b), \frac{1-k_I}{2}\}$, and $\gamma := \bigvee \{A_F(u * v), A_F(v), \frac{1-k_F}{2}\}$, then $\alpha \leq \frac{1-k_T}{2}$, $\beta \leq \frac{1-k_I}{2}$, $\gamma \geq \frac{1-k_F}{2}$, $x * y \in T_\infty(A; \alpha)$, $y \in T_\infty(A; \alpha)$, $a * b \in I_\infty(A; \beta)$, $b \in I_\infty(A; \beta)$, $u * v \in F_\infty(A; \gamma)$, and $v \in F_\infty(A; \gamma)$. But $x \notin T_\infty(A; \alpha)$, $a \notin I_\infty(A; \beta)$ and $u \notin F_\infty(A; \gamma)$, which induces a contradiction. Therefore $A_T(x) \geq \bigwedge \{A_T(x * y), A_T(y), \frac{1-k_T}{2}\}$, $A_I(x) \geq \bigwedge \{A_I(x * y), A_I(y), \frac{1-k_I}{2}\}$, and $A_F(x) \leq \bigvee \{A_F(x * y), A_F(y), \frac{1-k_F}{2}\}$ for all $x, y \in X$. Using Theorem 3.5, we conclude that $A = (A_T, A_I, A_F)$ is an $(\in, \in \vee q_{(k_T, k_I, k_F)})$ -neutrosophic ideal of $X \in \mathcal{B}(X)$. \square

COROLLARY 3.8 ([21]). A neutrosophic set $A = (A_T, A_I, A_F)$ in $X \in \mathcal{B}(X)$ is an $(\in, \in \vee q)$ -neutrosophic ideal of $X \in \mathcal{B}(X)$ if and only if the nonempty neutrosophic \in -subsets $T_\infty(A; \alpha)$, $I_\infty(A; \beta)$ and $F_\infty(A; \gamma)$ are ideals of X for all $\alpha, \beta \in (0, 0.5]$ and $\gamma \in [0.5, 1)$.

It is clear that every (\in, \in) -neutrosophic ideal is an $(\in, \in \vee q_{(k_T, k_I, k_F)})$ -neutrosophic ideal. But the converse is not true in general. For example, the $(\in, \in \vee q_{(k_T, k_I, k_F)})$ -neutrosophic ideal $A = (A_T, A_I, A_F)$ with $k_T = 0.24$, $k_I = 0.08$ and $k_F = 0.16$ in Example 3.4 is not an (\in, \in) -neutrosophic ideal since $2 \in I_\infty(A; 0.56)$ and $0 \notin I_\infty(A; 0.56)$.

We now consider conditions for an $(\in, \in \vee q_{(k_T, k_I, k_F)})$ -neutrosophic ideal to be an (\in, \in) -neutrosophic ideal.

THEOREM 3.9. *Let $A = (A_T, A_I, A_F)$ be an $(\in, \in \vee q_{(k_T, k_I, k_F)})$ -neutrosophic ideal of $X \in \mathcal{B}(X)$ such that*

$$(\forall x \in X) (A_T(x) < \frac{1-k_T}{2}, A_I(x) < \frac{1-k_I}{2}, A_F(x) > \frac{1-k_F}{2}).$$

Then $A = (A_T, A_I, A_F)$ is an (\in, \in) -neutrosophic ideal of $X \in \mathcal{B}(X)$.

PROOF: Let $x, y, z \in X$, $\alpha, \beta \in (0, 1]$ and $\gamma \in [0, 1)$ be such that $x \in T_{\in}(A; \alpha)$, $y \in I_{\in}(A; \beta)$ and $z \in F_{\in}(A; \gamma)$. Then $A_T(x) \geq \alpha$, $A_I(y) \geq \beta$ and $A_F(z) \leq \gamma$. It follows from (3.8) that

$$A_T(0) \geq A_T(x) \wedge \frac{1-k_T}{2} = A_T(x) \geq \alpha,$$

$$A_I(0) \geq A_I(y) \wedge \frac{1-k_I}{2} = A_I(y) \geq \beta,$$

$$A_F(0) \leq A_F(z) \vee \frac{1-k_F}{2} = A_F(z) \leq \gamma.$$

Hence $0 \in T_{\in}(A; \alpha)$, $0 \in I_{\in}(A; \beta)$ and $0 \in F_{\in}(A; \gamma)$. For any $x, y, a, b, u, v \in X$, let $\alpha_x, \alpha_y, \beta_a, \beta_b \in (0, 1]$ and $\gamma_u, \gamma_v \in [0, 1)$ be such that $x * y \in T_{\in}(A; \alpha_x)$, $y \in T_{\in}(A; \alpha_y)$, $a * b \in I_{\in}(A; \beta_a)$, $b \in I_{\in}(A; \beta_b)$, $u * v \in F_{\in}(A; \gamma_u)$, and $v \in F_{\in}(A; \gamma_v)$. Then $A_T(x * y) \geq \alpha_x$, $A_T(y) \geq \alpha_y$, $A_I(a * b) \geq \beta_a$, $A_I(b) \geq \beta_b$, $A_F(u * v) \leq \gamma_u$, and $A_F(v) \leq \gamma_v$. It follows from (3.9) that

$$A_T(x) \geq \bigwedge \{A_T(x * y), A_T(y), \frac{1-k_T}{2}\} = A_T(x * y) \wedge A_T(y) \geq \alpha_x \wedge \alpha_y,$$

$$A_I(a) \geq \bigwedge \{A_I(a * b), A_I(b), \frac{1-k_I}{2}\} = A_I(a * b) \wedge A_I(b) \geq \beta_a \wedge \beta_b,$$

$$A_F(u) \leq \bigvee \{A_F(u * v), A_F(v), \frac{1-k_F}{2}\} = A_F(u * v) \vee A_F(v) \leq \gamma_u \vee \gamma_v.$$

Thus $x \in T_{\in}(A; \alpha_x \wedge \alpha_y)$, $a \in I_{\in}(A; \beta_a \wedge \beta_b)$ and $u \in F_{\in}(A; \gamma_u \vee \gamma_v)$. Therefore $A = (A_T, A_I, A_F)$ is an (\in, \in) -neutrosophic ideal of $X \in \mathcal{B}(X)$. \square

COROLLARY 3.10 ([21]). *Let $A = (A_T, A_I, A_F)$ be an $(\in, \in \vee q)$ -neutrosophic ideal of $X \in \mathcal{B}(X)$ such that*

$$(\forall x \in X) (A_T(x) < 0.5, A_I(x) < 0.5, A_F(x) > 0.5).$$

Then $A = (A_T, A_I, A_F)$ is an (\in, \in) -neutrosophic ideal of $X \in \mathcal{B}(X)$.

THEOREM 3.11. *Given a neutrosophic set $A = (A_T, A_I, A_F)$ in $X \in \mathcal{B}(X)$, if the nonempty neutrosophic $\in \vee q_k$ -subsets $T_{\in \vee q_{k_T}}(A; \alpha)$, $I_{\in \vee q_{k_I}}(A; \beta)$*

and $F_{\in \vee q_{k_F}}(A; \gamma)$ are ideals of X for all $\alpha \in (0, \frac{1-k_T}{2}]$, $\beta \in (0, \frac{1-k_I}{2}]$ and $\gamma \in [\frac{1-k_F}{2}, 1)$, then $A = (A_T, A_I, A_F)$ is an $(\in, \in \vee q_{(k_T, k_I, k_F)})$ -neutrosophic ideal of X .

PROOF: Let $A = (A_T, A_I, A_F)$ be a neutrosophic set in $X \in \mathcal{B}(X)$ such that the nonempty neutrosophic $\in \vee q_k$ -subsets $T_{\in \vee q_{k_T}}(A; \alpha)$, $I_{\in \vee q_{k_I}}(A; \beta)$ and $F_{\in \vee q_{k_F}}(A; \gamma)$ are ideals of X for all $\alpha \in (0, \frac{1-k_T}{2}]$, $\beta \in (0, \frac{1-k_I}{2}]$ and $\gamma \in [\frac{1-k_F}{2}, 1)$. If $A_T(0) < A_T(x) \wedge \frac{1-k_T}{2} := \alpha_x$, $A_I(0) < A_I(y) \wedge \frac{1-k_I}{2} := \beta_y$ and $A_F(0) > A_F(z) \vee \frac{1-k_F}{2} := \gamma_z$ for some $x, y, z \in X$, then $x \in T_{\in}(A; \alpha_x) \subseteq T_{\in \vee q_{k_T}}(A; \alpha_x)$, $y \in I_{\in}(A; \beta_y) \subseteq I_{\in \vee q_{k_I}}(A; \beta_y)$, $z \in F_{\in}(A; \gamma_z) \subseteq F_{\in \vee q_{k_F}}(A; \gamma_z)$, $0 \notin T_{\in}(A; \alpha_x)$, $0 \notin I_{\in}(A; \beta_y)$, and $0 \notin F_{\in}(A; \gamma_z)$. Also, since $A_T(0) + \alpha_x + k_T < 2\alpha_x + k_T \leq 1$, i.e., $0 \notin T_{q_{k_T}}(A; \alpha_x)$, $A_I(0) + \beta_y + k_I < 2\beta_y + k_I \leq 1$, i.e., $0 \notin I_{q_{k_I}}(A; \beta_y)$, $A_F(0) + \gamma_z + k_F > 2\gamma_z + k_F \geq 1$, i.e., $0 \notin F_{q_{k_F}}(A; \gamma_z)$, we get $0 \notin T_{\in \vee q_{k_T}}(A; \alpha_x)$, $0 \notin I_{\in \vee q_{k_I}}(A; \beta_y)$, and $0 \notin F_{\in \vee q_{k_F}}(A; \gamma_z)$. This is a contradiction, and thus (3.8) is valid. Suppose that there exist $a, b \in X$ such that $A_I(a) < \bigwedge \{A_I(a * b), A_I(b), \frac{1-k_I}{2}\}$. Taking $\beta := \bigwedge \{A_I(a * b), A_I(b), \frac{1-k_I}{2}\}$ implies that $a * b \in I_{\in}(A; \beta) \subseteq I_{\in \vee q_{k_I}}(A; \beta)$, $b \in I_{\in}(A; \beta) \subseteq I_{\in \vee q_{k_I}}(A; \beta)$. Since $I_{\in \vee q_{k_I}}(A; \beta)$ is an ideal of X , it follows that $a \in I_{\in \vee q_{k_I}}(A; \beta)$, i.e., $a \in I_{\in}(A; \beta)$ or $a \in I_{q_{k_I}}(A; \beta)$, and so that $a \in I_{q_{k_I}}(A; \beta)$, i.e., $A_I(a) + \beta + k_I > 1$, since $a \notin I_{\in}(A; \beta)$. But $A_I(a) + \beta + k_I < 2\beta + k_I \leq 1$, a contradiction. Hence $A_I(x) \geq \bigwedge \{A_I(x * y), A_I(y), \frac{1-k_I}{2}\}$ for all $x, y \in X$. Similarly, we can verify that $A_T(x) \geq \bigwedge \{A_T(x * y), A_T(y), \frac{1-k_T}{2}\}$ for all $x, y \in X$. Assume that $A_F(a) > \bigvee \{A_F(a * b), A_F(b), \frac{1-k_F}{2}\} := \gamma$ for some $a, b \in X$. Then $a \notin F_{\in}(A; \gamma)$, $a * b \in F_{\in}(A; \gamma) \subseteq F_{\in \vee q_{k_F}}(A; \gamma)$, $b \in F_{\in}(A; \gamma) \subseteq F_{\in \vee q_{k_F}}(A; \gamma)$. Since $F_{\in \vee q_{k_F}}(A; \gamma)$ is an ideal of X , we have $a \in F_{\in \vee q_{k_F}}(A; \gamma)$. On the other hand, $A_F(a) + \gamma + k_F > 2\gamma + k_F \geq 1$, that is, $a \notin F_{q_{k_F}}(A; \gamma)$. Hence $a \notin F_{\in \vee q_{k_F}}(A; \gamma)$, a contradiction. Thus $A_F(x) \leq \bigvee \{A_F(x * y), A_F(y), \frac{1-k_F}{2}\}$ for all $x, y \in X$. Therefore (3.9) is valid, and consequently $A = (A_T, A_I, A_F)$ is an $(\in, \in \vee q_{(k_T, k_I, k_F)})$ -neutrosophic ideal of X by Theorem 3.5. \square

COROLLARY 3.12 ([21]). Given a neutrosophic set $A = (A_T, A_I, A_F)$ in $X \in \mathcal{B}(X)$, if the nonempty neutrosophic $\in \vee q$ -subsets $T_{\in \vee q}(A; \alpha)$, $I_{\in \vee q}(A; \beta)$ and $F_{\in \vee q}(A; \gamma)$ are ideals of X for all $\alpha, \beta \in (0, 0.5]$ and $\gamma \in [0.5, 1)$, then $A = (A_T, A_I, A_F)$ is an $(\in, \in \vee q)$ -neutrosophic ideal of X .

4. Conclusions

More general form of $(\in, \in \vee q)$ -neutrosophic ideal was introduced, and their properties were investigated. Relations between (\in, \in) -neutrosophic ideal and $(\in, \in \vee q_{(k_T, k_I, k_F)})$ -neutrosophic ideal were discussed. Characterizations of $(\in, \in \vee q_{(k_T, k_I, k_F)})$ -neutrosophic ideal were discussed, and conditions for a neutrosophic set to be an $(\in, \in \vee q_{(k_T, k_I, k_F)})$ -neutrosophic ideal were displayed.

These results can be applied to characterize the neutrosophic ideals in a BCK/BCI -algebra. In our future research, we will focus on some properties of ideal such as intersections, unions, maximality, primeness and height, and try to find the relations between these properties of ideals and the results of this paper. For instance, how we can define the prime and maximal neutrosophic ideals? What is the meaning of height of these types of ideals? For information about the maximality, primeness and height of ideals, please refer to [1, 2, 6, 5].

References

- [1] H. Bordbar, I. Cristea, *Height of prime hyperideals in Krasner hyperrings*, **Filomat**, vol. 31(1) (1944), pp. 6153–6163, DOI: <http://dx.doi.org/10.2298/FIL1719153B>.
- [2] H. Bordbar, I. Cristea, M. Novak, *Height of hyperideals in Noetherian Krasner hyperrings*, **Scientific Bulletin – “Politehnica” University of Bucharest. Series A, Applied mathematics and physics.**, vol. 79(2) (2017), pp. 31–42.
- [3] H. Bordbar, H. Harizavi, Y. Jun, *Uni-soft ideals in coresiduated lattices*, **Sigma Journal of Engineering and Natural Sciences**, vol. 9(1) (2018), pp. 69–75.
- [4] H. Bordbar, Y. B. Jun, S. Z. Song, *Homomorphic Image and Inverse Image of Weak Closure Operations on Ideals of BCK-Algebras*, **Mathematics**, vol. 8(4) (2020), p. 576, DOI: <http://dx.doi.org/10.3390/math8040567>.
- [5] H. Bordbar, G. Muhiuddin, A. M. Alanazi, *Primeness of Relative Annihilators in BCK-Algebra*, **Symmetry**, vol. 12(2) (2020), p. 286, DOI: <http://dx.doi.org/10.3390/sym12020286>.

- [6] H. Bordbar, M. Novak, I. Cristea, *A note on the support of a hypermodule*, **Journal of Algebra and Its Applications**, vol. 19(01) (2020), p. 2050019, DOI: <http://dx.doi.org/10.1142/S021949882050019X>.
- [7] H. Bordbar, M. Takallo, R. Borzooei, Y. B. Jun, *BMBJ-neutrosophic subalgebra in BCK/BCI -algebras*, **Neutrosophic Sets and System**, vol. 27(1) (2020).
- [8] H. Bordbar, M. M. Zahedi, Y. B. Jun, *Relative annihilators in lower BCK -semilattices*, **Mathematical Sciences Letters**, vol. 6(2) (2017), pp. 1–7, DOI: <http://dx.doi.org/10.18576/msl/BZJ-20151219R1>.
- [9] Y. S. Huang, *BCI-algebra*, Beijing, Science Press, Cambridge (2006).
- [10] Y. B. Jun, *Neutrosophic subalgebras of several types in BCK/BCI -algebras*, **Annals of Fuzzy Mathematics and Informatics**, vol. 14(1) (2017), pp. 75–86.
- [11] Y. B. Jun, F. Smarandache, H. Bordbar, *Neutrosophic \mathcal{N} -structures applied to BCK/BCI -algebras*, **Informations**, vol. 8(1) (2017), p. 128, DOI: <http://dx.doi.org/10.3390/info8040128>.
- [12] Y. B. Jun, F. Smarandache, H. Bordbar, *Neutrosophic falling shadows applied to subalgebras and ideals in BCK/BCI -algebras*, **Annals of Fuzzy Mathematics and Informatics**, vol. 15(3) (2018).
- [13] Y. B. Jun, F. Smarandache, S. Z. Song, H. Bordbar, *Neutrosophic Permeable Values and Energetic Subsets with Applications in BCK/BCI -Algebras*, **Mathematics**, vol. 5(6) (2018), pp. 74–90, DOI: <http://dx.doi.org/10.3390/math6050074>.
- [14] Y. B. Jun, S. Z. Song, F. Smarandache, H. Bordbar, *Neutrosophic Quadruple BCK/BCI -Algebras*, **Axioms**, vol. 7(2) (2018), p. 41, DOI: <http://dx.doi.org/10.3390/axioms7020041>.
- [15] J. Meng, Y. B. Jun, *BCK -algebra*, Kyungmoon Sa Co., Seoul (1994).
- [16] G. Muhiuddin, A. N. Al-kenani, E. H. Roh, Y. B. Jun, *Implicative neutrosophic quadruple BCK -algebras and ideals*, **Symmetry**, vol. 11(2) (2019), p. 277, DOI: <http://dx.doi.org/10.3390/sym11020277>.
- [17] G. Muhiuddin, H. Bordbar, F. Smarandache, Y. B. Jun, *Further results on (\in, \in) -neutrosophic subalgebras and ideals in BCK/BCI -algebras*, **Neutrosophic Sets and Systems**, vol. 20 (2018), pp. 36–43.

- [18] G. Muhiuddin, Y. B. Jun, *Further results of neutrosophic subalgebras in BCK/BCI-algebras based on neutrosophic point*, **TWMS Journal of Applied and Engineering Mathematics**, vol. 10(2) (2020), pp. 232–240.
- [19] G. Muhiuddin, S. J. Kim, Y. B. Jun, *Implicative N -ideals of BCK-algebras based on neutrosophic N -structures*, **Discrete Mathematics, Algorithms and Applications**, vol. 11(1) (2019), p. 1950011, DOI: <http://dx.doi.org/10.1142/S1793830919500113>.
- [20] G. Muhiuddin, F. Smarandache, Y. B. Jun, *Neutrosophic quadruple ideals in neutrosophic quadruple BCI-algebras*, **Neutrosophic Sets and Systems**, vol. 25 (2019), pp. 161–173.
- [21] M. A. Öztürk, Y. B. Jun, *Neutrosophic ideals in BCK/BCI-algebras based on neutrosophic points*, **Journal of International Mathematical Virtual Institute**, vol. 8(1) (2018), pp. 1–17.
- [22] A. B. Saeid, Y. B. Jun, *Neutrosophic subalgebras of BCK/BCI-algebras based on neutrosophic points*, **Annals of Fuzzy Mathematics and Informatics**, vol. 14(2) (2017), pp. 87–97.
- [23] F. Smarandache, **A Unifying Field in Logics: Neutrosophic Logic. Neutrosophy, Neutrosophic Set, Neutrosophic Probability**, American Reserch Press, Rehoboth, N. M. (1999).
- [24] F. Smarandache, *Neutrosophic set-a generalization of the intuitionistic fuzzy set*, **International Journal of Pure and Applied Mathematics**, vol. 24(3) (2005), pp. 287–297.
- [25] S. Z. Song, F. Smarandache, Y. B. Jun, *Neutrosophic commutative \mathcal{N} -ideals in BCK-algebras*, **Informations**, vol. 8 (2017), p. 130, DOI: <http://dx.doi.org/10.3390/info8040130>.
- [26] M. M. Takalloand, H. Bordbar, R. A. Borzooei, Y. B. Jun, *BMBJ-neutrosophic ideals in BCK/BCI-algebras*, **Neutrosophic Sets and Systems**, vol. 1(27) (2019).

Hashem Bordbar

University of Nova Gorica

Center for Information Technologies and Applied Mathematics

Nova Gorica, Slovenia

e-mail: Hashem.bordbar@ung.si

A general model of neutrosophic ideals in BCK/BCI-algebras...

Rajab Ali Borzooei

Shahid Beheshti University
Department of Mathematics
Tehran, Iran

e-mail: borzooei@sbu.ac.ir

Florentin Smarandache

University of New Mexico
Mathematics & Science Department
705 Gurley Ave.
Gallup, NM 87301, USA

e-mail: fsmarandache@gmail.com

Young Bae Jun

Gyeongsang National University
Department of Mathematics Education
Jinju 52828, Korea

e-mail: skywine@gmail.com