

Article

# Commutative Generalized Neutrosophic Ideals in *BCK*-Algebras

Rajab Ali Borzooei <sup>1</sup> , Xiaohong Zhang <sup>2,3,\*</sup> , Florentin Smarandache <sup>4</sup>  and Young Bae Jun <sup>1,5</sup>

<sup>1</sup> Department of Mathematics, Shahid Beheshti University, Tehran 1983963113, Iran; borzooei@sbu.ac.ir (R.A.B.); skywine@gmail.com (Y.B.J.)

<sup>2</sup> Department of Mathematics, Shaanxi University of Science & Technology, Xi'an 710021, China

<sup>3</sup> Department of Mathematics, Shanghai Maritime University, Shanghai 201306, China

<sup>4</sup> Department of Mathematics, University of New Mexico, Gallup, NM 87301, USA; fsmarandache@gmail.com

<sup>5</sup> Department of Mathematics Education, Gyeongsang National University, Jinju 52828, Korea

\* Correspondence: zhangxh@shmtu.edu.cn

Received: 19 July 2018; Accepted: 17 August 2018; Published: 20 August 2018



**Abstract:** The concept of a commutative generalized neutrosophic ideal in a *BCK*-algebra is proposed, and related properties are proved. Characterizations of a commutative generalized neutrosophic ideal are considered. Also, some equivalence relations on the family of all commutative generalized neutrosophic ideals in *BCK*-algebras are introduced, and some properties are investigated.

**Keywords:** (commutative) ideal; generalized neutrosophic set; generalized neutrosophic ideal; commutative generalized neutrosophic ideal

## 1. Introduction

In 1965, Zadeh introduced the concept of fuzzy set in which the degree of membership is expressed by one function (that is, truth or  $t$ ). The theory of fuzzy set is applied to many fields, including fuzzy logic algebra systems (such as pseudo-*BCI*-algebras by Zhang [1]). In 1986, Atanassov introduced the concept of intuitionistic fuzzy set in which there are two functions, membership function ( $t$ ) and nonmembership function ( $f$ ). In 1995, Smarandache introduced the new concept of neutrosophic set in which there are three functions, membership function ( $t$ ), nonmembership function ( $f$ ) and indeterminacy/neutral membership function ( $i$ ), that is, there are three components ( $t, i, f$ ) = (truth, indeterminacy, falsehood) and they are independent components.

Neutrosophic algebraic structures in *BCK/BCI*-algebras are discussed in the papers [2–10]. Moreover, Zhang et al. studied totally dependent-neutrosophic sets, neutrosophic duplet semi-group and cancellable neutrosophic triplet groups (see [11,12]). Song et al. proposed the notion of generalized neutrosophic set and applied it to *BCK/BCI*-algebras.

In this paper, we propose the notion of a commutative generalized neutrosophic ideal in a *BCK*-algebra, and investigate related properties. We consider characterizations of a commutative generalized neutrosophic ideal. Using a collection of commutative ideals in *BCK*-algebras, we obtain a commutative generalized neutrosophic ideal. We also establish some equivalence relations on the family of all commutative generalized neutrosophic ideals in *BCK*-algebras, and discuss related basic properties of these ideals.

## 2. Preliminaries

A set  $X$  with a constant element  $0$  and a binary operation  $*$  is called a *BCI*-algebra, if it satisfies  $(\forall x, y, z \in X)$ :

- (I)  $((x * y) * (x * z)) * (z * y) = 0,$   
 (II)  $(x * (x * y)) * y = 0,$   
 (III)  $x * x = 0,$   
 (IV)  $x * y = 0, y * x = 0 \Rightarrow x = y.$

A BCI-algebra  $X$  is called a BCK-algebra, if it satisfies  $(\forall x \in X)$ :

- (V)  $0 * x = 0,$

For any BCK/BCI-algebra  $X$ , the following conditions hold  $(\forall x, y, z \in X)$ :

$$x * 0 = x, \quad (1)$$

$$x \leq y \Rightarrow x * z \leq y * z, z * y \leq z * x, \quad (2)$$

$$(x * y) * z = (x * z) * y, \quad (3)$$

$$(x * z) * (y * z) \leq x * y \quad (4)$$

where the relation  $\leq$  is defined by:  $x \leq y \iff x * y = 0$ . If the following assertion is valid for a BCK-algebra  $X, \forall x, y \in X,$

$$x * (x * y) = y * (y * x). \quad (5)$$

then  $X$  is called a commutative BCK-algebra.

Assume  $I$  is a subset of a BCK/BCI-algebra  $X$ . If the following conditions are valid, then we call  $I$  is an ideal of  $X$ :

$$0 \in I, \quad (6)$$

$$(\forall x \in X) (\forall y \in I) (x * y \in I \Rightarrow x \in I). \quad (7)$$

A subset  $I$  of a BCK-algebra  $X$  is called a commutative ideal of  $X$  if it satisfies (6) and

$$(\forall x, y, z \in X) ((x * y) * z \in I, z \in I \Rightarrow x * (y * (y * x)) \in I). \quad (8)$$

Recall that any commutative ideal is an ideal, but the inverse is not true in general (see [7]).

**Lemma 1** ([7]). *Let  $I$  be an ideal of a BCK-algebra  $X$ . Then  $I$  is commutative ideal of  $X$  if and only if it satisfies the following condition for all  $x, y$  in  $X$ :*

$$x * y \in I \Rightarrow x * (y * (y * x)) \in I. \quad (9)$$

For further information regarding BCK/BCI-algebras, please see the books [7,13].

Let  $X$  be a nonempty set. A fuzzy set in  $X$  is a function  $\mu : X \rightarrow [0, 1]$ , and the complement of  $\mu$ , denoted by  $\mu^c$ , is defined by  $\mu^c(x) = 1 - \mu(x), \forall x \in X$ . A fuzzy set  $\mu$  in a BCK/BCI-algebra  $X$  is called a fuzzy ideal of  $X$  if

$$(\forall x \in X) (\mu(0) \geq \mu(x)), \quad (10)$$

$$(\forall x, y \in X) (\mu(x) \geq \min\{\mu(x * y), \mu(y)\}). \quad (11)$$

Assume that  $X$  is a non-empty set. A neutrosophic set (NS) in  $X$  (see [14]) is a structure of the form:

$$A := \{\langle x; A_T(x), A_I(x), A_F(x) \rangle \mid x \in X\}$$

where  $A_T : X \rightarrow [0, 1]$ ,  $A_I : X \rightarrow [0, 1]$ , and  $A_F : X \rightarrow [0, 1]$ . We shall use the symbol  $A = (A_T, A_I, A_F)$  for the neutrosophic set

$$A := \{ \langle x; A_T(x), A_I(x), A_F(x) \rangle \mid x \in X \}.$$

A generalized neutrosophic set (GNS) in a non-empty set  $X$  is a structure of the form (see [15]):

$$A := \{ \langle x; A_T(x), A_{IT}(x), A_{IF}(x), A_F(x) \rangle \mid x \in X, A_{IT}(x) + A_{IF}(x) \leq 1 \}$$

where  $A_T : X \rightarrow [0, 1]$ ,  $A_F : X \rightarrow [0, 1]$ ,  $A_{IT} : X \rightarrow [0, 1]$ , and  $A_{IF} : X \rightarrow [0, 1]$ .

We shall use the symbol  $A = (A_T, A_{IT}, A_{IF}, A_F)$  for the generalized neutrosophic set

$$A := \{ \langle x; A_T(x), A_{IT}(x), A_{IF}(x), A_F(x) \rangle \mid x \in X, A_{IT}(x) + A_{IF}(x) \leq 1 \}.$$

Note that, for every GNS  $A = (A_T, A_{IT}, A_{IF}, A_F)$  in  $X$ , we have (for all  $x$  in  $X$ )

$$(\forall x \in X) (0 \leq A_T(x) + A_{IT}(x) + A_{IF}(x) + A_F(x) \leq 3).$$

If  $A = (A_T, A_{IT}, A_{IF}, A_F)$  is a GNS in  $X$ , then  $\square A = (A_T, A_{IT}, A_{IT}^c, A_T^c)$  and  $\diamond A = (A_F^c, A_{IF}^c, A_{IF}, A_F)$  are also GNSs in  $X$ .

Given a GNS  $A = (A_T, A_{IT}, A_{IF}, A_F)$  in a BCK/BCI-algebra  $X$  and  $\alpha_T, \alpha_{IT}, \beta_F, \beta_{IF} \in [0, 1]$ , we define four sets as follows:

$$\begin{aligned} U_A(T, \alpha_T) &:= \{x \in X \mid A_T(x) \geq \alpha_T\}, \\ U_A(IT, \alpha_{IT}) &:= \{x \in X \mid A_{IT}(x) \geq \alpha_{IT}\}, \\ L_A(F, \beta_F) &:= \{x \in X \mid A_F(x) \leq \beta_F\}, \\ L_A(IF, \beta_{IF}) &:= \{x \in X \mid A_{IF}(x) \leq \beta_{IF}\}. \end{aligned}$$

A GNS  $A = (A_T, A_{IT}, A_{IF}, A_F)$  in a BCK/BCI-algebra  $X$  is called a generalized neutrosophic ideal of  $X$  (see [15]) if

$$(\forall x \in X) \begin{pmatrix} A_T(0) \geq A_T(x), A_{IT}(0) \geq A_{IT}(x) \\ A_{IF}(0) \leq A_{IF}(x), A_F(0) \leq A_F(x) \end{pmatrix}, \tag{12}$$

$$(\forall x, y \in X) \begin{pmatrix} A_T(x) \geq \min\{A_T(x * y), A_T(y)\} \\ A_{IT}(x) \geq \min\{A_{IT}(x * y), A_{IT}(y)\} \\ A_{IF}(x) \leq \max\{A_{IF}(x * y), A_{IF}(y)\} \\ A_F(x) \leq \max\{A_F(x * y), A_F(y)\} \end{pmatrix}. \tag{13}$$

### 3. Commutative Generalized Neutrosophic Ideals

Unless specified,  $X$  will always represent a BCK-algebra in the following discussion.

**Definition 1.** A GNS  $A = (A_T, A_{IT}, A_{IF}, A_F)$  in  $X$  is called a commutative generalized neutrosophic ideal of  $X$  if it satisfies the condition (12) and

$$(\forall x, y, z \in X) \begin{pmatrix} A_T(x * (y * (y * x))) \geq \min\{A_T((x * y) * z), A_T(z)\} \\ A_{IT}(x * (y * (y * x))) \geq \min\{A_{IT}((x * y) * z), A_{IT}(z)\} \\ A_{IF}(x * (y * (y * x))) \leq \max\{A_{IF}((x * y) * z), A_{IF}(z)\} \\ A_F(x * (y * (y * x))) \leq \max\{A_F((x * y) * z), A_F(z)\} \end{pmatrix}. \tag{14}$$

**Example 1.** Denote  $X = \{0, a, b, c\}$ . The binary operation  $*$  on  $X$  is defined in Table 1.

**Table 1.** The operation “\*”.

*	0	a	b	c
0	0	0	0	0
a	a	0	0	a
b	b	a	0	b
c	c	c	c	0

We can verify that  $(X, *, 0)$  is a BCK-algebra (see [7]). Define a GNS  $A = (A_T, A_{IT}, A_{IF}, A_F)$  in  $X$  by Table 2.

**Table 2.** GNS  $A = (A_T, A_{IT}, A_{IF}, A_F)$ .

X	$A_T(x)$	$A_{IT}(x)$	$A_{IF}(x)$	$A_F(x)$
0	0.7	0.6	0.1	0.3
a	0.5	0.5	0.2	0.4
b	0.3	0.2	0.4	0.6
c	0.3	0.2	0.4	0.6

Then  $A = (A_T, A_{IT}, A_{IF}, A_F)$  is a commutative generalized neutrosophic ideal of  $X$ .

**Theorem 1.** Every commutative generalized neutrosophic ideal is a generalized neutrosophic ideal.

**Proof.** Assume that  $A = (A_T, A_{IT}, A_{IF}, A_F)$  is a commutative generalized neutrosophic ideal of  $X$ .  $\forall x, z \in X$ , we have

$$A_T(x) = A_T(x * (0 * (0 * x))) \geq \min\{A_T((x * 0) * z), A_T(z)\} = \min\{A_T(x * z), A_T(z)\},$$

$$A_{IT}(x) = A_{IT}(x * (0 * (0 * x))) \geq \min\{A_{IT}((x * 0) * z), A_{IT}(z)\} = \min\{A_{IT}(x * z), A_{IT}(z)\},$$

$$A_{IF}(x) = A_{IF}(x * (0 * (0 * x))) \leq \max\{A_{IF}((x * 0) * z), A_{IF}(z)\} = \max\{A_{IF}(x * z), A_{IF}(z)\},$$

and

$$A_F(x) = A_F(x * (0 * (0 * x))) \leq \max\{A_F((x * 0) * z), A_F(z)\} = \max\{A_F(x * z), A_F(z)\}.$$

Therefore  $A = (A_T, A_{IT}, A_{IF}, A_F)$  is a generalized neutrosophic ideal.  $\square$

The following example shows that the inverse of Theorem 1 is not true.

**Example 2.** Let  $X = \{0, 1, 2, 3, 4\}$  be a set with the binary operation \* which is defined in Table 3.

**Table 3.** The operation “\*”.

*	0	1	2	3	4
0	0	0	0	0	0
1	1	0	1	0	0
2	2	2	0	0	0
3	3	3	3	0	0
4	4	4	4	3	0

We can verify that  $(X, *, 0)$  is a BCK-algebra (see [7]). We define a GNS  $A = (A_T, A_{IT}, A_{IF}, A_F)$  in  $X$  by Table 4.

**Table 4.** GNS  $A = (A_T, A_{IT}, A_{IF}, A_F)$ .

$X$	$A_T(x)$	$A_{IT}(x)$	$A_{IF}(x)$	$A_F(x)$
0	0.7	0.6	0.1	0.3
1	0.5	0.4	0.2	0.6
2	0.3	0.5	0.4	0.4
3	0.3	0.4	0.4	0.6
4	0.3	0.4	0.4	0.6

It is routine to verify that  $A = (A_T, A_{IT}, A_{IF}, A_F)$  is a generalized neutrosophic ideal of  $X$ , but  $A$  is not a commutative generalized neutrosophic ideal of  $X$  since

$$A_T(2 * (3 * (3 * 2))) = A_T(2) = 0.3 \not\geq \min\{A_T((2 * 3) * 0), A_T(0)\}$$

and/or

$$A_{IF}(2 * (3 * (3 * 2))) = A_{IF}(2) = 0.4 \not\leq \max\{A_{IF}((2 * 3) * 0), A_{IF}(0)\}.$$

**Theorem 2.** Suppose that  $A = (A_T, A_{IT}, A_{IF}, A_F)$  is a generalized neutrosophic ideal of  $X$ . Then  $A = (A_T, A_{IT}, A_{IF}, A_F)$  is commutative if and only if it satisfies the following condition.

$$(\forall x, y \in X) \left( \begin{array}{l} A_T(x * y) \leq A_T(x * (y * (y * x))) \\ A_{IT}(x * y) \leq A_{IT}(x * (y * (y * x))) \\ A_{IF}(x * y) \geq A_{IF}(x * (y * (y * x))) \\ A_F(x * y) \geq A_F(x * (y * (y * x))) \end{array} \right). \tag{15}$$

**Proof.** Assume that  $A = (A_T, A_{IT}, A_{IF}, A_F)$  is a commutative generalized neutrosophic ideal of  $X$ . Taking  $z = 0$  in (14) and using (12) and (1) induces (15).

Conversely, let  $A = (A_T, A_{IT}, A_{IF}, A_F)$  be a generalized neutrosophic ideal of  $X$  satisfying the condition (15). Then

$$A_T(x * (y * (y * x))) \geq A_T(x * y) \geq \min\{A_T((x * y) * z), A_T(z)\},$$

$$A_{IT}(x * (y * (y * x))) \geq A_{IT}(x * y) \geq \min\{A_{IT}((x * y) * z), A_{IT}(z)\},$$

$$A_{IF}(x * (y * (y * x))) \leq A_{IF}(x * y) \leq \max\{A_{IF}((x * y) * z), A_{IF}(z)\}$$

and

$$A_F(x * (y * (y * x))) \leq A_F(x * y) \leq \max\{A_F((x * y) * z), A_F(z)\}$$

for all  $x, y, z \in X$ . Therefore  $A = (A_T, A_{IT}, A_{IF}, A_F)$  is a commutative generalized neutrosophic ideal of  $X$ .  $\square$

**Lemma 2 ([15]).** Any generalized neutrosophic ideal  $A = (A_T, A_{IT}, A_{IF}, A_F)$  of  $X$  satisfies:

$$(\forall x, y, z \in X) \left( x * y \leq z \Rightarrow \left\{ \begin{array}{l} A_T(x) \geq \min\{A_T(y), A_T(z)\} \\ A_{IT}(x) \geq \min\{A_{IT}(y), A_{IT}(z)\} \\ A_{IF}(x) \leq \max\{A_{IF}(y), A_{IF}(z)\} \\ A_F(x) \leq \max\{A_F(y), A_F(z)\} \end{array} \right. \right). \tag{16}$$

We provide a condition for a generalized neutrosophic ideal to be commutative.

**Theorem 3.** For any commutative BCK-algebra, every generalized neutrosophic ideal is commutative.

**Proof.** Assume that  $A = (A_T, A_{IT}, A_{IF}, A_F)$  is a generalized neutrosophic ideal of a commutative BCK-algebra  $X$ . Note that

$$\begin{aligned} ((x * (y * (y * x))) * ((x * y) * z)) * z &= ((x * (y * (y * x))) * z) * ((x * y) * z) \\ &\leq (x * (y * (y * x))) * (x * y) \\ &= (x * (x * y)) * (y * (y * x)) = 0, \end{aligned}$$

thus,  $(x * (y * (y * x))) * ((x * y) * z) \leq z, \forall x, y, z \in X$ . By Lemma 2 we get

$$\begin{aligned} A_T(x * (y * (y * x))) &\geq \min\{A_T((x * y) * z), A_T(z)\}, \\ A_{IT}(x * (y * (y * x))) &\geq \min\{A_{IT}((x * y) * z), A_{IT}(z)\}, \\ A_{IF}(x * (y * (y * x))) &\leq \max\{A_{IF}((x * y) * z), A_{IF}(z)\}, \\ A_F(x * (y * (y * x))) &\leq \max\{A_F((x * y) * z), A_F(z)\}. \end{aligned}$$

Therefore  $A = (A_T, A_{IT}, A_{IF}, A_F)$  is a commutative generalized neutrosophic ideal of  $X$ .  $\square$

**Lemma 3 ([15]).** If a GNS  $A = (A_T, A_{IT}, A_{IF}, A_F)$  in  $X$  is a generalized neutrosophic ideal of  $X$ , then the sets  $U_A(T, \alpha_T)$ ,  $U_A(IT, \alpha_{IT})$ ,  $L_A(F, \beta_F)$  and  $L_A(IF, \beta_{IF})$  are ideals of  $X$  for all  $\alpha_T, \alpha_{IT}, \beta_F, \beta_{IF} \in [0, 1]$  whenever they are non-empty.

**Theorem 4.** If a GNS  $A = (A_T, A_{IT}, A_{IF}, A_F)$  in  $X$  is a commutative generalized neutrosophic ideal of  $X$ , then the sets  $U_A(T, \alpha_T)$ ,  $U_A(IT, \alpha_{IT})$ ,  $L_A(F, \beta_F)$  and  $L_A(IF, \beta_{IF})$  are commutative ideals of  $X$  for all  $\alpha_T, \alpha_{IT}, \beta_F, \beta_{IF} \in [0, 1]$  whenever they are non-empty.

The commutative ideals  $U_A(T, \alpha_T)$ ,  $U_A(IT, \alpha_{IT})$ ,  $L_A(F, \beta_F)$  and  $L_A(IF, \beta_{IF})$  are called *level neutrosophic commutative ideals* of  $A = (A_T, A_{IT}, A_{IF}, A_F)$ .

**Proof.** Assume that  $A = (A_T, A_{IT}, A_{IF}, A_F)$  is a commutative generalized neutrosophic ideal of  $X$ . Then  $A = (A_T, A_{IT}, A_{IF}, A_F)$  is a generalized neutrosophic ideal of  $X$ . Thus  $U_A(T, \alpha_T)$ ,  $U_A(IT, \alpha_{IT})$ ,  $L_A(F, \beta_F)$  and  $L_A(IF, \beta_{IF})$  are ideals of  $X$  whenever they are non-empty applying Lemma 3. Suppose that  $x, y \in X$  and  $x * y \in U_A(T, \alpha_T) \cap U_A(IT, \alpha_{IT})$ . Using (15),

$$\begin{aligned} A_T(x * (y * (y * x))) &\geq A_T(x * y) \geq \alpha_T, \\ A_{IT}(x * (y * (y * x))) &\geq A_{IT}(x * y) \geq \alpha_{IT}, \end{aligned}$$

and so  $x * (y * (y * x)) \in U_A(T, \alpha_T)$  and  $x * (y * (y * x)) \in U_A(IT, \alpha_{IT})$ . Suppose that  $a, b \in X$  and  $a * b \in L_A(IF, \beta_{IF}) \cap L_A(F, \beta_F)$ . It follows from (15) that  $A_{IF}(a * (b * (b * a))) \leq A_{IF}(a * b) \leq \beta_{IF}$  and  $A_F(a * (b * (b * a))) \leq A_F(a * b) \leq \beta_F$ . Hence  $a * (b * (b * a)) \in L_A(IF, \beta_{IF})$  and  $a * (b * (b * a)) \in L_A(F, \beta_F)$ . Therefore  $U_A(T, \alpha_T)$ ,  $U_A(IT, \alpha_{IT})$ ,  $L_A(F, \beta_F)$  and  $L_A(IF, \beta_{IF})$  are commutative ideals of  $X$ .  $\square$

**Lemma 4 ([15]).** Assume that  $A = (A_T, A_{IT}, A_{IF}, A_F)$  is a GNS in  $X$  and  $U_A(T, \alpha_T)$ ,  $U_A(IT, \alpha_{IT})$ ,  $L_A(F, \beta_F)$  and  $L_A(IF, \beta_{IF})$  are ideals of  $X$ ,  $\forall \alpha_T, \alpha_{IT}, \beta_F, \beta_{IF} \in [0, 1]$ . Then  $A = (A_T, A_{IT}, A_{IF}, A_F)$  is a generalized neutrosophic ideal of  $X$ .

**Theorem 5.** Let  $A = (A_T, A_{IT}, A_{IF}, A_F)$  be a GNS in  $X$  such that  $U_A(T, \alpha_T)$ ,  $U_A(IT, \alpha_{IT})$ ,  $L_A(F, \beta_F)$  and  $L_A(IF, \beta_{IF})$  are commutative ideals of  $X$  for all  $\alpha_T, \alpha_{IT}, \beta_F, \beta_{IF} \in [0, 1]$ . Then  $A = (A_T, A_{IT}, A_{IF}, A_F)$  is a commutative generalized neutrosophic ideal of  $X$ .

**Proof.** Let  $\alpha_T, \alpha_{IT}, \beta_F, \beta_{IF} \in [0, 1]$  be such that the non-empty sets  $U_A(T, \alpha_T), U_A(IT, \alpha_{IT}), L_A(F, \beta_F)$  and  $L_A(IF, \beta_{IF})$  are commutative ideals of  $X$ . Then  $U_A(T, \alpha_T), U_A(IT, \alpha_{IT}), L_A(F, \beta_F)$  and  $L_A(IF, \beta_{IF})$  are ideals of  $X$ . Hence  $A = (A_T, A_{IT}, A_{IF}, A_F)$  is a generalized neutrosophic ideal of  $X$  applying Lemma 4. For any  $x, y \in X$ , let  $A_T(x * y) = \alpha_T$ . Then  $x * y \in U_A(T, \alpha_T)$ , and so  $x * (y * (y * x)) \in U_A(T, \alpha_T)$  by (9). Hence  $A_T(x * (y * (y * x))) \geq \alpha_T = A_T(x * y)$ . Similarly, we can show that

$$(\forall x, y \in X)(A_{IT}(x * (y * (y * x))) \geq A_{IT}(x * y)).$$

For any  $x, y, a, b \in X$ , let  $A_F(x * y) = \beta_F$  and  $A_{IF}(a * b) = \beta_{IF}$ . Then  $x * y \in L_A(F, \beta_F)$  and  $a * b \in L_A(IF, \beta_{IF})$ . Using Lemma 1 we have  $x * (y * (y * x)) \in L_A(F, \beta_F)$  and  $a * (b * (b * a)) \in L_A(IF, \beta_{IF})$ . Thus  $A_F(x * y) = \beta_F \geq A_F(x * (y * (y * x)))$  and  $A_{IF}(a * b) = \beta_{IF} \geq A_{IF}((a * b) * b)$ . Therefore  $A = (A_T, A_{IT}, A_{IF}, A_F)$  is a commutative generalized neutrosophic ideal of  $X$ .  $\square$

**Theorem 6.** Every commutative generalized neutrosophic ideal can be realized as level neutrosophic commutative ideals of some commutative generalized neutrosophic ideal of  $X$ .

**Proof.** Given a commutative ideal  $C$  of  $X$ , define a GNS  $A = (A_T, A_{IT}, A_{IF}, A_F)$  as follows

$$A_T(x) = \begin{cases} \alpha_T & \text{if } x \in C, \\ 0 & \text{otherwise,} \end{cases} \quad A_{IT}(x) = \begin{cases} \alpha_{IT} & \text{if } x \in C, \\ 0 & \text{otherwise,} \end{cases}$$

$$A_{IF}(x) = \begin{cases} \beta_{IF} & \text{if } x \in C, \\ 1 & \text{otherwise,} \end{cases} \quad A_F(x) = \begin{cases} \beta_F & \text{if } x \in C, \\ 1 & \text{otherwise,} \end{cases}$$

where  $\alpha_T, \alpha_{IT} \in (0, 1]$  and  $\beta_F, \beta_{IF} \in [0, 1)$ . Let  $x, y, z \in X$ . If  $(x * y) * z \in C$  and  $z \in C$ , then  $x * (y * (y * x)) \in C$ . Thus

$$A_T(x * (y * (y * x))) = \alpha_T = \min\{A_T((x * y) * z), A_T(z)\},$$

$$A_{IT}(x * (y * (y * x))) = \alpha_{IT} = \min\{A_{IT}((x * y) * z), A_{IT}(z)\},$$

$$A_{IF}(x * (y * (y * x))) = \beta_{IF} = \max\{A_{IF}((x * y) * z), A_{IF}(z)\},$$

$$A_F(x * (y * (y * x))) = \beta_F = \max\{A_F((x * y) * z), A_F(z)\}.$$

Assume that  $(x * y) * z \notin C$  and  $z \notin C$ . Then  $A_T((x * y) * z) = 0, A_T(z) = 0, A_{IT}((x * y) * z) = 0, A_{IT}(z) = 0, A_{IF}((x * y) * z) = 1, A_{IF}(z) = 1,$  and  $A_F((x * y) * z) = 1, A_F(z) = 1$ . It follows that

$$A_T(x * (y * (y * x))) \geq \min\{A_T((x * y) * z), A_T(z)\},$$

$$A_{IT}(x * (y * (y * x))) \geq \min\{A_{IT}((x * y) * z), A_{IT}(z)\},$$

$$A_{IF}(x * (y * (y * x))) \leq \max\{A_{IF}((x * y) * z), A_{IF}(z)\},$$

$$A_F(x * (y * (y * x))) \leq \max\{A_F((x * y) * z), A_F(z)\}.$$

If exactly one of  $(x * y) * z$  and  $z$  belongs to  $C$ , then exactly one of  $A_T((x * y) * z)$  and  $A_T(z)$  is equal to 0; exactly one of  $A_{IT}((x * y) * z)$  and  $A_{IT}(z)$  is equal to 0; exactly one of  $A_F((x * y) * z)$  and  $A_F(z)$  is equal to 1 and exactly one of  $A_{IF}((x * y) * z)$  and  $A_{IF}(z)$  is equal to 1. Hence

$$A_T(x * (y * (y * x))) \geq \min\{A_T((x * y) * z), A_T(z)\},$$

$$A_{IT}(x * (y * (y * x))) \geq \min\{A_{IT}((x * y) * z), A_{IT}(z)\},$$

$$A_{IF}(x * (y * (y * x))) \leq \max\{A_{IF}((x * y) * z), A_{IF}(z)\},$$

$$A_F(x * (y * (y * x))) \leq \max\{A_F((x * y) * z), A_F(z)\}.$$

It is clear that  $A_T(0) \geq A_T(x), A_{IT}(0) \geq A_{IT}(x), A_{IF}(0) \leq A_{IF}(x)$  and  $A_F(0) \leq A_F(x)$  for all  $x \in X$ . Therefore  $A = (A_T, A_{IT}, A_{IF}, A_F)$  is a commutative generalized neutrosophic ideal of  $X$ .

Obviously,  $U_A(T, \alpha_T) = C$ ,  $U_A(IT, \alpha_{IT}) = C$ ,  $L_A(F, \beta_F) = C$  and  $L_A(IF, \beta_{IF}) = C$ . This completes the proof.  $\square$

**Theorem 7.** Let  $\{C_t \mid t \in \Lambda\}$  be a collection of commutative ideals of  $X$  such that

- (1)  $X = \bigcup_{t \in \Lambda} C_t$ ,
- (2)  $(\forall s, t \in \Lambda) (s > t \iff C_s \subset C_t)$

where  $\Lambda$  is any index set. Let  $A = (A_T, A_{IT}, A_{IF}, A_F)$  be a GNS in  $X$  given by

$$(\forall x \in X) \left( \begin{array}{l} A_T(x) = \sup\{t \in \Lambda \mid x \in C_t\} = A_{IT}(x) \\ A_{IF}(x) = \inf\{t \in \Lambda \mid x \in C_t\} = A_F(x) \end{array} \right). \tag{17}$$

Then  $A = (A_T, A_{IT}, A_{IF}, A_F)$  is a commutative generalized neutrosophic ideal of  $X$ .

**Proof.** According to Theorem 5, it is sufficient to show that  $U(T, t)$ ,  $U(IT, t)$ ,  $L(F, s)$  and  $L(IF, s)$  are commutative ideals of  $X$  for every  $t \in [0, A_T(0) = A_{IT}(0)]$  and  $s \in [A_{IF}(0) = A_F(0), 1]$ . In order to prove  $U(T, t)$  and  $U(IT, t)$  are commutative ideals of  $X$ , we consider two cases:

- (i)  $t = \sup\{q \in \Lambda \mid q < t\}$ ,
- (ii)  $t \neq \sup\{q \in \Lambda \mid q < t\}$ .

For the first case, we have

$$\begin{aligned} x \in U(T, t) &\iff (\forall q < t)(x \in C_q) \iff x \in \bigcap_{q < t} C_q, \\ x \in U(IT, t) &\iff (\forall q < t)(x \in C_q) \iff x \in \bigcap_{q < t} C_q. \end{aligned}$$

Hence  $U(T, t) = \bigcap_{q < t} C_q = U(IT, t)$ , and so  $U(T, t)$  and  $U(IT, t)$  are commutative ideals of  $X$ .

For the second case, we claim that  $U(T, t) = \bigcup_{q \geq t} C_q = U(IT, t)$ . If  $x \in \bigcup_{q \geq t} C_q$ , then  $x \in C_q$  for some  $q \geq t$ . It follows that  $A_{IT}(x) = A_T(x) \geq q \geq t$  and so that  $x \in U(T, t)$  and  $x \in U(IT, t)$ . This shows that  $\bigcup_{q \geq t} C_q \subseteq U(T, t)$  and  $\bigcup_{q \geq t} C_q \subseteq U(IT, t)$ . Now, suppose  $x \notin \bigcup_{q \geq t} C_q$ . Then  $x \notin C_q, \forall q \geq t$ . Since  $t \neq \sup\{q \in \Lambda \mid q < t\}$ , there exists  $\varepsilon > 0$  such that  $(t - \varepsilon, t) \cap \Lambda = \emptyset$ . Thus  $x \notin C_q, \forall q > t - \varepsilon$ , this means that if  $x \in C_q$ , then  $q \leq t - \varepsilon$ . So  $A_{IT}(x) = A_T(x) \leq t - \varepsilon < t$ , and so  $x \notin U(T, t) = U(IT, t)$ . Therefore  $U(T, t) = U(IT, t) \subseteq \bigcup_{q \geq t} C_q$ . Consequently,  $U(T, t) = U(IT, t) = \bigcup_{q \geq t} C_q$  which

is a commutative ideal of  $X$ . Next we show that  $L(F, s)$  and  $L(IF, s)$  are commutative ideals of  $X$ . We consider two cases as follows:

- (iii)  $s = \inf\{r \in \Lambda \mid s < r\}$ ,
- (iv)  $s \neq \inf\{r \in \Lambda \mid s < r\}$ .

Case (iii) implies that

$$\begin{aligned} x \in L(IF, s) &\iff (\forall s < r)(x \in C_r) \iff x \in \bigcap_{s < r} C_r, \\ x \in L(F, s) &\iff (\forall s < r)(x \in C_r) \iff x \in \bigcap_{s < r} C_r. \end{aligned}$$

It follows that  $L(IF, s) = L(F, s) = \bigcap_{s < r} C_r$ , which is a commutative ideal of  $X$ . Case (iv) induces  $(s, s + \varepsilon) \cap \Lambda = \emptyset$  for some  $\varepsilon > 0$ . If  $x \in \bigcup_{s \geq r} C_r$ , then  $x \in C_r$  for some  $r \leq s$ , and so  $A_{IF}(x) = A_F(x) \leq r \leq s$ , that is,  $x \in L(IF, s)$  and  $x \in L(F, s)$ . Hence  $\bigcup_{s \geq r} C_r \subseteq L(IF, s) = L(F, s)$ . If  $x \notin \bigcup_{s \geq r} C_r$ , then  $x \notin C_r$



for all  $r \leq s$  which implies that  $x \notin C_r$  for all  $r \leq s + \varepsilon$ , that is, if  $x \in C_r$  then  $r \geq s + \varepsilon$ . Hence  $A_{IF}(x) = A_F(x) \geq s + \varepsilon > s$ , and so  $x \notin L(A_{IF}, s) = L(A_F, s)$ . Hence  $L(A_{IF}, s) = L(A_F, s) = \bigcup_{s \geq r} C_r$  which is a commutative ideal of  $X$ . This completes the proof.  $\square$

Assume that  $f : X \rightarrow Y$  is a homomorphism of BCK/BCI-algebras ([7]). For any GNS  $A = (A_T, A_{IT}, A_{IF}, A_F)$  in  $Y$ , we define a new GNS  $A^f = (A_T^f, A_{IT}^f, A_{IF}^f, A_F^f)$  in  $X$ , which is called the *induced GNS*, by

$$(\forall x \in X) \left( \begin{array}{l} A_T^f(x) = A_T(f(x)), A_{IT}^f(x) = A_{IT}(f(x)) \\ A_{IF}^f(x) = A_{IF}(f(x)), A_F^f(x) = A_F(f(x)) \end{array} \right). \quad (18)$$

**Lemma 5** ([15]). *Let  $f : X \rightarrow Y$  be a homomorphism of BCK/BCI-algebras. If a GNS  $A = (A_T, A_{IT}, A_{IF}, A_F)$  in  $Y$  is a generalized neutrosophic ideal of  $Y$ , then the new GNS  $A^f = (A_T^f, A_{IT}^f, A_{IF}^f, A_F^f)$  in  $X$  is a generalized neutrosophic ideal of  $X$ .*

**Theorem 8.** *Let  $f : X \rightarrow Y$  be a homomorphism of BCK-algebras. If a GNS  $A = (A_T, A_{IT}, A_{IF}, A_F)$  in  $Y$  is a commutative generalized neutrosophic ideal of  $Y$ , then the new GNS  $A^f = (A_T^f, A_{IT}^f, A_{IF}^f, A_F^f)$  in  $X$  is a commutative generalized neutrosophic ideal of  $X$ .*

**Proof.** Suppose that  $A = (A_T, A_{IT}, A_{IF}, A_F)$  is a commutative generalized neutrosophic ideal of  $Y$ . Then  $A = (A_T, A_{IT}, A_{IF}, A_F)$  is a generalized neutrosophic ideal of  $Y$  by Theorem 1, and so  $A^f = (A_T^f, A_{IT}^f, A_{IF}^f, A_F^f)$  is a generalized neutrosophic ideal of  $Y$  by Lemma 5. For any  $x, y \in X$ , we have

$$\begin{aligned} A_T^f(x * (y * (y * x))) &= A_T(f(x * (y * (y * x)))) \\ &= A_T(f(x) * (f(y) * (f(y) * f(x)))) \\ &\geq A_T(f(x) * f(y)) \\ &= A_T(f(x * y)) = A_T^f(x * y), \end{aligned}$$

$$\begin{aligned} A_{IT}^f(x * (y * (y * x))) &= A_{IT}(f(x * (y * (y * x)))) \\ &= A_{IT}(f(x) * (f(y) * (f(y) * f(x)))) \\ &\geq A_{IT}(f(x) * f(y)) \\ &= A_{IT}(f(x * y)) = A_{IT}^f(x * y), \end{aligned}$$

$$\begin{aligned} A_{IF}^f(x * (y * (y * x))) &= A_{IF}(f(x * (y * (y * x)))) \\ &= A_{IF}(f(x) * (f(y) * (f(y) * f(x)))) \\ &\leq A_{IF}(f(x) * f(y)) \\ &= A_{IF}(f(x * y)) = A_{IF}^f(x * y), \end{aligned}$$

and

$$\begin{aligned} A_F^f(x * (y * (y * x))) &= A_F(f(x * (y * (y * x)))) \\ &= A_F(f(x) * (f(y) * (f(y) * f(x)))) \\ &\leq A_F(f(x) * f(y)) \\ &= A_F(f(x * y)) = A_F^f(x * y). \end{aligned}$$

Therefore  $A^f = (A_T^f, A_{IT}^f, A_{IF}^f, A_F^f)$  is a commutative generalized neutrosophic ideal of  $X$ .  $\square$

**Lemma 6** ([15]). *Let  $f : X \rightarrow Y$  be an onto homomorphism of BCK/BCI-algebras and let  $A = (A_T, A_{IT}, A_{IF}, A_F)$  be a GNS in  $Y$ . If the induced GNS  $A^f = (A_T^f, A_{IT}^f, A_{IF}^f, A_F^f)$  in  $X$  is a generalized neutrosophic ideal of  $X$ , then  $A = (A_T, A_{IT}, A_{IF}, A_F)$  is a generalized neutrosophic ideal of  $Y$ .*

**Theorem 9.** *Assume that  $f : X \rightarrow Y$  is an onto homomorphism of BCK-algebras and  $A = (A_T, A_{IT}, A_{IF}, A_F)$  is a GNS in  $Y$ . If the induced GNS  $A^f = (A_T^f, A_{IT}^f, A_{IF}^f, A_F^f)$  in  $X$  is a commutative generalized neutrosophic ideal of  $X$ , then  $A = (A_T, A_{IT}, A_{IF}, A_F)$  is a commutative generalized neutrosophic ideal of  $Y$ .*

**Proof.** Suppose that  $A^f = (A_T^f, A_{IT}^f, A_{IF}^f, A_F^f)$  is a commutative generalized neutrosophic ideal of  $X$ . Then  $A^f = (A_T^f, A_{IT}^f, A_{IF}^f, A_F^f)$  is a generalized neutrosophic ideal of  $X$ , and thus  $A = (A_T, A_{IT}, A_{IF}, A_F)$  is a generalized neutrosophic ideal of  $Y$ . For any  $a, b, c \in Y$ , there exist  $x, y, z \in X$  such that  $f(x) = a, f(y) = b$  and  $f(z) = c$ . Thus,

$$\begin{aligned} A_T(a * (b * (b * a))) &= A_T(f(x) * (f(y) * (f(y) * f(x)))) = A_T(f(x * (y * (y * x)))) \\ &= A_T^f(x * (y * (y * x))) \geq A_T^f(x * y) \\ &= A_T(f(x) * f(y)) = A_T(a * b), \end{aligned}$$

$$\begin{aligned} A_{IT}(a * (b * (b * a))) &= A_{IT}(f(x) * (f(y) * (f(y) * f(x)))) = A_{IT}(f(x * (y * (y * x)))) \\ &= A_{IT}^f(x * (y * (y * x))) \geq A_{IT}^f(x * y) \\ &= A_{IT}(f(x) * f(y)) = A_{IT}(a * b), \end{aligned}$$

$$\begin{aligned} A_{IF}(a * (b * (b * a))) &= A_{IF}(f(x) * (f(y) * (f(y) * f(x)))) = A_{IF}(f(x * (y * (y * x)))) \\ &= A_{IF}^f(x * (y * (y * x))) \leq A_{IF}^f(x * y) \\ &= A_{IF}(f(x) * f(y)) = A_{IF}(a * b), \end{aligned}$$

and

$$\begin{aligned} A_F(a * (b * (b * a))) &= A_F(f(x) * (f(y) * (f(y) * f(x)))) = A_F(f(x * (y * (y * x)))) \\ &= A_F^f(x * (y * (y * x))) \leq A_F^f(x * y) \\ &= A_F(f(x) * f(y)) = A_F(a * b). \end{aligned}$$

It follows from Theorem 2 that  $A = (A_T, A_{IT}, A_{IF}, A_F)$  is a commutative generalized neutrosophic ideal of  $Y$ .  $\square$

Let  $CGNI(X)$  denote the set of all commutative generalized neutrosophic ideals of  $X$  and  $t \in [0, 1]$ . Define binary relations  $U_T^t, U_{IT}^t, L_F^t$  and  $L_{IF}^t$  on  $CGNI(X)$  as follows:

$$\begin{aligned} (A, B) \in U_T^t &\Leftrightarrow U_A(T, t) = U_B(T, t), (A, B) \in U_{IT}^t \Leftrightarrow U_A(IT, t) = U_B(IT, t), \\ (A, B) \in L_F^t &\Leftrightarrow L_A(F, t) = L_B(F, t), (A, B) \in L_{IF}^t \Leftrightarrow L_A(IF, t) = L_B(IF, t) \end{aligned} \tag{19}$$

for  $A = (A_T, A_{IT}, A_{IF}, A_F)$  and  $B = (B_T, B_{IT}, B_{IF}, B_F)$  in  $CGNI(X)$ . Then clearly  $U_T^t, U_{IT}^t, L_F^t$  and  $L_{IF}^t$  are equivalence relations on  $CGNI(X)$ . For any  $A = (A_T, A_{IT}, A_{IF}, A_F) \in CGNI(X)$ , let  $[A]_{U_T^t}$  (resp.,  $[A]_{U_{IT}^t}, [A]_{L_F^t}$  and  $[A]_{L_{IF}^t}$ ) denote the equivalence class of  $A = (A_T, A_{IT}, A_{IF}, A_F)$  modulo  $U_T^t$  (resp.,  $U_{IT}^t, L_F^t$  and  $L_{IF}^t$ ). Denote by  $CGNI(X)/U_T^t$  (resp.,  $CGNI(X)/U_{IT}^t, CGNI(X)/L_F^t$  and  $CGNI(X)/L_{IF}^t$ ) the system of all equivalence classes modulo  $U_T^t$  (resp.,  $U_{IT}^t, L_F^t$  and  $L_{IF}^t$ ); so

$$CGNI(X)/U_T^t = \{[A]_{U_T^t} \mid A = (A_T, A_{IT}, A_{IF}, A_F) \in CGNI(X)\}, \tag{20}$$

$$CGNI(X)/U_{IT}^t = \{[A]_{U_{IT}^t} \mid A = (A_T, A_{IT}, A_{IF}, A_F) \in CGNI(X)\}, \quad (21)$$

$$CGNI(X)/L_F^t = \{[A]_{L_F^t} \mid A = (A_T, A_{IT}, A_{IF}, A_F) \in CGNI(X)\}, \quad (22)$$

and

$$CGNI(X)/L_{IF}^t = \{[A]_{L_{IF}^t} \mid A = (A_T, A_{IT}, A_{IF}, A_F) \in CGNI(X)\}, \quad (23)$$

respectively. Let  $CI(X)$  denote the family of all commutative ideals of  $X$  and let  $t \in [0, 1]$ . Define maps

$$f_t : CGNI(X) \rightarrow CI(X) \cup \{\emptyset\}, A \mapsto U_A(T, t), \quad (24)$$

$$g_t : CGNI(X) \rightarrow CI(X) \cup \{\emptyset\}, A \mapsto U_A(IT, t), \quad (25)$$

$$\alpha_t : CGNI(X) \rightarrow CI(X) \cup \{\emptyset\}, A \mapsto L_A(F, t), \quad (26)$$

and

$$\beta_t : CGNI(X) \rightarrow CI(X) \cup \{\emptyset\}, A \mapsto L_A(IF, t). \quad (27)$$

Then the definitions of  $f_t$ ,  $g_t$ ,  $\alpha_t$  and  $\beta_t$  are well.

**Theorem 10.** Suppose  $t \in (0, 1)$ , the definitions of  $f_t$ ,  $g_t$ ,  $\alpha_t$  and  $\beta_t$  are as above. Then the maps  $f_t$ ,  $g_t$ ,  $\alpha_t$  and  $\beta_t$  are surjective from  $CGNI(X)$  to  $CI(X) \cup \{\emptyset\}$ .

**Proof.** Assume  $t \in (0, 1)$ . We know that  $\mathbf{0}_{\sim} = (\mathbf{0}_T, \mathbf{0}_{IT}, \mathbf{1}_{IF}, \mathbf{1}_F)$  is in  $CGNI(X)$  where  $\mathbf{0}_T$ ,  $\mathbf{0}_{IT}$ ,  $\mathbf{1}_{IF}$  and  $\mathbf{1}_F$  are constant functions on  $X$  defined by  $\mathbf{0}_T(x) = 0$ ,  $\mathbf{0}_{IT}(x) = 0$ ,  $\mathbf{1}_{IF}(x) = 1$  and  $\mathbf{1}_F(x) = 1$  for all  $x \in X$ . Obviously  $f_t(\mathbf{0}_{\sim}) = U_{\mathbf{0}_{\sim}}(T, t)$ ,  $g_t(\mathbf{0}_{\sim}) = U_{\mathbf{0}_{\sim}}(IT, t)$ ,  $\alpha_t(\mathbf{0}_{\sim}) = L_{\mathbf{0}_{\sim}}(F, t)$  and  $\beta_t(\mathbf{0}_{\sim}) = L_{\mathbf{0}_{\sim}}(IF, t)$  are empty. Let  $G (\neq \emptyset) \in CGNI(X)$ , and consider functions:

$$G_T : X \rightarrow [0, 1], G \mapsto \begin{cases} 1 & \text{if } x \in G, \\ 0 & \text{otherwise,} \end{cases}$$

$$G_{IT} : X \rightarrow [0, 1], G \mapsto \begin{cases} 1 & \text{if } x \in G, \\ 0 & \text{otherwise,} \end{cases}$$

$$G_F : X \rightarrow [0, 1], G \mapsto \begin{cases} 0 & \text{if } x \in G, \\ 1 & \text{otherwise,} \end{cases}$$

and

$$G_{IF} : X \rightarrow [0, 1], G \mapsto \begin{cases} 0 & \text{if } x \in G, \\ 1 & \text{otherwise.} \end{cases}$$

Then  $G_{\sim} = (G_T, G_{IT}, G_{IF}, G_F)$  is a commutative generalized neutrosophic ideal of  $X$ , and  $f_t(G_{\sim}) = U_{G_{\sim}}(T, t) = G$ ,  $g_t(G_{\sim}) = U_{G_{\sim}}(IT, t) = G$ ,  $\alpha_t(G_{\sim}) = L_{G_{\sim}}(F, t) = G$  and  $\beta_t(G_{\sim}) = L_{G_{\sim}}(IF, t) = G$ . Therefore  $f_t$ ,  $g_t$ ,  $\alpha_t$  and  $\beta_t$  are surjective.  $\square$

**Theorem 11.** The quotient sets

$$CGNI(X)/U_T^t, CGNI(X)/U_{IT}^t, CGNI(X)/L_F^t \text{ and } CGNI(X)/L_{IF}^t$$

are equipotent to  $CI(X) \cup \{\emptyset\}$ .

**Proof.** For  $t \in (0,1)$ , let  $f_t^*$  (resp,  $g_t^*$ ,  $\alpha_t^*$  and  $\beta_t^*$ ) be a map from  $CGNI(X)/U_T^t$  (resp.,  $CGNI(X)/U_{IT}^t$ ,  $CGNI(X)/L_F^t$  and  $CGNI(X)/L_{IF}^t$ ) to  $CI(X) \cup \{\emptyset\}$  defined by  $f_t^*([A]_{U_T^t}) = f_t(A)$  (resp.,  $g_t^*([A]_{U_{IT}^t}) = g_t(A)$ ,  $\alpha_t^*([A]_{L_F^t}) = \alpha_t(A)$  and  $\beta_t^*([A]_{L_{IF}^t}) = \beta_t(A)$ ) for all  $A = (A_T, A_{IT}, A_{IF}, A_F) \in CGNI(X)$ . If  $U_A(T,t) = U_B(T,t)$ ,  $U_A(IT,t) = U_B(IT,t)$ ,  $L_A(F,t) = L_B(F,t)$  and  $L_A(IF,t) = L_B(IF,t)$  for  $A = (A_T, A_{IT}, A_{IF}, A_F)$  and  $B = (B_T, B_{IT}, B_{IF}, B_F)$  in  $CGNI(X)$ , then  $(A,B) \in U_T^t$ ,  $(A,B) \in U_{IT}^t$ ,  $(A,B) \in L_F^t$  and  $(A,B) \in L_{IF}^t$ . Hence  $[A]_{U_T^t} = [B]_{U_T^t}$ ,  $[A]_{U_{IT}^t} = [B]_{U_{IT}^t}$ ,  $[A]_{L_F^t} = [B]_{L_F^t}$  and  $[A]_{L_{IF}^t} = [B]_{L_{IF}^t}$ . Therefore  $f_t^*$  (resp,  $g_t^*$ ,  $\alpha_t^*$  and  $\beta_t^*$ ) is injective. Now let  $G(\neq \emptyset) \in CGNI(X)$ . For  $G_\sim = (G_T, G_{IT}, G_{IF}, G_F) \in CGNI(X)$ , we have

$$f_t^*([G_\sim]_{U_T^t}) = f_t(G_\sim) = U_{G_\sim}(T,t) = G,$$

$$g_t^*([G_\sim]_{U_{IT}^t}) = g_t(G_\sim) = U_{G_\sim}(IT,t) = G,$$

$$\alpha_t^*([G_\sim]_{L_F^t}) = \alpha_t(G_\sim) = L_{G_\sim}(F,t) = G$$

and

$$\beta_t^*([G_\sim]_{L_{IF}^t}) = \beta_t(G_\sim) = L_{G_\sim}(IF,t) = G.$$

Finally, for  $\mathbf{0}_\sim = (\mathbf{0}_T, \mathbf{0}_{IT}, \mathbf{1}_{IF}, \mathbf{1}_F) \in CGNI(X)$ , we have

$$f_t^*([\mathbf{0}_\sim]_{U_T^t}) = f_t(\mathbf{0}_\sim) = U_{\mathbf{0}_\sim}(T,t) = \emptyset,$$

$$g_t^*([\mathbf{0}_\sim]_{U_{IT}^t}) = g_t(\mathbf{0}_\sim) = U_{\mathbf{0}_\sim}(IT,t) = \emptyset,$$

$$\alpha_t^*([\mathbf{0}_\sim]_{L_F^t}) = \alpha_t(\mathbf{0}_\sim) = L_{\mathbf{0}_\sim}(F,t) = \emptyset$$

and

$$\beta_t^*([\mathbf{0}_\sim]_{L_{IF}^t}) = \beta_t(\mathbf{0}_\sim) = L_{\mathbf{0}_\sim}(IF,t) = \emptyset.$$

Therefore,  $f_t^*$  (resp,  $g_t^*$ ,  $\alpha_t^*$  and  $\beta_t^*$ ) is surjective.  $\square$

$\forall t \in [0,1]$ , define another relations  $R^t$  and  $Q^t$  on  $CGNI(X)$  as follows:

$$(A,B) \in R^t \Leftrightarrow U_A(T,t) \cap L_A(F,t) = U_B(T,t) \cap L_B(F,t)$$

and

$$(A,B) \in Q^t \Leftrightarrow U_A(IT,t) \cap L_A(IF,t) = U_B(IT,t) \cap L_B(IF,t)$$

for any  $A = (A_T, A_{IT}, A_{IF}, A_F)$  and  $B = (B_T, B_{IT}, B_{IF}, B_F)$  in  $CGNI(X)$ . Then  $R^t$  and  $Q^t$  are equivalence relations on  $CGNI(X)$ .

**Theorem 12.** Suppose  $t \in (0, 1)$ , consider the following maps

$$\varphi_t : CGNI(X) \rightarrow CI(X) \cup \{\emptyset\}, A \mapsto f_t(A) \cap \alpha_t(A), \tag{28}$$

and

$$\psi_t : CGNI(X) \rightarrow CI(X) \cup \{\emptyset\}, A \mapsto g_t(A) \cap \beta_t(A) \tag{29}$$

for each  $A = (A_T, A_{IT}, A_{IF}, A_F) \in CGNI(X)$ . Then  $\varphi_t$  and  $\psi_t$  are surjective.

**Proof.** Assume  $t \in (0, 1)$ . For  $\mathbf{0}_{\sim} = (\mathbf{0}_T, \mathbf{0}_{IT}, \mathbf{1}_{IF}, \mathbf{1}_F) \in CGNI(X)$ ,

$$\varphi_t(\mathbf{0}_{\sim}) = f_t(\mathbf{0}_{\sim}) \cap \alpha_t(\mathbf{0}_{\sim}) = U_{\mathbf{0}_{\sim}}(T, t) \cap L_{\mathbf{0}_{\sim}}(F, t) = \emptyset$$

and

$$\psi_t(\mathbf{0}_{\sim}) = g_t(\mathbf{0}_{\sim}) \cap \beta_t(\mathbf{0}_{\sim}) = U_{\mathbf{0}_{\sim}}(IT, t) \cap L_{\mathbf{0}_{\sim}}(IF, t) = \emptyset.$$

For any  $G \in CI(X)$ , there exists  $G_{\sim} = (G_T, G_{IT}, G_{IF}, G_F) \in CGNI(X)$  such that

$$\varphi_t(G_{\sim}) = f_t(G_{\sim}) \cap \alpha_t(G_{\sim}) = U_{G_{\sim}}(T, t) \cap L_{G_{\sim}}(F, t) = G$$

and

$$\psi_t(G_{\sim}) = g_t(G_{\sim}) \cap \beta_t(G_{\sim}) = U_{G_{\sim}}(IT, t) \cap L_{G_{\sim}}(IF, t) = G.$$

Therefore  $\varphi_t$  and  $\psi_t$  are surjective.  $\square$

**Theorem 13.** For any  $t \in (0, 1)$ , the quotient sets  $CGNI(X)/R^t$  and  $CGNI(X)/Q^t$  are equipotent to  $CI(X) \cup \{\emptyset\}$ .

**Proof.** Let  $t \in (0, 1)$  and define maps

$$\varphi_t^* : CGNI(X)/R^t \rightarrow CI(X) \cup \{\emptyset\}, [A]_{R^t} \mapsto \varphi_t(A)$$

and

$$\psi_t^* : CGNI(X)/Q^t \rightarrow CI(X) \cup \{\emptyset\}, [A]_{Q^t} \mapsto \psi_t(A).$$

If  $\varphi_t^*([A]_{R^t}) = \varphi_t^*([B]_{R^t})$  and  $\psi_t^*([A]_{Q^t}) = \psi_t^*([B]_{Q^t})$  for all  $[A]_{R^t}, [B]_{R^t} \in CGNI(X)/R^t$  and  $[A]_{Q^t}, [B]_{Q^t} \in CGNI(X)/Q^t$ , then  $f_t(A) \cap \alpha_t(A) = f_t(B) \cap \alpha_t(B)$  and  $g_t(A) \cap \beta_t(A) = g_t(B) \cap \beta_t(B)$ , that is,  $U_A(T, t) \cap L_A(F, t) = U_B(T, t) \cap L_B(F, t)$  and  $U_A(IT, t) \cap L_A(IF, t) = U_B(IT, t) \cap L_B(IF, t)$ . Hence  $(A, B) \in R^t, (A, B) \in Q^t$ . So  $[A]_{R^t} = [B]_{R^t}, [A]_{Q^t} = [B]_{Q^t}$ , which shows that  $\varphi_t^*$  and  $\psi_t^*$  are injective. For  $\mathbf{0}_{\sim} = (\mathbf{0}_T, \mathbf{0}_{IT}, \mathbf{1}_{IF}, \mathbf{1}_F) \in CGNI(X)$ ,

$$\varphi_t^*([\mathbf{0}_{\sim}]_{R^t}) = \varphi_t(\mathbf{0}_{\sim}) = f_t(\mathbf{0}_{\sim}) \cap \alpha_t(\mathbf{0}_{\sim}) = U_{\mathbf{0}_{\sim}}(\mathbf{0}_T, t) \cap L_{\mathbf{0}_{\sim}}(\mathbf{1}_F, t) = \emptyset$$

and

$$\psi_t^*([\mathbf{0}_{\sim}]_{Q^t}) = \psi_t(\mathbf{0}_{\sim}) = g_t(\mathbf{0}_{\sim}) \cap \beta_t(\mathbf{0}_{\sim}) = U_{\mathbf{0}_{\sim}}(\mathbf{0}_{IT}, t) \cap L_{\mathbf{0}_{\sim}}(\mathbf{1}_{IF}, t) = \emptyset.$$

If  $G \in CI(X)$ , then  $G_{\sim} = (G_T, G_{IT}, G_{IF}, G_F) \in CGNI(X)$ , and so

$$\varphi_t^*([G_{\sim}]_{R^t}) = \varphi_t(G_{\sim}) = f_t(G_{\sim}) \cap \alpha_t(G_{\sim}) = U_{G_{\sim}}(G_T, t) \cap L_{G_{\sim}}(G_F, t) = G$$

and

$$\psi_t^* \left( [G_{\sim}]_{Q_t} \right) = \psi_t(G_{\sim}) = g_t(G_{\sim}) \cap \beta_t(G_{\sim}) = U_{G_{\sim}}(G_{IT}, t) \cap L_{G_{\sim}}(G_{IF}, t) = G.$$

Hence  $\varphi_t^*$  and  $\psi_t^*$  are surjective, and the proof is complete.  $\square$

#### 4. Conclusions

Based on the theory of generalized neutrosophic sets, we proposed the new concept of commutative generalized neutrosophic ideal in a BCK-algebra, and obtained some characterizations. Moreover, we investigated some homomorphism properties related to commutative generalized neutrosophic ideals.

The research ideas of this paper can be extended to a wide range of logical algebraic systems such as pseudo-BCI algebras (see [1,16]). At the same time, the concept of generalized neutrosophic set involved in this paper can be further studied according to the thought in [11,17], which will be the direction of our next research work.

**Author Contributions:** R.A.B. and Y.B.J. wrote the draft version of the paper, X.Z. wrote the revised version and completed submission of the paper, F.S. as a supervisor gave some guidance.

**Funding:** This research was funded by the National Natural Science Foundation of China grant number 61573240.

**Conflicts of Interest:** The authors declare no conflicts of interest.

#### References

- Zhang, X.H. Fuzzy anti-grouped filters and fuzzy normal filters in pseudo-BCI algebras. *J. Intell. Fuzzy Syst.* **2017**, *33*, 1767–1774. [[CrossRef](#)]
- Jun, Y.B. Neutrosophic subalgebras of several types in BCK/BCI-algebras. *Ann. Fuzzy Math. Inform.* **2017**, *14*, 75–86.
- Jun, Y.B.; Kim, S.J.; Smarandache, F. Interval neutrosophic sets with applications in BCK/BCI-algebra. *Axioms* **2018**, *7*, 23. [[CrossRef](#)]
- Jun, Y.B.; Smarandache, F.; Bordbar, H. Neutrosophic  $\mathcal{N}$ -structures applied to BCK/BCI-algebras. *Information* **2017**, *8*, 128. [[CrossRef](#)]
- Jun, Y.B.; Smarandache, F.; Song, S.Z.; Khan, M. Neutrosophic positive implicative  $\mathcal{N}$ -ideals in BCK/BCI-algebras. *Axioms* **2018**, *7*, 3. [[CrossRef](#)]
- Khan, M.; Anis, S.; Smarandache, F.; Jun, Y.B. Neutrosophic  $\mathcal{N}$ -structures and their applications in semigroups. *Ann. Fuzzy Math. Inform.* **2017**, *14*, 583–598.
- Meng, J.; Jun, Y.B. *BCK-Algebras*; Kyung Moon Sa Co.: Seoul, Korea, 1994.
- Öztürk, M.A.; Jun, Y.B. Neutrosophic ideals in BCK/BCI-algebras based on neutrosophic points. *J. Int. Math. Virtual Inst.* **2018**, *8*, 1–17.
- Saeid, A.B.; Jun, Y.B. Neutrosophic subalgebras of BCK/BCI-algebras based on neutrosophic points. *Ann. Fuzzy Math. Inform.* **2017**, *14*, 87–97.
- Song, S.Z.; Smarandache, F.; Jun, Y.B. Neutrosophic commutative  $\mathcal{N}$ -ideals in BCK-algebras. *Information* **2017**, *8*, 130. [[CrossRef](#)]
- Zhang, X.H.; Bo, C.X.; Smarandache, F.; Park, C. New operations of totally dependent- neutrosophic sets and totally dependent-neutrosophic soft sets. *Symmetry* **2018**, *10*, 187. [[CrossRef](#)]
- Zhang, X.H.; Smarandache, F.; Liang, X.L. Neutrosophic duplet semi-group and cancellable neutrosophic triplet groups. *Symmetry* **2017**, *9*, 275. [[CrossRef](#)]
- Huang, Y.S. *BCI-Algebra*; Science Press: Beijing, China, 2006.
- Smarandache, F. *A Unifying Field in Logics. Neutrosophy: Neutrosophic Probability, Set and Logic*; American Research Press: Rehoboth, NM, USA, 1999.
- Song, S.Z.; Khan, M.; Smarandache, F.; Jun, Y.B. A novel extension of neutrosophic sets and its application in BCK/BCI-algebras. In *New Trends in Neutrosophic Theory and Applications (Volume II)*; Pons Editions; EU: Brussels, Belgium, 2018; pp. 308–326.

16. Zhang, X.H.; Park, C.; Wu, S.P. Soft set theoretical approach to pseudo-BCI algebras. *J. Intell. Fuzzy Syst.* **2018**, *34*, 559–568. [[CrossRef](#)]
17. Zhang, X.H.; Bo, C.X.; Smarandache, F.; Dai, J.H. New inclusion relation of neutrosophic sets with applications and related lattice structure. *Int. J. Mach. Learn. Cybern.* **2018**. [[CrossRef](#)]



© 2018 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).