

# BASES OF SOLUTIONS FOR LINEAR CONGRUENCES

Florentin Smarandache  
University of New Mexico  
200 College Road  
Gallup, NM 87301, USA  
E-mail: smarand@unm.edu

In this article we establish some properties regarding the solutions of a linear congruence, bases of solutions of a linear congruence, and the finding of other solutions starting from these bases.

This article is a continuation of my article “On linear congruences”.

## §1. Introductory Notions

**Definition 1.** (linear congruence)

We call linear congruence with  $n$  unknowns a congruence of the following form:

$$a_1x_1 + \dots + a_nx_n \equiv b \pmod{m} \quad (1)$$

where  $a_1, \dots, a_n, m \in \mathbb{Z}$ ,  $n \geq 1$ , and  $x_i, i = \overline{1, n}$ , are the unknowns.

The following theorems are known:

**Theorem 1.** The linear congruence (1) has solutions if and only if  $(a_1, \dots, a_n, m, b) | b$ .

**Theorem 2.** If the linear congruence (1) has solutions, then:  $|d| \cdot |m|^{n-1}$  is its number of distinct solutions. (See the article “On the linear congruences”).

**Definition 2.** Two solutions  $X = (x_1, \dots, x_n)$  and  $Y = (y_1, \dots, y_n)$  of the linear congruence (1) are distinct (different) if  $\exists i \in \overline{1, n}$  such that  $x_i \not\equiv y_i \pmod{m}$ .

## §2. Definitions and proprieties of congruences

We'll present some arithmetic properties, which will be used later.

**Lemma 1.** If  $a_1, \dots, a_n \in \mathbb{Z}$ ,  $m \in \mathbb{Z}$ , then:

$$\frac{(a_1, \dots, a_n, m) \cdot m^{n-1}}{(a_1, m) \cdot \dots \cdot (a_n, m)} \in \mathbb{Z}$$

The proof is done using complete induction for  $n \in \mathbb{N}^*$ .

When  $n = 1$  it is evident.

Considering that it is true for values smaller or equal to  $n$ , let's proof that it is true for  $n + 1$ .

Let's note  $x = (a_1, \dots, a_n)$ . Then:

$(a_1, \dots, a_n, a_{n+1}, m) \cdot m^n = [(x, a_{n+1}, m) \cdot m^{2-1}] \cdot m^{n-1}$ , which, in accordance to the induction hypothesis, is divisible by:  
 $[(x, m) \cdot (a_{n+1}, m)] \cdot m^{n-1} = [(a_1, \dots, a_n, m) \cdot (a_{n+1}, m)] \cdot m^{n-1} = [(a_1, \dots, a_n, m) \cdot m^{n-1}] \cdot (a_{n+1}, m)$ ,  
 which is divisible, also in accordance with the induction hypothesis, by  
 $[(a_1, m) \cdot \dots \cdot (a_n, m)] \cdot (a_{n+1}, m) = (a_1, m) \cdot \dots \cdot (a_n, m) \cdot (a_{n+1}, m)$ .

**Theorem 3.** If  $X^0$  constitutes a (particular) solution of the linear congruence (1), and  $p = \prod_{i=1}^n (a_i, m)$ , then:

$$X_i \equiv x_i^0 + \frac{m}{(a_i, m)} t_i, \quad 0 \leq t_i < (a_i, m), \quad t_i \in \mathbb{N} \quad (*)$$

( $i$  taking values from 1 to  $n$ ) constitute  $p$  distinct solutions of (1).

*Proof:*

Because the module of the congruence ( $m$ ) is sub-understood, we omitted it, and we will continue to omit it.

$$\sum_{i=1}^n a_i x_i = \sum_{i=1}^n a_i x_i^0 + \sum_{i=1}^n \frac{a_i m}{(a_i, m)} t_i \equiv b + 0, \text{ therefore there are solutions. Let's show}$$

that they are also distinct.

$$x_i^0 + \frac{m}{(a_i, m)} \alpha \not\equiv x_i^0 + \frac{m}{(a_i, m)} \beta, \quad \text{for } \alpha, \beta \in \mathbb{N}, \alpha \neq \beta, \text{ and } 0 \leq \alpha, \beta < (a_i, m),$$

because the set:

$$\left\{ \frac{m}{(a_i, m)} t_i \mid 0 \leq t_i < (a_i, m), t_i \in \mathbb{N} \right\} \subseteq \{0, 1, \dots, n-1\}, \text{ which constitutes a complete}$$

system of residues modulo  $m$ , and  $\frac{m}{(a_i, m)} \alpha \neq \frac{m}{(a_i, m)} \beta$ , for  $\alpha$  and  $\beta$  previously defined.

Therefore the theorem is proved.

\*

\* \*

One considers the  $Z$ -module  $A$  generated by the vectors  $V_i$ , where

$$V_i^* = \left( \underbrace{0, \dots, 0}_{i-1 \text{ times}}, \frac{m}{(a_i, m)}, \underbrace{0, \dots, 0}_{n-i \text{ times}} \right), \quad i = \overline{1, n}, \text{ from } \mathbb{Z}^n. \text{ The module } A \text{ has the rank } n, (n \geq 1).$$

We could note it  $A = \{v_1, \dots, v_n\}$ .

We'll introduce a few new terms.

**Definition 3.** Two solutions (vectors solution)  $X$  and  $Y$  of congruence (1) are called independent if  $X - Y \notin A$ . Otherwise, they are called dependent solutions.

**Remark 1.** In other words, if  $X$  is a solution of the congruence (1), then the solution  $Y$  of the same congruence is independent of  $X$ , if it was not obtained from  $X$  by applying the formula (\*) for certain values of the parameters  $t_1, \dots, t_n$ .

**Definition 4.** The solutions  $X^1, \dots, X^n$  are called **independent (all together)** if they are independent two by two.

Otherwise, they are called **dependent solutions (all together)**.

**Definition 5.** The solutions  $X^1, \dots, X^n$  of the congruence (1) constitute a base for this congruence, if  $X^1, \dots, X^n$  are independent amongst them, and with their help one obtains all (distinct) solutions of the congruence with the procedure (\*) using the parameters  $t_1, \dots, t_n$ .

**Some proprieties of the linear congruences solutions:**

- 1) If the solution  $X^1$  is independent with the solution  $X^2$  then  $X^2$  is independent with  $X^1$  (the commutative property of the relation “independent”).
- 2)  $X^1$  is not independent with  $X^1$ .
- 3) If  $X^1$  is independent with  $X^2$ ,  $X^2$  is independent with  $X^3$ , it does not imply that  $X^1$  is independent with  $X^3$  (the relation is not transitive).
- 4) If  $X$  is independent with  $Y$ , then  $X$  is independent with  $Y$ .

Indeed, if  $Y$  is dependent with  $Y$ , then  $X - Y = \underbrace{(X - Y)}_{\notin A} + \underbrace{(Y - Y_1)}_{\in A} = Z$ .

If  $Z \in A$ , it results that  $(X - Y) = Z - (Y - Y_1) \in A$  because  $A$  is a  $Z$ - module. Absurdity.

\*

\* \*

**Theorem 4.** Let's note  $P_1 = (a_1, \dots, a_n, m) \cdot |m|^{n-1}$  and  $P_2 = (a_1, m) \cdot \dots \cdot (a_n, m)$  then the linear congruence (1) has the base formed of:  $\frac{P_1}{P_2}$  solutions.

*Proof:*

$P_1 > 0$  and  $P_2 > 0$ , from Lemma 1 we have  $\frac{P_1}{P_2} \in \mathbb{N}^*$ , therefore the theorem has

sense (we consider LCD as a positive number).

$P_1$  represents the number of distinct solutions (in total) of congruence (1), in accordance to theorem 2.

$P_2$  represents the number of distinct solutions obtained for congruence (1) by applying the procedure (\*) (allocating to parameters  $t_1, \dots, t_n$  all possible values) to a single particular solution.

Therefore we must apply the procedure (\*)  $\frac{P_1}{P_2}$  times to obtain all solutions of the congruence, that is, it is necessary of exact  $\frac{P_1}{P_2}$  independent particular solutions of the congruence. That is, the base has  $\frac{P_1}{P_2}$  solutions.

**Remark 2.** Any base of solutions (for the same linear congruence) has the same number of vectors.

### §3. Method of solving the linear congruences

In this paragraph we will utilize the results obtained in the precedent paragraphs.

Let's consider the linear congruence (1) with  $(a_1, \dots, a_n, m) = d \mid b, m \neq 0$ .

- we determine the number of distinct solutions of the congruence:

$$P_1 = |d| \cdot |m|^{n-1};$$

- we determine the number of solutions from the base:  $S = \frac{P_1}{\prod_{i=1}^n (a_i, m)}$ ;

- we construct the  $Z$ -module  $A = \{V_1, \dots, V_n\}$ , where

$$V_i^t = \left( \underbrace{0, \dots, 0}_{i-1 \text{ times}}, \frac{m}{(a_i, m)}, \underbrace{0, \dots, 0}_{n-i \text{ times}} \right), \quad i = \overline{1, n};$$

- we search to find  $s$  independent (particular) solutions of the congruence;

- we apply the procedure (\*) as follows:

if  $X^j, j = \overline{1, s}$ , are the  $s$  independent solutions from the base, it results that

$$X^{j(t_1, \dots, t_n)} = \left( x_i^j + \frac{m}{(a_i, m)} t_i \right), \quad i = \overline{1, n}, \quad (*)$$

are all  $P_1$  solutions of the linear congruence (1),

$$j = \overline{1, s}, \quad t_1 \times \dots \times t_n \in \{0, 1, 2, \dots, d_1 - 1\} \times \dots \times \{0, 1, 2, \dots, d_n - 1\},$$

where  $d_i = |(a_i, m)|, i = \overline{1, n}$ .

**Remark 3.** The correctness of this method results from the anterior paragraphs.

**Application.** Let's consider the linear non-homogeneous congruence  $2x - 6y \equiv 2 \pmod{12}$ . It has  $(2, 6, 12) \cdot 12^{2-1} = 24$  distinct solutions. Its base will have  $24 : [(2, 12) \cdot (6, 12)] = 2$  solutions.

$$V_1^t = (6, 0), \quad V_2^t = (0, 2) \text{ and } A = \{V_1, V_2\} = \{(6t_1, 2t_2)^t \mid t_1, t_2 \in \mathbb{Z}\}.$$

The solutions  $x \equiv 7 \pmod{12}$  and  $y \equiv 4 \pmod{12}$ ,  $x \equiv 1$  and  $y \equiv 0$  are dependent because:

$$\begin{pmatrix} 7 \\ 0 \end{pmatrix} - \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 6 \\ 4 \end{pmatrix} = 1 \begin{pmatrix} 6 \\ 0 \end{pmatrix} + 2 \begin{pmatrix} 0 \\ 2 \end{pmatrix} \in A.$$

But  $\begin{pmatrix} 4 \\ 1 \end{pmatrix}$  is independent with  $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$  because  $\begin{pmatrix} 4 \\ 1 \end{pmatrix} - \begin{pmatrix} 0 \\ 1 \end{pmatrix} \notin A$ .

Therefore, the 24 solutions of the congruence can be obtained from:

$$\begin{cases} x \equiv 1 + 6t_1, & 0 \leq t_1 < 2, & t_1 \in \mathbb{N} \\ y \equiv 0 + 2t_2, & 0 \leq t_2 < 6, & t_2 \in \mathbb{N} \end{cases}$$

and

$$\begin{cases} x \equiv 4 + 6t_1, & 0 \leq t_1 < 2, & t_1 \in \mathbb{N} \\ y \equiv 1 + 2t_2, & 0 \leq t_2 < 6, & t_2 \in \mathbb{N} \end{cases}$$

by the parameterization  $(t_1, t_2) \in \{0, 1\} \times \{0, 1, 2, 3, 4, 5\}$ .

$$\begin{cases} x \equiv 1 + 6t_1 \\ y \equiv 0 + 2t_2 \end{cases} \Rightarrow \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 2 \end{pmatrix}, \begin{pmatrix} 1 \\ 4 \end{pmatrix}, \begin{pmatrix} 1 \\ 6 \end{pmatrix}, \begin{pmatrix} 1 \\ 8 \end{pmatrix}, \begin{pmatrix} 1 \\ 10 \end{pmatrix}, \begin{pmatrix} 7 \\ 0 \end{pmatrix}, \begin{pmatrix} 7 \\ 2 \end{pmatrix}, \begin{pmatrix} 7 \\ 4 \end{pmatrix}, \begin{pmatrix} 7 \\ 6 \end{pmatrix}, \begin{pmatrix} 7 \\ 8 \end{pmatrix}, \begin{pmatrix} 7 \\ 10 \end{pmatrix}.$$

$$\begin{cases} x \equiv 4 + 6t_1 \\ y \equiv 1 + 2t_2 \end{cases} \Rightarrow \begin{pmatrix} 4 \\ 1 \end{pmatrix}, \begin{pmatrix} 4 \\ 3 \end{pmatrix}, \begin{pmatrix} 4 \\ 5 \end{pmatrix}, \begin{pmatrix} 4 \\ 7 \end{pmatrix}, \begin{pmatrix} 4 \\ 9 \end{pmatrix}, \begin{pmatrix} 4 \\ 11 \end{pmatrix}, \begin{pmatrix} 10 \\ 1 \end{pmatrix}, \begin{pmatrix} 10 \\ 3 \end{pmatrix}, \begin{pmatrix} 10 \\ 5 \end{pmatrix}, \begin{pmatrix} 10 \\ 7 \end{pmatrix}, \begin{pmatrix} 10 \\ 9 \end{pmatrix}, \begin{pmatrix} 10 \\ 11 \end{pmatrix};$$

which constitute all 24 distinct solutions of the given congruence;  $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$  means:

$x \equiv 1(\text{mod}12)$  and  $y \equiv 0(\text{mod}12)$ ; etc.

## REFERENCES

- [1] C. P. Popovici – “Teoria numerelor”, Editura didactică și pedagogică, București, 1973.
- [2] F. Gh. Smarandache – “Rezolvarea ecuațiilor și a sistemelor de ecuații liniare în numere întregi”, Lucrare de licență, Universitatea din Craiova, 1979.

[Published in “Bulet. Univ. Brașov”, series C, vol. XXII, pp. 25-31, 1980; and in “Bulet. Șt. și Tehn. al Institut. Polit. “Traian Vuia”, Timișoara”, fascicolul 2, tomul 26 (40), pp. 13-6; MR: 83c: 10024]