

GENERALIZATIONS OF DEGARGUES THEOREM*

Florentin Smarandache
University of New Mexico
200 College Road
Gallup, NM 87301, USA
E-mail: smarand@unm.edu

Let's consider the points A_1, \dots, A_n situated on the same plane, and B_1, \dots, B_n situated on another plane, such that the lines $A_i B_i$ are concurrent. Let's prove that if the lines $A_i A_j$ and $B_i B_j$ are concurrent, then their intersecting points are collinear.

Solution. Let α be the plane that contains the points A_1, \dots, A_n (in the case in which the points are non-collinear α is unique), and analogously, let $\beta = P(B_1, \dots, B_n)$, and consider $\alpha \cap \beta = d$.

Because the lines $A_i A_j$ and $B_i B_j$ are concurrent, $A_i A_j \subset \alpha$, and $B_i B_j \subset \beta$, therefore their intersection belongs to line d .

Remark 1.

For $n = 3$ and A_1, A_2, A_3 non-collinear, B_1, B_2, B_3 non-collinear, and $A_i \neq B_j$ we obtain Desargues theorem.

Remark 2.

An extension of this generalization is: If we consider A_1, \dots, A_n situated in a plane, and B_1, \dots, B_m situated on another plane, prove that if $A_i A_j$ and $B_k B_r$ are concurrent, then their intersection points are concurrent.

Remark 3.

For $n = m$, and $A_i B_i$ concurrent lines, we obtain the first generalization.

Remark 4.

If in addition we also have $n = m = 3$ along with the previous conditions, we obtain the Desargues theorem.

* Gamma, Anul X, nr. 1-2, Oct. 1987.