



Regular Single Valued Neutrosophic Hypergraphs

Muhammad Aslam Malik¹, Ali Hassan², Said Broumi³ and F. Smarandache⁴

¹Department of mathematics, University of Punjab, Lahore (Pakistan), E-mail: aslam@math.pu.edu.pk, malikpu@yahoo.com.

²Department of mathematics, University of Punjab, Lahore (Pakistan), E-mail: alihassan.iui.math@gmail.com.

³Laboratory of Information Processing, Faculty of Science Ben M'Sik, University Hassan II, B.P 7955, Sidi Othman, Casablanca, Morocco.

⁴University of New Mexico, Mathematics & Science Department, 705 Gurley Ave., Gallup, NM 87301, USA. E-mail: fsmarandache@gmail.com

Abstract. In this paper, we define the regular and totally regular single valued neutrosophic hypergraphs, and discuss the order and size along with properties of regular

and totally regular single valued neutrosophic hypergraphs. We also extend work on completeness of single valued neutrosophic hypergraphs.

Keywords: Single valued neutrosophic hypergraphs, regular single valued neutrosophic hypergraphs and totally regular single valued neutrosophic hypergraphs.

1 Introduction

The notion of neutrosophic sets (NSs) was proposed by Smarandache [8] as a generalization of the fuzzy sets [14], intuitionistic fuzzy sets [12], interval valued fuzzy set [11] and interval-valued intuitionistic fuzzy sets [13] theories. The neutrosophic set is a powerful mathematical tool for dealing with incomplete, indeterminate and inconsistent information in real world. The neutrosophic sets are characterized by a truth-membership function (t), an indeterminacy-membership function (i) and a falsity membership function (f) independently, which are within the real standard or nonstandard unit interval $]0, 1^+[$. In order to conveniently use NS in real life applications, Wang et al. [9] introduced the concept of the single-valued neutrosophic set (SVNS), a subclass of the neutrosophic sets. The same authors [10] introduced the concept of the interval valued neutrosophic set (IVNS), which is more precise and flexible than the single valued neutrosophic set. The IVNS is a generalization of the single valued neutrosophic set, in which the three membership functions are independent and their value belong to the unit interval $[0, 1]$. More works on single valued neutrosophic sets, interval valued neutrosophic sets and their applications can be found on <http://fs.gallup.unm.edu/NSS/>.

Hypergraph is a graph in which an edge can connect more than two vertices, hypergraphs can be applied to analyse

architecture structures and to represent system partitions, Mordesen J.N and P.S Nasir gave the definitions for fuzzy hypergraphs. Parvathy. R and M. G. Karunambigai's paper introduced the concepts of Intuitionistic fuzzy hypergraphs and analyse its components, Nagoor Gani. A and Sajith Begum. S defined degree, order and size in intuitionistic fuzzy graphs and extend the properties. Nagoor Gani. A and Latha. R introduced irregular fuzzy graphs and discussed some of its properties.

Regular intuitionistic fuzzy hypergraphs and totally regular intuitionistic fuzzy hypergraphs are introduced by Pra-deepa. I and Vimala. S in [0]. In this paper we extend regularity and totally regularity on single valued neutrosophic hypergraphs.

2 Preliminaries

Definition 2.1 Let X be a space of points (objects) with generic elements in X denoted by x . A single valued neutrosophic set A (SVNS A) is characterized by truth membership function $T_A(x)$, indeterminacy membership function $I_A(x)$ and a falsity membership function $F_A(x)$. For each point $x \in X$; $T_A(x), I_A(x), F_A(x) \in [0, 1]$.

Definition 2.2 Let A be a SVNS on X then support of A is denoted and defined by, $Supp(A) = \{x : x \in X, T_A(x) > 0, I_A(x) > 0, F_A(x) > 0\}$.

Definition 2.3 A hypergraph is an ordered pair $H = (X, E)$, where

- (1) $X = \{x_1, x_2, \dots, x_n\}$ be a finite set of vertices.
- (2) $E = \{E_1, E_2, \dots, E_m\}$ be a family of subsets of X .
- (3) E_j for $j = 1, 2, 3, \dots, m$ and $\cup_j(E_j) = X$.

The set X is called set of vertices and E is the set of edges(or hyper edges).

Definition 2.4 The single valued neutrosophic hypergraph is an ordered pair $H = (X, E)$, where

- (1) $X = \{x_1, x_2, \dots, x_n\}$ be a finite set of vertices.
- (2) $E = \{E_1, E_2, \dots, E_m\}$ be a family of SVN-edges of X .
- (3) $E_j \neq O = (0, 0, 0)$ for $j = 1, 2, 3, \dots, m$ and $\cup_j Supp(E_j) = X$.

The set X is called set of vertices and E is the set of SVN-edges(or SVN-hyperedges).

Proposition 2.5 The single valued neutrosophic hypergraph is the generalization of fuzzy hypergraphs and intuitionistic fuzzy hypergraphs.

3 Regular and totally regular SVNHG

Definition 3.1 The open neighbourhood of a vertex x in single valued neutrosophic hypergraphs (SVNHGs) is the set of adjacent vertices of x , excluding that vertex and is denoted by $N(x)$.

Definition 3.2 The closed neighbourhood of a vertex x in single valued neutrosophic hypergraphs (SVNHGs) is the set of adjacent vertices of x , including that vertex and is denoted by $N[x]$.

Example 3.3 Consider a single valued neutrosophic hypergraphs $H = (X, E)$, where $X = \{a, b, c, d, e\}$ and $E = \{P, Q, R, S\}$, which are defined by

$$P = \{(a, .1, .2, .3), (b, .4, .5, .6)\}$$

$$Q = \{(c, .1, .2, .3), (d, .4, .5, .6), (e, .7, .8, .9)\}$$

$$R = \{(b, .1, .2, .3), (c, .4, .5, .6)\}$$

$$S = \{(a, .1, .2, .3), (d, .4, .5, .6)\}$$

Then the open neighbourhood of a vertex a contain b and d . The closed neighbourhood of a vertex b contain b, a and c .

Definition 3.4 Let $H = (X, E)$ be a SVNHG, the open neighbourhood degree of a vertex x , which is denoted and defined by $deg(x) = (deg_T(x), deg_I(x), deg_F(x))$, where

$$deg_T(x) = \sum_{x \in X} T_E(x)$$

$$deg_I(x) = \sum_{x \in X} I_E(x)$$

$$deg_F(x) = \sum_{x \in X} F_E(x)$$

Example 3.5 Consider a single valued neutrosophic hypergraphs $H = (X, E)$ where, $X = \{a, b, c, d, e\}$ and $E = \{P, Q, R, S\}$, which are defined by

$$P = \{(a, .1, .2, .3), (b, .4, .5, .6)\}$$

$$Q = \{(c, .1, .2, .3), (d, .4, .5, .6), (e, .7, .8, .9)\}$$

$$R = \{(b, .1, .2, .3), (c, .4, .5, .6)\}$$

$$S = \{(a, .1, .2, .3), (d, .4, .5, .6)\}$$

Then the open neighbourhood of a vertex a is b and d , and therefore the open neighbourhood degree degree of a vertex a is $(.8, 1, 1.2)$.

Definition 3.6 Let $H = (X, E)$ be a SVNHG, the closed neighbourhood degree of a vertex x , which is denoted and defined by

$$deg[x] = (deg_T[x], deg_I[x], deg_F[x])$$

where

$$deg_T[x] = deg_T(x) + T_E(x)$$

$$deg_I[x] = deg_I(x) + I_E(x)$$

$$deg_F[x] = deg_F(x) + F_E(x)$$

Example 3.7 Consider a single valued neutrosophic hypergraphs $H = (X, E)$, where $X = \{a, b, c, d, e\}$ and $E = \{P, Q, R, S\}$, which is defined by

$$P = \{(a, .1, .2, .3), (b, .4, .5, .6)\}$$

$$Q = \{(c, .1, .2, .3), (d, .4, .5, .6), (e, .7, .8, .9)\}$$

$$R = \{(b, .1, .2, .3), (c, .4, .5, .6)\}$$

$$S = \{(a, .1, .2, .3), (d, .4, .5, .6)\}$$

The closed neighbourhood of a vertex b contain b, a and c , hence the closed neighbourhood degree of a vertex a is $(.9, .1.2, 1.5)$.

Definition 3.8 Let $H = (X, E)$ be a SVNHG, then H is said to be an n -regular SVNHG if all the vertices have the same open neighbourhood degree $n = (n_1, n_2, n_3)$.

Definition 3.9 Let $H = (X, E)$ be a SVNHG, then H is said to be an m -totally regular SVNHG if all the vertices have the same closed neighbourhood degree $m = (m_1, m_2, m_3)$.

Proposition 3.10 A regular SVNHG is the generalization of regular fuzzy hypergraphs and regular intuitionistic fuzzy hypergraphs.

Proposition 3.11 A totally regular SVNHG is the generalization of totally regular fuzzy hypergraphs and totally regular intuitionistic fuzzy hypergraphs.

Example 3.12 Consider a single valued neutrosophic hypergraphs $H = (X, E)$, where $X = \{a, b, c, d\}$ and $E = \{P, Q, R, S\}$, which are defined by

$$P = \{(a, .8, .2, .3), (b, .8, .2, .3)\}$$

$$Q = \{(b, .8, .2, .3), (c, .8, .2, .3)\}$$

$$R = \{(c, .8, .2, .3), (d, .8, .2, .3)\}$$

$$S = \{(d, .8, .2, .3), (a, .8, .2, .3)\}$$

Here the open neighbourhood degree of every vertex is $(1.6, .4, .6)$, hence H is regular SVNHG and closed neighbourhood degree of every vertex is $(2.4, .6, .9)$. Hence H is both regular and totally regular SVNHG.

Theorem 3.13 Let $H = (X, E)$ be a SVNHG which is both regular and totally regular SVNHG then E is constant.

Proof: Suppose H is an n -regular and an m -totally regular SVNHG. Then

$$deg(x) = n = (n_1, n_2, n_3)$$

$$deg[x] = m = (m_1, m_2, m_3)$$

for all $x \in E_i$. Consider the $deg[x] = m$, hence by definition $deg(x) + E_i(x) = m$ this implies that $E_i(x) = m - n$ for all x in E_i . Therefore E is constant.

Remark 3.14 The converse of above theorem need not to be true in general.

Example 3.15 Consider a SVNHG $H = (X, E)$, where $X = \{a,$

$b, c, d\}$ and $E = \{P, Q, R, S\}$, which is defined by

$$P = \{(a, .8, .2, .3), (b, .8, .2, .3)\}$$

$$Q = \{(b, .8, .2, .3), (d, .8, .2, .3)\}$$

$$R = \{(c, .8, .2, .3), (d, .8, .2, .3)\}$$

$$S = \{(d, .8, .2, .3), (a, .8, .2, .3)\}$$

Here E is constant but $deg(a) = (1.6, .4, .6)$ and $deg(d) = (2.4, .6, .9)$. Therefore $deg(a)$ and $deg(d)$ are not equal, hence H is not regular SVNHG. Also $deg[a] = (2.4, .6, .9)$ and $deg[d] = (3.2, .8, 1.2)$, thus $deg[a]$ and $deg[d]$ are not equal, hence H is not totally regular SVNHG, we conclude that H is neither regular and nor totally regular SVNHG.

Theorem 3.16 Let $H = (X, E)$ be a SVNHG, then E is constant on X if and only if following are equivalent

(1) H is regular SVNHG.

(2) H is totally regular SVNHG.

Proof: Suppose $H = (X, E)$ be a SVNHG and E is constant in H , then $E_i(x) = c = (c_1, c_2, c_3)$ for all $x \in E_i$. Suppose H is an n -regular SVNHG, then $deg(x) = n = (n_1, n_2, n_3)$ for all $x \in E_i$. Consider $deg[x] = deg(x) + E_i(x) = n + c$ for all $x \in E_i$. Hence H is totally regular SVNHG. Next suppose that H is an m -totally regular SVNHG, then $deg[x] = m = (m_1, m_2, m_3)$ for all $x \in E_i$, that is $deg(x) + E_i(x) = m$ for all $x \in E_i$. This implies that $deg(x) = m - c$ for all $x \in E_i$. Thus H is regular SVNHG.

Conversely: Suppose contrary E is not constant, that is $E_i(x)$ and $E_i(y)$ not equal for some x and y in X . Let $H = (X, E)$ be an n -regular SVNHG, then $deg(x) = n = (n_1, n_2, n_3)$ for all $x \in E_i$. Consider

$$deg[x] = deg(x) + E_i(x) = n + E_i(x)$$

$$deg[y] = deg(y) + E_i(y) = n + E_i(y)$$

since $E_i(x)$ and $E_i(y)$ are not equal for some x and y in X , hence $deg[x]$ and $deg[y]$ are not equal, thus H is not totally regular SVNHG, which contradict to our assumption. Next let H be totally regular SVNHG, then $deg[x] = deg[y]$, that is

$$deg(x) + E_i(x) = deg(y) + E_i(y)$$

$$deg(x) - deg(y) = E_i(y) - E_i(x)$$

since RHS of above equation is nonzero, hence LHS of above equation is also nonzero, thus $deg(x)$ and $deg(y)$ are not equals, so H is not regular SVNHG, which is again contradict to our assumption, thus our supposition was

wrong, hence E must be constant, this completes the proof.

Definition 3.17 Let $H = (X, E)$ be a regular SVNHG, then the order of SVNHG H , which is denoted and defined by $O(H) = (p, q, r)$, where

$$p = \sum_{x \in X} T_{E_i}(x)$$

$$q = \sum_{x \in X} I_{E_i}(x)$$

$$r = \sum_{x \in X} F_{E_i}(x)$$

for every $x \in X$ and the size of regular SVNHG, which is denoted and defined by $S(H) = \sum_{i=1}^n (S_{E_i})$, where $S(E_i) = (a, b, c)$, which is defined by

$$a = \sum_{x \in E_i} T_{E_i}(x)$$

$$b = \sum_{x \in E_i} I_{E_i}(x)$$

$$c = \sum_{x \in E_i} F_{E_i}(x).$$

Example 3.18 Consider the SVNHG $H = (X, E)$, where $X = \{a, b, c, d\}$ and $E = \{P, Q, R, S\}$, which is defined by

$$P = \{(a, .8, .2, .3), (b, .8, .2, .3)\}$$

$$Q = \{(b, .8, .2, .3), (c, .8, .2, .3)\}$$

$$R = \{(c, .8, .2, .3), (d, .8, .2, .3)\}$$

$$S = \{(d, .8, .2, .3), (a, .8, .2, .3)\}$$

Here the order and the size of H are given (3.2, .8, 1.2) and (6.4, 1.6, 2.4), respectively.

Proposition 3.19 The size of an n -regular SVNHG $H = (X, E)$ is $nk/2$, where $|X|=k$.

Proposition 3.20 Let $H = (X, E)$ be an m -totally regular SVNHG, then $2S(H) + O(H) = mk$, where $|X|=k$.

Corollary 3.21 Let $H = (X, E)$ be an n -regular and an m -totally regular SVNHG, then $O(H) = k(m - n)$, where $|X|=k$.

Proposition 3.22 The dual of an n -regular and an m -totally regular SVNHG $H = (X, E)$ is again an n -regular and an m -totally regular SVNHG.

Definition 3.23 The SVNHG is said to be complete SVNHG if for every x in X , $N(x) = \{x: x \text{ in } X - \{x\}\}$, that is $N(x)$ contains all remaining vertices of X except x .

Example 3.24 Consider a single valued neutrosophic hypergraphs $H = (X, E)$, where $X = \{a, b, c, d\}$ and $E = \{P, Q, R\}$, which is defined by

$$P = \{(a, .4, .6, .3), (c, .8, .2, .3)\}$$

$$Q = \{(a, .8, .8, .3), (b, .8, .2, .1), (d, .8, .2, .1)\}$$

$$R = \{(c, .4, .9, .9), (d, .7, .2, .1), (b, .4, .2, .1)\}$$

Here $N(a) = \{b, c, d\}$, $N(b) = \{a, c, d\}$, $N(c) = \{a, b, d\}$, $N(d) = \{a, b, c\}$. Hence H is complete SVNHG.

Remark 3.25 In a complete SVNHG $H = (X, E)$ the cardinality of $N(x)$ is same for every vertex.

Theorem 3.26 Every complete SVNHG $H = (X, E)$ is both regular and totally regular if is constant in H .

Proof: Let $H = (X, E)$ be a complete SVNHG, suppose E is constant in H , so $\forall x \in E_i, E_i(x) = c = (c_1, c_2, c_3)$, since SVNHG is complete, then by definition for every vertex x in X , $N(x) = \{x : x \text{ in } X - \{x\}\}$, the open neighbourhood degree of every vertex is same. Hence $deg(x) = n = (n_1, n_2, n_3)$ for all $x \in E_i$. Hence complete SVNHG is regular SVNHG. Also $deg[x] = deg(x) + E_i(x) = n + c$ for all $x \in E_i$. Thus H is totally regular SVNHG.

Remark 3.27 Every complete SVNHG is totally regular even if E is not constant.

Definition 3.28 The SVNHG is said to be k -uniform if all the hyperedges have same cardinality.

Example 3.29 Consider a SVNHG $H = (X, E)$, where $X = \{a, b, c, d\}$ and $E = \{P, Q, R\}$, which is defined by

$$P = \{(a, .8, .2, .3), (b, .7, .5, .3)\}$$

$$Q = \{(b, .8, .1, .8), (c, .8, .4, .2)\}$$

$$R = \{(c, .8, .1, .4), (d, .8, .9, .5)\}$$

4 Conclusion

Theoretical concepts of graphs and hypergraphs are highly utilized by computer science applications. The SVNHG are more flexible than fuzzy hypergraphs and intuitionistic fuzzy hypergraphs. The concepts of SVNHGs can be applied in various areas of engineering and computer science. In this paper, we defined the concept of regular and totally regular SVNHGs. We plan to extend our research work to regular and totally regular on Bipolar SVNHGs, regular and totally regular on interval valued neutrosophic hypergraphs, irregular and totally irregular on SVNHGs, irregular and totally irregular on bipolar SVNHGs.

References

- [0] I. Pradeepa and S.Vimala, regular and totally regular intuitionistic fuzzy hyper graphs, international journal of mathematics and applications, volume 4 , issue 1-C (2016), 137-142.
- [1] A. V. Devadoss, A. Rajkumar & N. J. P. Praveena. A Study on Miracles through Holy Bible using Neutrosophic Cognitive Maps (NCMS). In: International Journal of Computer Applications, 69(3) (2013).
- [2] A. Nagoor Gani and M. B. Ahamed. Order and Size in Fuzzy Graphs. In: Bulletin of Pure and Applied Sciences, Vol 22E (No.1) (2003) 145-148.
- [3] A. N. Gani. A. and S. Shajitha Begum. Degree, Order and Size in Intuitionistic Fuzzy Graphs. In: Intl. Journal of Algorithms, Computing and Mathematics, (3)3 (2010).
- [4] A. Nagoor Gani and S.R Latha. On Irregular Fuzzy Graphs. In: Applied Mathematical Sciences, Vol. 6, no.11 (2012) 517-523.
- [5] F. Smarandache. Refined Literal Indeterminacy and the Multiplication Law of Sub-Indeterminacies. In: Neutrosophic Sets and Systems, Vol. 9 (2015) 58-63.
- [6] F. Smarandache. Types of Neutrosophic Graphs and Neutrosophic Algebraic Structures together with their Applications in Technology, Seminar, Universitatea Transilvania din Brasov, Facultatea de Design de Produs si Mediu, Brasov, Romania 06 June 2015.
- [7] F. Smarandache. Symbolic Neutrosophic Theory. Brussels: Europanova, 2015, 195 p.
- [8] F. Smarandache. Neutrosophic set - a generalization of the intuitionistic fuzzy set. In: Granular Computing, 2006 IEEE Intl. Conference, (2006) 38 - 42, DOI: 10.1109/GRC.2006.1635754.
- [9] H. Wang, F. Smarandache, Y. Zhang, and R. Sunderraman. Single Valued Neutrosophic Sets. In: Multispace and Multistructure, 4 (2010) 410-413.
- [10] H. Wang, F. Smarandache, Zhang, Y.-Q. and R. Sunderraman. Interval Neutrosophic Sets and Logic: Theory and Applications in Computing. Phoenix: Hexis, 2005.
- [11] I. Turksen. Interval valued fuzzy sets based on normal forms. In: Fuzzy Sets and Systems, vol. 20 (1986) 191-210.
- [12] K. Atanassov. Intuitionistic fuzzy sets. In: Fuzzy Sets and Systems. vol. 20 (1986) 87-96.
- [13] K. Atanassov and G. Gargov. Interval valued intuitionistic fuzzy sets. In: Fuzzy Sets and Systems, vol. 31 (1989) 343-349.
- [14] L. Zadeh. Fuzzy sets. In: Information and Control, 8 (1965) 338-353.
- [15] P. Bhattacharya. Some remarks on fuzzy graphs. In: Pattern Recognition Letters 6 (1987) 297-302.
- [16] R. Parvathi and M. G. Karunambigai. Intuitionistic Fuzzy Graphs. In: Computational Intelligence. In: Theory and applications, International Conference in Germany, Sept 18-20, 2006.
- [17] R. A. Borzooei, H. Rashmanlou. More Results On Vague Graphs, U.P.B. Sci. Bull., Series A, Vol. 78, Issue 1, 2016, 109-122.
- [18] S. Broumi, M. Talea, F. Smarandache, A. Bakali. Single Valued Neutrosophic Graphs: Degree, Order and Size, FUZZ IEEE Conference (2016), 8 page.
- [19] S. Broumi, M. Talea, A. Bakali, F. Smarandache. Single Valued Neutrosophic Graphs. In: Journal of New Theory, no. 10, 68-101 (2016).
- [20] S. Broumi, M. Talea, A. Bakali, F. Smarandache. On Bipolar Single Valued Neutrosophic Graphs. In: Journal of New Theory, no. 11, 84-102 (2016).
- [21] S. Broumi, M. Talea, A. Bakali, F. Smarandache. Interval Valued Neutrosophic Graphs. SISOM Conference (2016), in press.
- [22] S. Broumi, F. Smarandache, M. Talea, A. Bakali. An Introduction to Bipolar Single Valued Neutrosophic Graph Theory. OPTIROB conference, 2016.
- [23] S. Broumi, M. Talea, A. Bakali, F. Smarandache. Operations on Interval Valued Neutrosophic Graphs (2016), submitted.
- [24] S. Broumi, M. Talea, A. Bakali, F. Smarandache. Strong Interval Valued Neutrosophic Graphs, (2016) , submitted.
- [25] S. N. Mishra and A. Pal. Product of Interval Valued Intuitionistic fuzzy graph. In: Annals of Pure and Applied Mathematics, Vol. 5, No.1 (2013) 37-46.
- [26] S. Rahurikar. On Isolated Fuzzy Graph. In: Intl. Journal of Research in Engineering Technology and Management, 3 pages.
- [27] W. B. Vasantha Kandasamy, K. Ilanthenral F. Smarandache. Neutrosophic Graphs: A New Dimension to Graph Theory. Kindle Edition, 2015.

- [28] F. Smarandache, *Neutrosophy / Neutrosophic probability, set, and logic*, American Res. Press, see pages 7-8, 1998, <http://fs.gallup.unm.edu/eBook-neutrosophics6.pdf> (last edition online); reviewed in Zentralblatt fuer Mathematik (Berlin, Germany): <https://zbmath.org/?q=an:01273000>
- [29] Denis Howe, England, *The Free Online Dictionary of Computing*, 1999.

Received: December 01, 2016. Accepted: December 28, 2016